You can contact CarePlus for the most recent list of drugs by calling 1-800-794-5907; TTY: 711. From October 1 - March 31, we are open 7 days a week, 8 a.m. to 8 p.m. From April 1 - September 30, we are open Monday - Friday, 8 a.m. to 8 p.m. You may always leave a voicemail after hours, Saturdays, Sundays, and holidays and we will return your call within one business day. You may also visit www.careplushealthplans.com.

Prior Authorization Criteria

Effective 01/01/2027

abiraterone - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Prostate Cancer: One of the following applies: The member has metastatic castration-resistant prostate cancer (mCRPC) or The member has a diagnosis of castration-sensitive prostate cancer (CSPC).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

abirtega - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Prostate Cancer: One of the following applies: The member has metastatic castration-resistant prostate cancer (mCRPC) or The member has a diagnosis of castration-sensitive prostate cancer (CSPC).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

acitretin - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Member must have a diagnosis of severe cutaneous psoriasis including plaque, guttate, erythrodermic, palmar-plantar, and pustular types AND the member has had previous treatment, contraindication, or intolerance to methotrexate or cyclosporine.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ACTIMMUNE - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Chronic Granulomatous Disease (CGD): The member has chronic granulomatous disease (CGD). The member is using Actimmune to reduce the frequency and severity of infections. Severe Malignant Osteopetrosis: The member has severe malignant osteopetrosis. The member is using Actimmune to delay time to disease progression.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ADALIMUMAB-ADAZ - PEN INJECTOR (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Ankylosing Spondylitis: Diagnosis of active ankylosing spondylitis. Member has tried or cannot use a non-steroidal anti-inflammatory drug (NSAIDs) (e.g. ibuprofen, meloxicam, naproxen). Psoriatic Arthritis: Diagnosis of active psoriatic arthritis. Rheumatoid Arthritis: Diagnosis of moderately to severely active rheumatoid arthritis. Member has tried a single DMARD (e.g. methotrexate, sulfasalazine, hydroxychloroquine, leflunomide) or cannot use any DMARDs. Polyarticular Juvenile Idiopathic Arthritis: Diagnosis of moderately to severely active polyarticular juvenile idiopathic arthritis. Member has tried or cannot use a DMARD (e.g. methotrexate, cyclosporine, leflunomide). Moderate to severe Chronic Plaque Psoriasis: Diagnosis of moderate to severe chronic plaque psoriasis. The member has tried at least one conventional oral systemic treatment (e.g. methotrexate, cyclosporine), or cannot use any conventional oral systemic treatments. Oral Ulcers Associated with Behcets Disease: The member has a diagnosis of Behcets disease AND The member will be using for the treatment of oral ulcers AND has tried or cannot use two of the following: topical corticosteroid therapy (e.g. triamcinolone oral paste), azathioprine, or colchicine.
Age Restriction	Must be 18 years of age for Rheumatoid Arthritis, Moderate to severe Chronic Plaque Psoriasis, Psoriatic Arthritis, Ankylosing Spondylitis, Oral Ulcers Associated with Behcets Disease. Must be two years of age or older for moderately to severely active polyarticular juvenile idiopathic arthritis and Uveitis. Must be 6 years or older for Crohns Disease. Must be 5 years of age or older for Ulcerative Colitis. Must be 12 years of age or older for Hidradenitis Suppurativa.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration

ADALIMUMAB-ADAZ - PEN INJECTOR (ML)

Other Criteria	Ulcerative colitis: Member has a diagnosis of moderate to severely active ulcerative colitis. Crohn's Disease: The member must have moderately to severely active Crohn's disease. Hidradenitis Suppurativa: The member must have a diagnosis of moderate to severe Hidradenitis Suppurativa. Uveitis: The member must have a diagnosis of non-infectious, intermediate, posterior, or pan-uveitis. The member has tried or cannot use one of the following: an intravitreal steroid (e.g. triamcinolone, dexamethasone) OR a systemic corticosteroid (e.g. prednisone, methylprednisolone) OR an anti-metabolite (e.g. methotrexate, mycophenolate) OR a calcineurin inhibitor (e.g. cyclosporine).
Part B Prerequisite	0

ADALIMUMAB-ADBM - PEN INJECTOR KIT (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Ankylosing Spondylitis: Diagnosis of active ankylosing spondylitis. Member has tried or cannot use a non-steroidal anti-inflammatory drug (NSAIDs) (e.g. ibuprofen, meloxicam, naproxen). Psoriatic Arthritis: Diagnosis of active psoriatic arthritis. Rheumatoid Arthritis: Diagnosis of moderately to severely active rheumatoid arthritis. Member has tried a single DMARD (e.g. methotrexate, sulfasalazine, hydroxychloroquine, leflunomide) or cannot use any DMARDs. Polyarticular Juvenile Idiopathic Arthritis: Diagnosis of moderately to severely active polyarticular juvenile idiopathic arthritis. Member has tried or cannot use a DMARD (e.g. methotrexate, cyclosporine, leflunomide). Moderate to severe Chronic Plaque Psoriasis: Diagnosis of moderate to severe chronic plaque psoriasis. The member has tried at least one conventional oral systemic treatment (e.g. methotrexate, cyclosporine), or cannot use any conventional oral systemic treatments. Oral Ulcers Associated with Behcets Disease: The member has a diagnosis of Behcets disease AND The member will be using for the treatment of oral ulcers AND has tried or cannot use two of the following: topical corticosteroid therapy (e.g. triamcinolone oral paste), azathioprine, or colchicine.
Age Restriction	Must be 18 years of age for Rheumatoid Arthritis, Moderate to severe Chronic Plaque Psoriasis, Psoriatic Arthritis, Ankylosing Spondylitis, Oral Ulcers Associated with Behcets Disease. Must be two years of age or older for moderately to severely active polyarticular juvenile idiopathic arthritis and Uveitis. Must be 6 years or older for Crohns Disease. Must be 5 years of age or older for Ulcerative Colitis. Must be 12 years of age or older for Hidradenitis Suppurativa.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration

ADALIMUMAB-ADBM - PEN INJECTOR KIT (EA)

Other Criteria	Ulcerative colitis: Member has a diagnosis of moderate to severely active ulcerative colitis. Crohn's Disease: The member must have moderately to severely active Crohn's disease. Hidradenitis Suppurativa: The member must have a diagnosis of moderate to severe Hidradenitis Suppurativa. Uveitis: The member must have a diagnosis of non-infectious, intermediate, posterior, or pan-uveitis. The member has tried or cannot use one of the following: an intravitreal steroid (e.g. triamcinolone, dexamethasone) OR a systemic corticosteroid (e.g. prednisone, methylprednisolone) OR an anti-metabolite (e.g. methotrexate, mycophenolate) OR a calcineurin inhibitor (e.g. cyclosporine).
Part B Prerequisite	0

ADALIMUMAB-ADBM(CF) PEN CROHNS - PEN INJECTOR KIT (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Ankylosing Spondylitis: Diagnosis of active ankylosing spondylitis. Member has tried or cannot use a non-steroidal anti-inflammatory drug (NSAIDs) (e.g. ibuprofen, meloxicam, naproxen). Psoriatic Arthritis: Diagnosis of active psoriatic arthritis. Rheumatoid Arthritis: Diagnosis of moderately to severely active rheumatoid arthritis. Member has tried a single DMARD (e.g. methotrexate, sulfasalazine, hydroxychloroquine, leflunomide) or cannot use any DMARDs. Polyarticular Juvenile Idiopathic Arthritis: Diagnosis of moderately to severely active polyarticular juvenile idiopathic arthritis. Member has tried or cannot use a DMARD (e.g. methotrexate, cyclosporine, leflunomide). Moderate to severe Chronic Plaque Psoriasis: Diagnosis of moderate to severe chronic plaque psoriasis. The member has tried at least one conventional oral systemic treatment (e.g. methotrexate, cyclosporine), or cannot use any conventional oral systemic treatments. Oral Ulcers Associated with Behcets Disease: The member has a diagnosis of Behcets disease AND The member will be using for the treatment of oral ulcers AND has tried or cannot use two of the following: topical corticosteroid therapy (e.g. triamcinolone oral paste), azathioprine, or colchicine.
Age Restriction	Must be 18 years of age for Rheumatoid Arthritis, Moderate to severe Chronic Plaque Psoriasis, Psoriatic Arthritis, Ankylosing Spondylitis, Oral Ulcers Associated with Behcets Disease. Must be two years of age or older for moderately to severely active polyarticular juvenile idiopathic arthritis and Uveitis. Must be 6 years or older for Crohns Disease. Must be 5 years of age or older for Ulcerative Colitis. Must be 12 years of age or older for Hidradenitis Suppurativa.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration

ADALIMUMAB-ADBM(CF) PEN CROHNS - PEN INJECTOR KIT (EA)

Other Criteria	Ulcerative colitis: Member has a diagnosis of moderate to severely active ulcerative colitis. Crohn's Disease: The member must have moderately to severely active Crohn's disease. Hidradenitis Suppurativa: The member must have a diagnosis of moderate to severe Hidradenitis Suppurativa. Uveitis: The member must have a diagnosis of non-infectious, intermediate, posterior, or pan-uveitis. The member has tried or cannot use one of the following: an intravitreal steroid (e.g. triamcinolone, dexamethasone) OR a systemic corticosteroid (e.g. prednisone, methylprednisolone) OR an anti-metabolite (e.g. methotrexate, mycophenolate) OR a calcineurin inhibitor (e.g. cyclosporine).
Part B Prerequisite	0

ADALIMUMAB-ADBM(CF) PEN PS-UV - PEN INJECTOR KIT (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Ankylosing Spondylitis: Diagnosis of active ankylosing spondylitis. Member has tried or cannot use a non-steroidal anti-inflammatory drug (NSAIDs) (e.g. ibuprofen, meloxicam, naproxen). Psoriatic Arthritis: Diagnosis of active psoriatic arthritis. Rheumatoid Arthritis: Diagnosis of moderately to severely active rheumatoid arthritis. Member has tried a single DMARD (e.g. methotrexate, sulfasalazine, hydroxychloroquine, leflunomide) or cannot use any DMARDs. Polyarticular Juvenile Idiopathic Arthritis: Diagnosis of moderately to severely active polyarticular juvenile idiopathic arthritis. Member has tried or cannot use a DMARD (e.g. methotrexate, cyclosporine, leflunomide). Moderate to severe Chronic Plaque Psoriasis: Diagnosis of moderate to severe chronic plaque psoriasis. The member has tried at least one conventional oral systemic treatment (e.g. methotrexate, cyclosporine), or cannot use any conventional oral systemic treatments. Oral Ulcers Associated with Behcets Disease: The member has a diagnosis of Behcets disease AND The member will be using for the treatment of oral ulcers AND has tried or cannot use two of the following: topical corticosteroid therapy (e.g. triamcinolone oral paste), azathioprine, or colchicine.
Age Restriction	Must be 18 years of age for Rheumatoid Arthritis, Moderate to severe Chronic Plaque Psoriasis, Psoriatic Arthritis, Ankylosing Spondylitis, Oral Ulcers Associated with Behcets Disease. Must be two years of age or older for moderately to severely active polyarticular juvenile idiopathic arthritis and Uveitis. Must be 6 years or older for Crohns Disease. Must be 5 years of age or older for Ulcerative Colitis. Must be 12 years of age or older for Hidradenitis Suppurativa.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration

ADALIMUMAB-ADBM(CF) PEN PS-UV - PEN INJECTOR KIT (EA)

Other Criteria	Ulcerative colitis: Member has a diagnosis of moderate to severely active ulcerative colitis. Crohn's Disease: The member must have moderately to severely active Crohn's disease. Hidradenitis Suppurativa: The member must have a diagnosis of moderate to severe Hidradenitis Suppurativa. Uveitis: The member must have a diagnosis of non-infectious, intermediate, posterior, or pan-uveitis. The member has tried or cannot use one of the following: an intravitreal steroid (e.g. triamcinolone, dexamethasone) OR a systemic corticosteroid (e.g. prednisone, methylprednisolone) OR an anti-metabolite (e.g. methotrexate, mycophenolate) OR a calcineurin inhibitor (e.g. cyclosporine).
Part B Prerequisite	0

ADEMPAS - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Chronic Thromboembolic Pulmonary Hypertension (CTEPH). The member must have a diagnosis of Chronic Thromboembolic Pulmonary Hypertension (CTEPH) (WHO Group 4) AND The member must have CTEPH classified as inoperable or persistent/recurrent after surgical treatment (i.e. pulmonary endarterectomy). Pulmonary Arterial Hypertension (PAH). The member must have a diagnosis of pulmonary arterial hypertension (WHO Group 1) confirmed by right heart catheterization.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ADSTILADRIN - VIAL (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member has experienced disease progression while on Adstiladrin (nadofaragene firadenovec-vncg)
Required Medical Information	Non-muscle invasive bladder cancer (NMIBC): Member has a diagnosis of high-risk Bacillus Calmette-Guerin (BCG)-unresponsive disease, which is defined as persistent or recurrent disease following adequat BCG therapy (adequate BGC is defined as the administration of at least five of six doses of an initial induction course plus either of: at least two of three doses of maintenance therapy or at least two of six doses of a second induction course) AND member has non-muscle invasive bladder cancer (NMIBC) with documented carcinoma in situ (CIS) AND member is ineligible for or has elected not to undergo cystectomy.
Age Restriction	Member is 18 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 months duration
Other Criteria	
Part B Prerequisite	0

afatinib - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Non-small cell lung cancer (NSCLC): The member has a diagnosis of metastatic non-small cell lung cancer (NSCLC) AND the following apply: The member has a documented non-resistant epidermal growth factor receptor (EGFR) mutation (sensitizing EGFR mutation e.g., exon 19 deletion, L861Q, S768I, G719X, L858R) AND The member is using Gilotrif (afatinib) as monotherapy (without concomitant chemotherapy) OR squamous cell histology after disease progression on platinum containing chemotherapy and is using Gilotrif (afatinib) as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Six month duration
Other Criteria	
Part B Prerequisite	0

AKEEGA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on or following PARP inhibitor therapy [e.g., Rubraca (rucaparib), Lynparza (olaparib), Zejula (niraparib)].
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

ALECENSA - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Non-Small Cell Lung Cancer (Adjuvant) only: The member exceeds two years of total Alecensa (alectinib) treatment and member has experienced disease progression while on Alecensa (alectinib).
Required Medical Information	Non-small Cell Lung Cancer: The member has recurrent, advanced or metastatic non-small cell lung cancer AND The member has documented anaplastic lymphoma kinase (ALK)-positive disease AND The member will be using Alecensa (alectinib) as monotherapy. Non-small Cell Lung Cancer (Adjuvant): The member has a diagnosis of non-small cell lung cancer (NSCLC) [i.e., tumors greater than or equal to 4 cm or node positive] AND The member will be using Alecensa (alectinib) for adjuvant treatment following complete tumor resection AND The member has documented anaplastic lymphoma kinase (ALK)-positive disease AND The member will be using Alecensa (alectinib) as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

alosetron - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	The member has a diagnosis of severe diarrhea-predominant irritable bowel syndrome, AND The member has chronic IBS symptoms (lasting 6 months or longer), AND The member has had previous treatment with, contraindication, or intolerance to at least two of the following: loperamide, a tricyclic antidepressant (e.g. amitriptyline, nortriptyline, imipramine HCl, desipramine), or Xifaxan.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ALUNBRIG - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Non-Small cell lung cancer: The member has a diagnosis of recurrent, advanced or metastatic NSCLC AND The member has documented anaplastic lymphoma kinase (ALK)-positive disease AND Alunbrig (brigatinib) will be given as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

alyq - TABLET

PA Criteria	Criteria Details
Off-Label Uses	NA
Exclusion Criteria	NA
Required Medical Information	Pulmonary Arterial Hypertension (PAH). The member must have a diagnosis of pulmonary arterial hypertension (WHO Group I) confirmed by right heart catheterization.
Age Restriction	NA
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	NA
Part B Prerequisite	0

ambrisentan - TABLET

PA Criteria	Criteria Details
Off-Label Uses	NA
Exclusion Criteria	Member has a diagnosis of idiopathic pulmonary fibrosis.
Required Medical Information	Pulmonary Arterial Hypertension (PAH). The member has a diagnosis of pulmonary arterial hypertension (WHO Group I) confirmed by right heart catheterization.
Age Restriction	NA
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	NA
Part B Prerequisite	0

ANKTIVA - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member has experienced disease progression while on Anktiva (nogapendekin alfa inbakicept-pmln).
Required Medical Information	Non-muscle invasive bladder cancer (NMIBC): Member has a diagnosis of non-muscle invasive bladder cancer (NMIBC) with documented carcinoma in situ (CIS) AND The disease is documented as Bacillus Calmette-Guerin (BCG)-unresponsive, which is defined as persistent or recurrent disease following adequate BCG therapy (Adequate BCG is defined as the administration of at least five of six doses of an initial induction course plus either of: at least two of three doses of maintenance therapy or at least two of six doses of a second induction course) AND Member is ineligible for or has elected not to undergo cystectomy AND Anktiva (nogapendekin alfa inbakicept-pmln) will be used concurrently with Bacillus Calmette-Guerin (BCG).
Age Restriction	The member is 18 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Six months duration
Other Criteria	
Part B Prerequisite	0

apomorphine - CARTRIDGE (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Parkinson's Disease off episodes: The member has a diagnosis of Parkinson's disease AND is currently taking carbidopa/levodopa and will continue taking carbidopa/levodopa with Apokyn (apomorphine) AND is experiencing breakthrough off periods related to their Parkinson's disease.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ARCALYST - VIAL (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Familial Cold Autoinflammatory Syndrome or Muckle-Wells Syndrome: The member has a diagnosis of Cryopryin-Associated Periodic Syndrome (CAPS), including Familial Cold Autoinflammatory Syndrome (FCAS) and Muckle-Wells Syndrome (MWS). Recurrent Pericarditis: Member has a diagnosis of recurrent pericarditis defined by: presentation of symptoms of acute pericarditis after a symptom-free interval of at least 4 weeks. Deficiency of Interleukin-1 Receptor Antagonist (DIRA) The member has a diagnosis of deficiency of interleukin-1 receptor antagonist (DIRA).
Age Restriction	Member must be 12 years of age or older for Familial Cold Autoinflammatory Syndrome, Muckle-Wells Syndrome, and Recurrent Pericarditis indications. Age restriction does not apply to DIRA.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ARIKAYCE - VIAL, NEBULIZER (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Use of Arikayce as monotherapy. Use of Arikayce for non-MAC infections or for non-refractory or disseminated MAC. Continued use of Arikayce in members that failed to achieve negative sputum cultures after 6 months of therapy with Arikayce.
Required Medical Information	Mycobacterium avium Complex (MAC) Lung Disease. Member must have a diagnosis of Mycobacterium avium complex (MAC) lung disease. Member must have failed to achieve negative sputum cultures after a minimum of 6 consecutive months of a multi-drug regimen. Member must have limited or no alternative treatment options. Arikayce will be used as part of a multi-drug regimen.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ATROVENT HFA - HFA AEROSOL WITH ADAPTER (GRAM)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Chronic Obstructive Pulmonary Disease (COPD): Diagnosis of Chronic Obstructive Pulmonary Disease (COPD), including chronic bronchitis and emphysema. The member has had previous treatment, contraindication, or intolerance to Spiriva (i.e. Spiriva Handihaler, Spiriva Respimat).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

AUGTYRO - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	ROS1 Positive NSCLC: Member has diagnosis of locally advanced or metastatic non-small cell lung cancer (NSCLC) AND Member has disease which is ROS-1 rearrangement(s) positive AND Member will be using Augtyro (repotrectinib) as monotherapy. NTRK Gene Fusion-Positive Solid Tumors: The member has a diagnosis of solid tumors which are metastatic OR the member is not a candidate for surgical resection AND The member has a documented neurotrophic tyrosine receptor kinase (NTRK) gene fusion AND The members disease has progressed following treatment or does not have satisfactory alternative therapy options AND Member will be using Augtyro (repotrectinib) as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

AUSTEDO - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	TD: The member is utilizing Austedo (deutetrabenazine) or Austedo XR for the treatment of tardive dyskinesia that is associated with the use of dopamine receptor blocking agents AND Symptoms persist despite one of the following: Discontinuation or reduction in dose of dopamine blocking agent(s) OR Discontinuation or reduction in dose of dopamine blocking agent(s) is not possible. OR the member is utilizing Austedo (deutetrabenazine) or Austedo XR for the treatment of TD that is not associated with other medication therapies (e.g. dopamine receptor-blocking agents). Chorea with HD: Diagnosis of chorea associated with Huntington's disease.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

AUSTEDO XR - TABLET, EXTENDED RELEASE 24 HR

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	TD: The member is utilizing Austedo (deutetrabenazine) or Austedo XR for the treatment of tardive dyskinesia that is associated with the use of dopamine receptor blocking agents AND Symptoms persist despite one of the following: Discontinuation or reduction in dose of dopamine blocking agent(s) OR Discontinuation or reduction in dose of dopamine blocking agent(s) is not possible. OR the member is utilizing Austedo (deutetrabenazine) or Austedo XR for the treatment of TD that is not associated with other medication therapies (e.g. dopamine receptor-blocking agents). Chorea with HD: Diagnosis of chorea associated with Huntington's disease.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

AUSTEDO XR TITRATION KT(WK1-4) - TABLET, EXTENDED RELEASE 24 HR DOSE PACK

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	TD: The member is utilizing Austedo (deutetrabenazine) or Austedo XR for the treatment of tardive dyskinesia that is associated with the use of dopamine receptor blocking agents AND Symptoms persist despite one of the following: Discontinuation or reduction in dose of dopamine blocking agent(s) OR Discontinuation or reduction in dose of dopamine blocking agent(s) is not possible. OR the member is utilizing Austedo (deutetrabenazine) or Austedo XR for the treatment of TD that is not associated with other medication therapies (e.g. dopamine receptor-blocking agents). Chorea with HD: Diagnosis of chorea associated with Huntington's disease.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

AVMAPKI-FAKZYNJA - COMBINATION PACKAGE (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on Avmapki (avutometinib)-Fakzynja (defactinib).
Required Medical Information	Ovarian Cancer: The member has a diagnosis of recurrent low-grade serous ovarian cancer (LGSOC) AND The disease is documented as KRAS mutation-positive AND The member has received previous systemic therapy AND Avmapki (avutometinib) will be used in combination with Fakzynja (defactinib).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

AYVAKIT - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member experiences disease progression on Ayvakit (avapritinib). [Applicable to all indications except Gastrointestinal stromal tumor (GIST)]
Required Medical Information	Gastrointestinal Stromal tumor. The member has documented PDGFRA exon 18 mutation-positive unresectable or metastatic gastrointestinal stromal tumor (including PDGFRA D842V) AND Ayvakit (avapritinib) will be given as monotherapy. Advanced systemic mastocytosis: The member has a diagnosis of advanced systemic mastocytosis (AdvSM), including systemic mastocytosis with an associated hematological neoplasm and mast cell leukemia AND Avyakit is not recommended for the treatment of members with AdvSM with platelet counts of less than 50 X 109 /L AND Ayvakit (avapritinib) is administered as monotherapy. Indolent Systemic Mastocytosis (ISM). The member has a diagnosis of Indolent Systemic Mastocytosis (ISM) AND is not recommended for treatment of members with documented platelet counts of less than 50 X 109 /L AND Ayvakit (avapritinib) is administered as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Six month duration
Other Criteria	
Part B Prerequisite	0

azacitidine - VIAL (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Myelodysplastic Syndromes. The member has a diagnosis of myelodysplastic syndrome. Myeloproliferative Neoplasms: The member has a diagnosis of myelofibrosis (MF)-accelerated phase or MF-blast phase/acute myeloid leukemia. Acute Myelogenous Leukemia (AML). The member has a diagnosis of AML.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

BALVERSA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Urothelial Cancer: The member has a diagnosis of locally advanced or metastatic urothelial carcinoma AND the member has identification of a susceptible FGFR3 genetic alteration documented in the medical record [e.g., FGFR3 gene mutations (R248C, S249C, G370C, Y373C), FGFR3 gene fusions (FGFR3-TACC3v1, FGFR3-TACCv3, FGFR3-TACC3, FGFR3-BAIAP2L1)] AND the member will be using Balversa (erdafitinib) as a single agent for subsequent therapy after disease progression during or following at least one prior line of systemic therapy, including a PD-1 (Programmed cell death protein 1) or PD-L1 (Programmed death-ligand 1) inhibitor [e.g., avelumab, pembrolizumab] (if eligible).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 Months Duration
Other Criteria	
Part B Prerequisite	0

BAVENCIO - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression while on or following PD-1/PD-L1 therapy (e.g Keytruda, Opdivo, Tecentriq, Imfinzi).
Required Medical Information	Merkel Cell Carcinoma. The member has a diagnosis of metastatic Merkel cell carcinoma AND Member will be using Bavencio as monotherapy. Urothelial Cancer. The member has a diagnosis of locally advanced or metastatic urothelial cancer AND the member will be using Bavencio (avelumab) as monotherapy AND One of the following apply: The member will be using Bavencio (avelumab) as second or subsequent line systemic therapy OR the member has had disease progression within 12 months of neoadjuvant or adjuvant chemotherapy OR The member will be using Bavencio (avelumab) as maintenance treatment if there is no disease progression with first-line platinum-containing chemotherapy. Renal Cell Carcinoma. The member has a diagnosis of advanced or metastatic renal cell carcinoma AND Bavencio (avelumab) will be given in combination with Inlyta (axitinib) as first-line therapy.
Age Restriction	Merkel Cell Carcinoma – member must be 12 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 months duration
Other Criteria	
Part B Prerequisite	0

BENLYSTA - AUTO-INJECTOR (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Severe active central nervous system lupus.
Required Medical Information	Systemic Lupus Erythematosus (SLE): The member must have a diagnosis of active systemic lupus erythematosus (SLE) AND The member must be positive for autoantibodies relevant to SLE (e.g., ANA, anti-ds DNA, anti-Sm) AND The member must be using Benlysta with standard treatment regimens for SLE (e.g., glucocorticoids, hydroxychloroquine, azathioprine, methotrexate, and/or mycophenolate). Lupus Nephritis: The member must have a diagnosis of active lupus nephritis AND The member must be using Benlysta with standard therapy (e.g. corticosteroids with mycophenolate or cyclophosphamide).
Age Restriction	The member must be 5 years of age or older for Lupus Nephritis and Systemic Lupus Erythematosus (SLE).
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

BEQALZI - TABLET

PA Criteria	Criteria Details
Off-Label Uses	Pending CMS Review
Exclusion Criteria	Pending CMS Review
Required Medical Information	Pending CMS Review
Age Restriction	Pending CMS Review
Prescriber Restriction	Pending CMS Review
Coverage Duration	Pending CMS Review
Other Criteria	Pending CMS Review
Part B Prerequisite	Pending CMS Review

BEQALZI STARTER PACK - TABLET, DOSE PACK

PA Criteria	Criteria Details
Off-Label Uses	Pending CMS Review
Exclusion Criteria	Pending CMS Review
Required Medical Information	Pending CMS Review
Age Restriction	Pending CMS Review
Prescriber Restriction	Pending CMS Review
Coverage Duration	Pending CMS Review
Other Criteria	Pending CMS Review
Part B Prerequisite	Pending CMS Review

BESREMI - SYRINGE (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members with hepatic decompensation (Child-Pugh score greater than 6 [class B and C]). Members that have experienced disease progression while on Besremi (ropeginterferon alfa-2b-njft). Members with existence of, or history of severe psychiatric disorders, particularly severe depression, suicidal ideation or suicide attempt. Members that are immunosuppressed transplant recipients. Members with Hypersensitivity to interferon or to any component of Besremi (ropeginterferon alfa-2b-njft). Members with history or presence of active serious or untreated autoimmune disease.
Required Medical Information	Polycythemia Vera: The member has a diagnosis of Polycythemia Vera.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

BESREMI PEN - PEN INJECTOR (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members with hepatic decompensation (Child-Pugh score greater than 6 [class B and C]). Members that have experienced disease progression while on Besremi (ropeginterferon alfa-2b-njft). Members with existence of, or history of severe psychiatric disorders, particularly severe depression, suicidal ideation or suicide attempt. Members that are immunosuppressed transplant recipients. Members with Hypersensitivity to interferon or to any component of Besremi (ropeginterferon alfa-2b-njft). Members with history or presence of active serious or untreated autoimmune disease.
Required Medical Information	Polycythemia Vera: The member has a diagnosis of Polycythemia Vera.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

bexarotene - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that are pregnant.
Required Medical Information	Cutaneous T-cell Lymphoma (CTCL). (bexarotene capsules). The member will be using bexarotene as primary treatment OR Member has experienced disease progression, contraindication, or intolerance to at least one prior systemic therapy for cutaneous manifestations of cutaneous T-cell lymphoma. Cutaneous T-cell Lymphoma. (bexarotene 1% topical gel/jelly). The member will be using bexarotene as primary treatment OR Member has experienced disease progression, contraindications, or intolerance to at least one prior CTCL therapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

BOMYNTRA - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Concurrent use of bisphosphonate therapy (e.g., zoledronic acid, pamidronate)
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	1

bortezomib - VIAL (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression while on bortezomib.
Required Medical Information	Mantle Cell Lymphoma (MCL): The member has a diagnosis of Mantle Cell Lymphoma(MCL). Multiple Myeloma. The member has a diagnosis of Multiple Myeloma. Waldenström's Macroglobulinemia. The member has a diagnosis of Waldenström's macroglobulinemia AND Velcade (bortezomib) is being used for primary therapy, progressive or relapsed disease or salvage therapy for disease that does not respond to primary therapy AND Velcade (bortezomib) is being used as monotherapy, in combination with Dexamethasone, or in combination with a rituximab product.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

BOSULIF - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on bosutinib. The member has one of the following mutations: T315I, V299L, G250E, or F317L.
Required Medical Information	Chronic Myelogenous Leukemia. The member has a diagnosis of Philadelphia chromosome positive chronic myeloid leukemia (CML) AND One of the following applies: The member has accelerated or blast phase CML OR The member has a diagnosis of chronic phase CML that has not been previously treated and one of the following applies: Intermediate- or high-risk score for disease progression OR Low risk score for disease progression and has contraindication to, intolerance to, or unable to achieve treatment goals with imatinib OR The member has a diagnosis of chronic phase CML that has received previous treatment. Brand Bosulif requests only: Members must have an intolerance or contraindication with generic bosutinib.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

bosutinib - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on bosutinib. The member has one of the following mutations: T315I, V299L, G250E, or F317L.
Required Medical Information	Chronic Myelogenous Leukemia. The member has a diagnosis of Philadelphia chromosome positive chronic myeloid leukemia (CML) AND One of the following applies: The member has accelerated or blast phase CML OR The member has a diagnosis of chronic phase CML that has not been previously treated and one of the following applies: Intermediate- or high-risk score for disease progression OR Low risk score for disease progression and has contraindication to, intolerance to, or unable to achieve treatment goals with imatinib OR The member has a diagnosis of chronic phase CML that has received previous treatment. Brand Bosulif requests only: Members must have an intolerance or contraindication with generic bosutinib.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

BRAFTOVI - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members on concomitant Yervoy (ipilimumab), Opdivo (nivolumab), Keytruda (pembrolizumab), Zelboraf (vemurafenib), Cotellic (cobimetinib), Tafinlar (dabrafenib), or Mekinist (trametinib). Members that have experienced disease progression while on prior anti-BRAF/MEK combination therapy [e.g. Cotellic (cobimetinib) with Zelboraf (vemurafenib) or Tafinlar (dabrafenib) with Mekinist (trametinib)].
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

brivaracetam - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Partial-Onset (Focal) Seizures: Member has a diagnosis of partial-onset (focal) seizures AND Member has been unable to achieve seizure control with levetiracetam AND at least ONE other drug for controlling partial-onset seizures (e.g., lamotrigine, carbamazepine, zonisamide, divalproex, gabapentin or topiramate).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

BRIVIACT - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Partial-Onset (Focal) Seizures: Member has a diagnosis of partial-onset (focal) seizures AND Member has been unable to achieve seizure control with levetiracetam AND at least ONE other drug for controlling partial-onset seizures (e.g., lamotrigine, carbamazepine, zonisamide, divalproex, gabapentin or topiramate).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

BRUKINSA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression while on or following Brukinsa (zanubrutinib).
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

budesonide - TABLET, DELAYED AND EXTENDED RELEASE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Mild to moderate active ulcerative colitis: The member must have a diagnosis of mild to moderate active ulcerative colitis AND the member must have had previous treatment or intolerance to at least two of the following: sulfasalazine, balsalazide capsules, mesalamine enema or mesalamine 0.375g extended-release.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

buprenorphine - PATCH, TRANSDERMAL WEEKLY

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Severe and Persistent Pain: Has persistent, non-acute pain that requires around-the-clock treatment AND Has pain severe enough to require an opioid pain medication AND Has tried or cannot take any one of the following: oxycodone immediate release tablet (non-abuse deterrent formulation, e.g., generic Roxicodone tablet), hydromorphone immediate-release tablet, OR morphine immediate-release tablet.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

BYSANTI - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Schizophrenia: The member has a diagnosis of schizophrenia AND The member has tried or cannot use at least 2 of the following: risperidone, olanzapine, quetiapine, ziprasidone, aripiprazole, or lurasidone. Bipolar I Disorder (manic or mixed episodes): The member has a diagnosis of Bipolar I Disorder (manic or mixed episodes) AND The member has tried or cannot use at least 2 of the following: risperidone, olanzapine, quetiapine, ziprasidone, or aripiprazole.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

BYSANTI TITRATION PACK A - TABLET, DOSE PACK

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Schizophrenia: The member has a diagnosis of schizophrenia AND The member has tried or cannot use at least 2 of the following: risperidone, olanzapine, quetiapine, ziprasidone, aripiprazole, or lurasidone. Bipolar I Disorder (manic or mixed episodes): The member has a diagnosis of Bipolar I Disorder (manic or mixed episodes) AND The member has tried or cannot use at least 2 of the following: risperidone, olanzapine, quetiapine, ziprasidone, or aripiprazole.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

BYSANTI TITRATION PACK B - TABLET, DOSE PACK

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Schizophrenia: The member has a diagnosis of schizophrenia AND The member has tried or cannot use at least 2 of the following: risperidone, olanzapine, quetiapine, ziprasidone, aripiprazole, or lurasidone. Bipolar I Disorder (manic or mixed episodes): The member has a diagnosis of Bipolar I Disorder (manic or mixed episodes) AND The member has tried or cannot use at least 2 of the following: risperidone, olanzapine, quetiapine, ziprasidone, or aripiprazole.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

BYSANTI TITRATION PACK C - TABLET, DOSE PACK

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Schizophrenia: The member has a diagnosis of schizophrenia AND The member has tried or cannot use at least 2 of the following: risperidone, olanzapine, quetiapine, ziprasidone, aripiprazole, or lurasidone. Bipolar I Disorder (manic or mixed episodes): The member has a diagnosis of Bipolar I Disorder (manic or mixed episodes) AND The member has tried or cannot use at least 2 of the following: risperidone, olanzapine, quetiapine, ziprasidone, or aripiprazole.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

CABOMETYX - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Hepatocellular carcinoma, Renal cell carcinoma, and Thyroid cancer: Member experiences disease progression on cabozantinib.
Required Medical Information	Renal cell carcinoma: The member has advanced renal cell carcinoma AND one of the following applies: the member will be using Cabometyx (cabozanitinib) as monotherapy OR the member will be using Cabometyx (cabozanitinib) in combination with a nivolumab product (Opdivo or Opdivo Qvantig) as first line therapy. Hepatocellular carcinoma. The member has a diagnosis of hepatocellular carcinoma AND The member has been previously treated with a first line therapy (e.g., sorafenib) AND Cabometyx (cabozantinib) will be given as monotherapy. Thyroid Cancer: The member has a diagnosis of locally advanced or metastatic differentiated thyroid cancer AND Member has experienced disease progression following prior anti-VEGF targeted therapy AND The disease no longer responds to radioactive iodine treatment (if radioactive iodine is appropriate) AND Cabometyx (cabozantinib) will be administered as monotherapy. Neuroendocrine Tumors (NETs): The member has a diagnosis of unresectable, locally advanced, or metastatic, well-differentiated neuroendocrine tumor (i.e., pNET and epNET).
Age Restriction	Thyroid Cancer and Neuroendocrine Tumors (NETs): Member is 12 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

CALQUENCE (ACALABRUTINIB MAL) - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression while on or following Calquence (acalabrutinib).
Required Medical Information	Mantle Cell Lymphoma: The member has a diagnosis of mantle cell lymphoma AND One of the following: The member has received at least one prior therapy AND the member will be using Calquence (acalabrutinib) as monotherapy OR The member has previously untreated mantle cell lymphoma AND member is ineligible for autologous hematopoietic stem cell transplantation (HSCT) AND Calquence (acalabrutinib) will be used in combination with bendamustine and a rituximab product, followed by maintenance Calquence (acalabrutinib) with or without a rituximab product. Chronic lymphocytic leukemia (CLL) / Small lymphocytic lymphoma (SLL): member has a diagnosis of chronic lymphocytic leukemia (CLL) or small lymphocytic lymphoma (SLL).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

CAPLYTA - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Schizophrenia. The member must have a diagnosis of schizophrenia AND The member has tried or cannot use at least two of the following: risperidone, olanzapine, quetiapine, ziprasidone, aripiprazole, or lurasidone. Bipolar I or II Disorder (Bipolar Depression): The member must have a diagnosis of bipolar I or II disorder (bipolar depression) AND the member has tried or cannot use quetiapine or lurasidone. Major Depressive Disorder: The member has a clinical diagnosis of major depressive disorder (MDD) AND The member has tried or cannot use at least one antidepressant used as monotherapy for MDD AND The member has tried or cannot use at least one generic atypical antipsychotic that has been used with antidepressant therapy AND Caplyta (lumateperone) will be used with antidepressant therapy and not as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

CAPRELSA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on Vandetanib.
Required Medical Information	Thyroid Cancer. The member has a diagnosis of locally advanced or metastatic medullary thyroid cancer AND The member has symptomatic or progressive disease OR the member has a diagnosis of symptomatic or progressive follicular carcinoma or oncocytic carcinoma or papillary carcinoma AND unresectable recurrent or persistent locoregional disease or metastatic disease AND the disease no longer responds to radioactive iodine treatment (if radioactive iodine is appropriate).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

carglumic acid - TABLET, DISPERSIBLE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Acute or chronic hyperammonemia due to NAGS deficiency: the member has acute or chronic hyperammonemia due to deficiency of hepatic enzyme N-acetylglutamate synthase (NAGS). Acute hyperammonemia due to PA or MMA deficiency: the member has acute hyperammonemia due to propionic acidemia (PA) or methylmalonic acidemia (MMA).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

CAVHANZA - TABLET,DISINTEGRATING

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression on any nilotinib product.
Required Medical Information	Chronic Myelogenous Leukemia (CML). The member has a diagnosis of Philadelphia chromosome positive chronic myeloid leukemia (CML) AND One of the following applies: The member has accelerated or blast phase CML OR The member has a diagnosis of chronic phase CML that has not been previously treated, and one of the following applies: Intermediate- or high-risk score for disease progression OR Low-risk score for disease progression and has contraindication to, intolerance to, or unable to achieve treatment goals with imatinib OR The members has a diagnosis of chronic phase CML that has received previous treatment.
Age Restriction	The member is 18 years or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

CAYSTON - VIAL, NEBULIZER (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Patients with a forced expiratory volume in one second (FEV1) less than 25% or greater than 75% predicted. Patients colonized with Burkholderia cepacia.
Required Medical Information	Cystic Fibrosis. The member must have a diagnosis of cystic fibrosis (CF). The member is colonized with Pseudomonas aeruginosa. The member must have a short or long-acting beta-agonist bronchodilator (e.g. albuterol or formoterol), and will be utilized prior to Cayston.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year duration
Other Criteria	
Part B Prerequisite	0

CHORIONIC GONADOTROPIN, HUMAN - VIAL (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Obesity, Female or male infertility, Erectile Dysfunction, Precocious puberty, Prostatic carcinoma or other androgen-dependent neoplasm.
Required Medical Information	NA
Age Restriction	NA
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	NA
Part B Prerequisite	0

clobazam - SUSPENSION, ORAL (FINAL DOSE FORM)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

clozapine - TABLET,DISINTEGRATING

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	The member must be using clozapine ODT or Versacloz (clozapine oral solution) for treatment-resistant schizophrenia. The member has tried or cannot use generic clozapine tablets.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

COBENFY - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Urinary retention. Moderate or severe hepatic impairment. Gastric retention. Untreated narrow-angle glaucoma.
Required Medical Information	Schizophrenia: The member will be using Cobenfy for the treatment of schizophrenia AND The member has tried or cannot use at least two of the following generic atypical antipsychotics: olanzapine, risperidone, aripiprazole, quetiapine, lurasidone, or ziprasidone.
Age Restriction	The member must be 18 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

COBENFY STARTER PACK - CAPSULE, DOSE PACK

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Urinary retention. Moderate or severe hepatic impairment. Gastric retention. Untreated narrow-angle glaucoma.
Required Medical Information	Schizophrenia: The member will be using Cobenfy for the treatment of schizophrenia AND The member has tried or cannot use at least two of the following generic atypical antipsychotics: olanzapine, risperidone, aripiprazole, quetiapine, lurasidone, or ziprasidone.
Age Restriction	The member must be 18 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

COMETRIQ - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression while on Cometriq (cabozantinib).
Required Medical Information	Metastatic Medullary Thyroid Carcinoma. The member has a diagnosis of progressive, metastatic medullary thyroid carcinoma MTC.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

COPIKTRA - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression while on or following a PI3K inhibitor (e.g. idelalisib, copanlisib, duvelisib).
Required Medical Information	Chronic lymphocytic leukemia. The member has a diagnosis of chronic lymphocytic leukemia (CLL) or small lymphocytic lymphoma (SLL) AND the member has relapsed or refractory disease AND The member will be using Copiktra (duvelisib) as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

COSENTYX - SYRINGE (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Moderate to severe chronic plaque psoriasis: The member has a diagnosis of moderate to severe plaque psoriasis AND The member has tried at least one conventional oral systemic treatment (e.g. methotrexate, cyclosporine), or cannot use any conventional oral systemic treatments. Psoriatic Arthritis: The member has a diagnosis of active psoriatic arthritis. Active ankylosing spondylitis: The member has a diagnosis of active ankylosing spondylitis AND the member has tried or cannot use a non-steroidal anti-inflammatory drug (NSAID) (e.g. ibuprofen, meloxicam, naproxen). Non-radiographic Axial Spondyloarthritis: The member has a diagnosis of non-radiographic axial spondyloarthritis with signs of inflammation AND the member has tried or cannot use a nonsteroidal anti-inflammatory drug (NSAID) (e.g. ibuprofen, meloxicam, naproxen). Active Enthesitis-related Juvenile Idiopathic Arthritis: The member has a diagnosis of active enthesitis-related arthritis (ERA) AND the member has tried or cannot use a non-steroidal anti-inflammatory drugs (NSAID) (e.g. ibuprofen, meloxicam, naproxen). Hidradenitis suppurativa: The member must have a diagnosis of moderate to severe hidradenitis suppurativa.
Age Restriction	Plaque Psoriasis: Member must be 6 years of age or older. Non-radiographic axial spondyloarthritis: Member must be 18 years of age or older. Psoriatic Arthritis: member must be 2 years of age or older. Active Enthesitis-related Juvenile Idiopathic Arthritis: member must be 4 years of age or older. Ankylosing Spondylitis and Hidradenitis suppurativa: Member must be 12 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

COSENTYX (2 SYRINGES) - SYRINGE (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Moderate to severe chronic plaque psoriasis: The member has a diagnosis of moderate to severe plaque psoriasis AND The member has tried at least one conventional oral systemic treatment (e.g. methotrexate, cyclosporine), or cannot use any conventional oral systemic treatments. Psoriatic Arthritis: The member has a diagnosis of active psoriatic arthritis. Active ankylosing spondylitis: The member has a diagnosis of active ankylosing spondylitis AND the member has tried or cannot use a non-steroidal anti-inflammatory drug (NSAID) (e.g. ibuprofen, meloxicam, naproxen). Non-radiographic Axial Spondyloarthritis: The member has a diagnosis of non-radiographic axial spondyloarthritis with signs of inflammation AND the member has tried or cannot use a nonsteroidal anti-inflammatory drug (NSAID) (e.g. ibuprofen, meloxicam, naproxen). Active Enthesitis-related Juvenile Idiopathic Arthritis: The member has a diagnosis of active enthesitis-related arthritis (ERA) AND the member has tried or cannot use a non-steroidal anti-inflammatory drugs (NSAID) (e.g. ibuprofen, meloxicam, naproxen). Hidradenitis suppurativa: The member must have a diagnosis of moderate to severe hidradenitis suppurativa.
Age Restriction	Plaque Psoriasis: Member must be 6 years of age or older. Non-radiographic axial spondyloarthritis: Member must be 18 years of age or older. Psoriatic Arthritis: member must be 2 years of age or older. Active Enthesitis-related Juvenile Idiopathic Arthritis: member must be 4 years of age or older. Ankylosing Spondylitis and Hidradenitis suppurativa: Member must be 12 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

COSENTYX PEN - PEN INJECTOR (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Moderate to severe chronic plaque psoriasis: The member has a diagnosis of moderate to severe plaque psoriasis AND The member has tried at least one conventional oral systemic treatment (e.g. methotrexate, cyclosporine), or cannot use any conventional oral systemic treatments. Psoriatic Arthritis: The member has a diagnosis of active psoriatic arthritis. Active ankylosing spondylitis: The member has a diagnosis of active ankylosing spondylitis AND the member has tried or cannot use a non-steroidal anti-inflammatory drug (NSAID) (e.g. ibuprofen, meloxicam, naproxen). Non-radiographic Axial Spondyloarthritis: The member has a diagnosis of non-radiographic axial spondyloarthritis with signs of inflammation AND the member has tried or cannot use a nonsteroidal anti-inflammatory drug (NSAID) (e.g. ibuprofen, meloxicam, naproxen). Active Enthesitis-related Juvenile Idiopathic Arthritis: The member has a diagnosis of active enthesitis-related arthritis (ERA) AND the member has tried or cannot use a non-steroidal anti-inflammatory drugs (NSAID) (e.g. ibuprofen, meloxicam, naproxen). Hidradenitis suppurativa: The member must have a diagnosis of moderate to severe hidradenitis suppurativa.
Age Restriction	Plaque Psoriasis: Member must be 6 years of age or older. Non-radiographic axial spondyloarthritis: Member must be 18 years of age or older. Psoriatic Arthritis: member must be 2 years of age or older. Active Enthesitis-related Juvenile Idiopathic Arthritis: member must be 4 years of age or older. Ankylosing Spondylitis and Hidradenitis suppurativa: Member must be 12 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

COSENTYX PEN (2 PENS) - PEN INJECTOR (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Moderate to severe chronic plaque psoriasis: The member has a diagnosis of moderate to severe plaque psoriasis AND The member has tried at least one conventional oral systemic treatment (e.g. methotrexate, cyclosporine), or cannot use any conventional oral systemic treatments. Psoriatic Arthritis: The member has a diagnosis of active psoriatic arthritis. Active ankylosing spondylitis: The member has a diagnosis of active ankylosing spondylitis AND the member has tried or cannot use a non-steroidal anti-inflammatory drug (NSAID) (e.g. ibuprofen, meloxicam, naproxen). Non-radiographic Axial Spondyloarthritis: The member has a diagnosis of non-radiographic axial spondyloarthritis with signs of inflammation AND the member has tried or cannot use a nonsteroidal anti-inflammatory drug (NSAID) (e.g. ibuprofen, meloxicam, naproxen). Active Enthesitis-related Juvenile Idiopathic Arthritis: The member has a diagnosis of active enthesitis-related arthritis (ERA) AND the member has tried or cannot use a non-steroidal anti-inflammatory drugs (NSAID) (e.g. ibuprofen, meloxicam, naproxen). Hidradenitis suppurativa: The member must have a diagnosis of moderate to severe hidradenitis suppurativa.
Age Restriction	Plaque Psoriasis: Member must be 6 years of age or older. Non-radiographic axial spondyloarthritis: Member must be 18 years of age or older. Psoriatic Arthritis: member must be 2 years of age or older. Active Enthesitis-related Juvenile Idiopathic Arthritis: member must be 4 years of age or older. Ankylosing Spondylitis and Hidradenitis suppurativa: Member must be 12 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

COSENTYX UNOREADY PEN - PEN INJECTOR (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Moderate to severe chronic plaque psoriasis: The member has a diagnosis of moderate to severe plaque psoriasis AND The member has tried at least one conventional oral systemic treatment (e.g. methotrexate, cyclosporine), or cannot use any conventional oral systemic treatments. Psoriatic Arthritis: The member has a diagnosis of active psoriatic arthritis. Active ankylosing spondylitis: The member has a diagnosis of active ankylosing spondylitis AND the member has tried or cannot use a non-steroidal anti-inflammatory drug (NSAID) (e.g. ibuprofen, meloxicam, naproxen). Non-radiographic Axial Spondyloarthritis: The member has a diagnosis of non-radiographic axial spondyloarthritis with signs of inflammation AND the member has tried or cannot use a nonsteroidal anti-inflammatory drug (NSAID) (e.g. ibuprofen, meloxicam, naproxen). Active Enthesitis-related Juvenile Idiopathic Arthritis: The member has a diagnosis of active enthesitis-related arthritis (ERA) AND the member has tried or cannot use a non-steroidal anti-inflammatory drugs (NSAID) (e.g. ibuprofen, meloxicam, naproxen). Hidradenitis suppurativa: The member must have a diagnosis of moderate to severe hidradenitis suppurativa.
Age Restriction	Plaque Psoriasis: Member must be 6 years of age or older. Non-radiographic axial spondyloarthritis: Member must be 18 years of age or older. Psoriatic Arthritis: member must be 2 years of age or older. Active Enthesitis-related Juvenile Idiopathic Arthritis: member must be 4 years of age or older. Ankylosing Spondylitis and Hidradenitis suppurativa: Member must be 12 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

COTELLIC - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Melanoma indication only: Members on Cotellic as a single agent. Members on concomitant Yervoy (ipilimumab), Opdivo (nivolumab), Keytruda(pembrolizumab), Tafinlar (dabrafenib), Mekinist (trametinib), Braftovi (encorafenib), or Mektovi (binimetinib). Members that have experienced disease progression while on Cotellic. Members that have experienced disease progression while on prior anti-BRAF/MEK combination therapy [e.g.,Braftovi (encorafenib) with Mektovi (binimetinib) or Tafinlar (dabrafenib) with Mekinist (trametinib)].
Required Medical Information	Melanoma: The member has a diagnosis of unresectable or stage IV metastatic melanoma AND The member has a documented BRAF V600 activating mutation AND The member will be using Cotellic (cobimetinib) in combination with Zelboraf(vemurafenib). Histiocytic Neoplasms: the member has a diagnosis of histiocytic neoplasms AND the member will be using Cotellic (cobimetinib) as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

CRESEMBA - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Familial short QT syndrome.
Required Medical Information	Invasive Aspergillosis and Invasive Mucormycosis: Member must have diagnosis of invasive aspergillosis or invasive mucormycosis.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

CYRAMZA - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on Cyramza (ramucirumab).
Required Medical Information	Gastric Cancer: member has a diagnosis of advanced or metastatic gastric cancer or gastro-esophageal adenocarcinoma AND the member has disease progression or intolerance on or after prior therapy with platinum-based or fluoropyrimidine-based chemotherapy AND Cyramza (ramucirumab) will be used as subsequent therapy AND will be used as monotherapy or in combination with paclitaxel. Non-Small Cell Lung Cancer: The member has a diagnosis of metastatic non-small cell lung cancer AND The member has disease progression or intolerance on or after prior therapy with platinum-based chemotherapy AND For members with EGFR (epidermal growth factor receptor) or ALK (anaplastic lymphoma kinase) genomic aberrations: the member has disease progression on FDA-approved therapy for these aberrations and Cyramza will be used in combination with Docetaxel OR member has documented epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) mutations AND Cyramza (ramucirumab) is given in combo with erlotinib as first line therapy. Colorectal Cancer: The member has a diagnosis of unresectable or metastatic colorectal cancer AND Cyramza (ramucirumab) will be given as primary treatment in combination with irinotecan or FOLFIRI (fluorouracil, leucovorin calcium, and irinotecan) for unresectable metachronous metastases and previous treatment with FOLFOX (fluorouracil, leucovorin calcium, and oxaliplatin) or CapeOX (capecitabine, oxaliplatin) as adjuvant therapy has been given OR The member has disease progression on or after prior therapy with a bevacizumab product, oxaliplatin, and a fluoropyrimidine (e.g. 5-fluorouracil, capecitabine) AND Cyramza is given in combination with FOLFIRI (irinotecan, folinic acid,and 5-fluorouracil) or irinotecan as therapy after first progression of disease if irinotecan was not previously given.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration

CYRAMZA - VIAL (ML)

Other Criteria	Esophageal Cancer: The member has a diagnosis of unresectable locally advanced, recurrent, or metastatic esophageal adenocarcinoma AND Cyramza will be used as second or subsequent line therapy with or without paclitaxel. Hepatocellular Carcinoma: The member has a diagnosis of metastatic or unresectable hepatocellular carcinoma AND the member has received prior treatment with a first line therapy (e.g.,sorafenib) AND the member has alpha fetoprotein greater than or equal to 400 ng/ml AND Cyramza (ramucirumab) will be given as a single agent as subsequent therapy.
Part B Prerequisite	0

CYSTARAN - DROPS

PA Criteria	Criteria Details
Off-Label Uses	NA
Exclusion Criteria	NA
Required Medical Information	Cystinosis: The member has a diagnosis of cystinosis AND The member is using cysteamine ophthalmic solution in the treatment of corneal cystine crystal accumulation.
Age Restriction	NA
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	NA
Part B Prerequisite	0

DANYELZA - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members receiving Danyelza (naxitamab) as monotherapy. Members that have experienced disease progression while on Danyelza (naxitamabgqgk).
Required Medical Information	Relapsed or Refractory Neuroblastoma: The member has a diagnosis of relapsed or refractory high-risk neuroblastoma AND The disease is in the bone or bone marrow AND The member has achieved a partial or minor response or stable disease to prior therapy AND Danyelza (naxitamab-gqgk) will be used in combination with Leukine (sargramostim).
Age Restriction	The member is 1 year of age and older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

DARZALEX - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Disease progression while taking Darzalex (daratumumab).
Required Medical Information	Multiple Myeloma:The member has a diagnosis of multiple myeloma AND The member will be using Darzalex (daratumumab) for newly diagnosed disease AND the member is ineligible for autologous stem cell transplant AND the member will be using Darzalex (daratumumab) in combination with bortezomib, melphalan, and prednisone OR the member will be using Darzalex in combination with lenalidomide and dexamethasone OR the member is eligible for autologous stem cell transplant AND the member will be using Darzalex in combination with bortezomib, thalidomide, and dexamethasone OR the member will be using Darzalex (daratumumab) for relapsed, progressive, or refractory disease in one of the following scenarios: The member will be using Darzalex (daratumumab) in combination with Pomalyst (pomalidomide) and dexamethasone AND the member has received at least two prior therapies, including lenalidomide and a proteasome inhibitor (e.g. bortezomib, carfilzomib, or ixazomib) OR The member will be using Darzalex (daratumumab) in combination with Velcade (bortezomib) and dexamethasone OR The member will be using Darzalex (daratumumab) in combination with Revlimid (lenalidomide) and dexamethasone OR the member will be using Darzalex (daratumumab) in combination with Kyprolis (carfilzomib) OR The member will be using Darzalex (daratumumab) as monotherapy and one of the following applies: The member has received at least three prior lines of therapy, which must have included a proteasome inhibitor (e.g. bortezomib or carfilzomib) and an immunomodulatory drug (e.g. thalidomide, lenalidomide, or pomalidomide) OR The member is double-refractory to a proteasome inhibitor (e.g. bortezomib or carfilzomib) and an immunomodulatory drug (e.g. thalidomide, lenalidomide, or pomalidomide). Omission of corticosteroid from regimen is allowed if intolerance/contraindication.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

DARZALEX FASPRO - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Disease progression while taking daratumumab. Members who have NYHA Class IIIB or Class IV cardiac disease or Mayo Stage IIIB (Applicable to Light chain (AL) amyloidosis indication only).
Required Medical Information	Multiple Myeloma: The member has a diagnosis of multiple myeloma AND The member will be using Darzalex (daratumumab) Faspro for newly diagnosed disease in one of the following scenarios AND the member is ineligible for autologous stem cell transplant AND the member will be using Darzalex Faspro in combination with bortezomib, melphalan, and prednisone OR the member will be using Darzalex Faspro in combination with lenalidomide and dexamethasone OR the member will be using Darzalex Faspro in combination with bortezomib, lenalidomide, and dexamethasone OR the member is eligible for autologous stem cell transplant AND the member will be using in combination with bortezomib, thalidomide, and dexamethasone OR the member will be using in combination with bortezomib, lenalidomide, and dexamethasone OR the member will be using Darzalex Faspro for relapsed or progressive disease in one of the following scenarios: in combination with Velcade (bortezomib) and dexamethasone OR in combination with Kyprolis (carfilzomib) and dexamethasone OR in combination with Revlimid (lenalidomide) and dexamethasone OR in combination with Pomalyst (pomalidomide) and dexamethasone OR as monotherapy and one of the following applies: The member has received at least three prior lines of therapy, which must have included a proteasome inhibitor (e.g. bortezomib or carfilzomib) and an immunomodulatory drug (e.g. thalidomide, lenalidomide, or pomalidomide) OR The member is double-refractory to a proteasome inhibitor (e.g. bortezomib or carfilzomib) and an immunomodulatory drug (e.g. thalidomide, lenalidomide, or pomalidomide). Omission of corticosteroid from regimen is allowed if intolerance/contraindication.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration

DARZALEX FASPRO - VIAL (ML)

Other Criteria	Light Chain Amyloidosis: The member has a diagnosis of light chain amyloidosis AND The member will be using Darzalex Faspro for newly diagnosed disease AND The member will be using in combination with bortezomib, cyclophosphamide, and dexamethasone. Omission of corticosteroid from regimen is allowed if intolerance/contraindication. Smoldering Multiple Myeloma: the member has a diagnosis of smoldering multiple myeloma AND the member's disease is categorized as high-risk: e.g. According to the Mayo 2018 & International Myeloma Working Group (IMWG) 20/2/20 criteria, high-risk is defined as the presence of at least two of the following: serum M-protein greater than 2g/dL OR bone marrow plasma cells greater than 20% OR free light chain ratio greater than 20 AND The member will be using Darzalex Faspro as monotherapy.
Part B Prerequisite	0

dasatinib - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on dasatinib. For ALL and CML: The member has one of the following mutations: T315I/A, F317L/V/I/C or V299L.
Required Medical Information	Chronic Myelogenous Leukemia (CML): The member has a diagnosis of Ph+ chronic myeloid leukemia (CML) and one of the following applies: The member has accelerated or blast phase CML OR The member has a diagnosis of chronic phase CML that has not been previously treated and one of the following applies: Intermediate- or high-risk score for disease progression OR Low risk score for disease progression and has contraindication to, intolerance to, or unable to achieve treatment goals with imatinib OR The member has a diagnosis of chronic phase CML that has received previous treatment. Acute Lymphoblastic Leukemia (ALL): The member has ALL (Philadelphia Chromosome positive) and dasatinib is being used for induction or consolidation treatment in combination with chemotherapy or corticosteroids OR treatment is for maintenance therapy or the treatment of members with resistance or intolerance to prior therapy. Advanced Gastrointestinal Stromal Tumor (GIST): The member has a diagnosis of advanced unresectable GIST. The member has progressive disease or is intolerant to prior therapy with imatinib or sunitinib or Stivarga (regorafenib). [Pediatric] Chronic Myelogenous Leukemia (CML). The member has a diagnosis of chronic myeloid leukemia (CML) that is Philadelphia chromosome positive (Ph+) AND the member is in chronic phase. [Pediatric] Acute lymphoblastic leukemia (ALL). The member has a diagnosis of acute lymphoblastic leukemia (ALL) AND the member has Philadelphia chromosome positive (Ph+) disease AND the member has newly-diagnosed disease AND The member will be using dasatinib in combination with chemotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

DAURISMO - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression while on Daurismo (glasdegib).
Required Medical Information	Acute Myeloid Leukemia. The member has a diagnosis of newly-diagnosed acute myeloid leukemia (AML) AND One of the following applies: The member is age 75 years or older OR The member has comorbidities that preclude the use of intensive induction chemotherapy (e.g. severe cardiac disease, baseline Eastern Cooperative Oncology Group (ECOG) performance status of 2, or baseline serum creatinine greater than 1.3 mg/dL) AND The member will be using Daurismo (glasdegib) in combination with low-dose Cytarabine.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

decitabine - VIAL (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Myelodysplastic Syndromes. The member has a diagnosis of myelodysplastic syndrome. Myeloproliferative Neoplasms: The member has a diagnosis of myelofibrosis (MF)-accelerated phase or MF-blast phase/acute myeloid leukemia. Acute Myelogenous Leukemia (AML). The member has a diagnosis of AML.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

deferasirox - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Platelet counts less than 50,000/mcL. (Per FDA label, Deferasirox is contraindicated in patients with platelets below 50,000).
Required Medical Information	Chronic Iron Toxicity (hemosiderosis) Secondary to Transfusional Iron Overload. (Initial Request): The member must meet ALL of the following criteria: Diagnosis of chronic iron overload (hemosiderosis) secondary to multiple RBC transfusions AND Ferritin level is consistently greater than 1000mcg/L. (Continuation of Therapy): The member must meet ALL of the following criteria: Ferritin level remains consistently at or above 500mcg/L while taking deferasirox therapy. Chronic iron overload in patients with non-transfusion dependent thalassemia (NTDT) syndromes: The member must meet ALL of the following criteria: Has a diagnosis of chronic iron overload with non-transfusion dependent thalassemia (NTDT) syndromes AND Has a liver iron (Fe) concentration of at least 5mg Fe/gram of liver dry weight AND Has a serum ferritin greater than 300 mcg/L. (Continuation of Therapy): The member must meet ALL of the following criteria: Has a liver iron (Fe) concentration of at least 3 mg/gram of liver dry weight AND has a serum ferritin greater than or equal to 300 mcg/L.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Initial: Plan Year Duration. Reauthorization: Plan year duration.
Other Criteria	
Part B Prerequisite	0

dextromethorphan-quinidine - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	Pending CMS Review
Exclusion Criteria	Pending CMS Review
Required Medical Information	Pending CMS Review
Age Restriction	Pending CMS Review
Prescriber Restriction	Pending CMS Review
Coverage Duration	Pending CMS Review
Other Criteria	Pending CMS Review
Part B Prerequisite	Pending CMS Review

DIACOMIT - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	The member has a diagnosis of seizures associated with Dravet syndrome AND The member is refractory on current therapy (e.g experiencing generalized tonic clonic or clonic seizures within the past 28 days) AND The member is taking concomitant clobazam.
Age Restriction	
Prescriber Restriction	Diacomit is prescribed by or in consultation with a specialist (i.e. neurologist, epileptologist).
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

diclofenac sodium - GEL (GRAM)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Actinic Keratosis: The member has a diagnosis of actinic keratosis. The member has trial, intolerance, or contraindication to generic imiquimod 5% cream or topical fluorouracil.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

dihydroergotamine - AEROSOL, SPRAY WITH PUMP (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Acute treatment of moderate to severe migraine headaches with or without aura AND has tried or cannot use two of the following: naproxen tablet, naratriptan tablet, rizatriptan tablet, sumatriptan tablet, eletriptan tablet.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

dimethyl fumarate - CAPSULE,DELAYED RELEASE (ENTERIC COATED)

PA Criteria	Criteria Details
Off-Label Uses	NA
Exclusion Criteria	NA
Required Medical Information	The member has a diagnosis of a relapsing form of multiple sclerosis, to include relapsing-remitting or active secondary progressive disease OR the member has a diagnosis of a clinically isolated syndrome (CIS).
Age Restriction	NA
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	NA
Part B Prerequisite	0

DRIZALMA SPRINKLE - CAPSULE, DELAYED RELEASE SPRINKLE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Major Depressive Disorder, Generalized Anxiety Disorder, or Diabetic Peripheral Neuropathic Pain: The member has a diagnosis of Major Depressive Disorder (MDD), Generalized Anxiety Disorder (GAD), or Diabetic Peripheral Neuropathic Pain (DPNP). The member has prior therapy, intolerance, or contraindication with venlafaxine (IR or ER) AND duloxetine 20mg, 30mg or 60 mg (generic Cymbalta). Chronic Musculoskeletal Pain, Fibromyalgia: The member has a diagnosis of Chronic Musculoskeletal Pain or Fibromyalgia (FM). The member has prior therapy, intolerance, or contraindication with duloxetine 20mg, 30mg or 60 mg (generic Cymbalta).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

droxidopa - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Neurogenic Orthostatic Hypotension: The member has symptomatic, neurogenic orthostatic hypotension (NOH) caused by: Primary autonomic failure- Parkinson's disease (PD), multiple system atrophy or pure autonomic failure OR Dopamine beta-hydroxylase deficiency OR Non-diabetic neuropathy. AND the member must have had previous treatment, intolerance, or contraindication to one of the following agents: fludrocortisone OR midodrine.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

DUAVEE - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Treatment of moderate to severe vasomotor symptoms associated with menopause:Diagnosis of moderate to severe vasomotor symptoms associated with menopause AND The member must have had previous treatment, intolerance or contraindication to a SSRI [e.g. citalopram, fluoxetine, paroxetine hydrochloride] or venlafaxine. Prevention of osteoporosis: For the prevention of osteoporosis in a member who is postmenopausal AND the member must have had previous treatment, intolerance, or contraindication to either alendronate or Evista (raloxifene).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

DUPIXENT PEN - PEN INJECTOR (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	Moderate-to-severe asthma with an eosinophilic phenotypic or with oral corticosteroid dependent asthma: Member is 6 years of age or older, Atopic dermatitis: Member is 6 months of age or older. Chronic rhinosinusitis with nasal polyposis: Member is 12 years of age or older. Prurigo Nodularis, Bullous Pemphigoid, and COPD with an eosinophilic phenotype: Member is 18 years of age or older. Eosinophilic Esophagitis: 1 year of age or older. CSU: Member is 2 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration

Other Criteria	Moderate-to-severe asthma with an eosinophilic phenotypic or with oral corticosteroid dependent asthma: The member has a diagnosis of moderate-to-severe asthma AND The member has an eosinophilic phenotype, defined by an elevated peripheral blood eosinophil level of: greater than or equal to 150 cells/uL at therapy initiation, OR greater than or equal to 300 cells/uL within 12 months prior to therapy initiation, OR The member has oral corticosteroid-dependent asthma AND The member has been unable to achieve adequate control of asthma while on maximum tolerated inhaled corticosteroid therapy in combination with a long acting beta agonist (eg formoterol). Chronic Rhinosinusitis with Nasal Polyposis: The member must have a diagnosis of Chronic Rhinosinusitis with Nasal Polyposis AND Dupixent (dupilumab) will be used in conjunction with a daily intranasal corticosteroid spray AND The member has been unable to achieve adequate control of symptoms with maximum tolerated intranasal corticosteroid therapy. Prurigo Nodularis: member must meet all of the following criteria: diagnosis of Prurigo Nodularis Chronic Obstructive Pulmonary Disease (COPD) with an eosinophilic phenotype: The member has a diagnosis of Chronic Obstructive Pulmonary Disease (COPD) AND The member has an eosinophilic phenotype defined as an elevated peripheral blood eosinophil level of greater than or equal to 300 cells/uL within 12 months prior to therapy initiation AND The member has been unable to achieve adequate control of COPD while on an: Inhaled corticosteroid therapy in combination with a long-acting beta agonist (LABA) and long-acting muscarinic antagonist (LAMA) OR LABA-LAMA therapy if inhaled corticosteroids are contraindicated. Bullous Pemphigoid: The member has a diagnosis of bullous pemphigoid confirmed by biopsy or serologic tests AND Prescribed by or in consultation with a dermatologist, allergist, or immunologist AND member has had previous treatment with, intolerance or contraindication to one high potency topical corticosteroid (e.g. augmented betamethasone dipropionate 0.05%, clobetasol cream/ointment 0.05%, triamcinolone acetonide 0.5%) OR oral corticosteroid (e.g. prednisone). Allergic Fungal Rhinosinusitis (AFRS): Member has a diagnosis of allergic fungal rhinosinusitis (AFRS) AND Member has a history of sino-nasal surgery.
Part B Prerequisite	0

DUPIXENT SYRINGE - SYRINGE (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	Moderate-to-severe asthma with an eosinophilic phenotypic or with oral corticosteroid dependent asthma: Member is 6 years of age or older, Atopic dermatitis: Member is 6 months of age or older. Chronic rhinosinusitis with nasal polyposis: Member is 12 years of age or older. Prurigo Nodularis, Bullous Pemphigoid, and COPD with an eosinophilic phenotype: Member is 18 years of age or older. Eosinophilic Esophagitis: 1 year of age or older. CSU: Member is 2 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration

DUPIXENT SYRINGE - SYRINGE (ML)

Other Criteria	Moderate-to-severe asthma with an eosinophilic phenotypic or with oral corticosteroid dependent asthma: The member has a diagnosis of moderate-to-severe asthma AND The member has an eosinophilic phenotype, defined by an elevated peripheral blood eosinophil level of: greater than or equal to 150 cells/uL at therapy initiation, OR greater than or equal to 300 cells/uL within 12 months prior to therapy initiation, OR The member has oral corticosteroid-dependent asthma AND The member has been unable to achieve adequate control of asthma while on maximum tolerated inhaled corticosteroid therapy in combination with a long acting beta agonist (eg formoterol). Chronic Rhinosinusitis with Nasal Polyposis: The member must have a diagnosis of Chronic Rhinosinusitis with Nasal Polyposis AND Dupixent (dupilumab) will be used in conjunction with a daily intranasal corticosteroid spray AND The member has been unable to achieve adequate control of symptoms with maximum tolerated intranasal corticosteroid therapy. Prurigo Nodularis: member must meet all of the following criteria: diagnosis of Prurigo Nodularis Chronic Obstructive Pulmonary Disease (COPD) with an eosinophilic phenotype: The member has a diagnosis of Chronic Obstructive Pulmonary Disease (COPD) AND The member has an eosinophilic phenotype defined as an elevated peripheral blood eosinophil level of greater than or equal to 300 cells/uL within 12 months prior to therapy initiation AND The member has been unable to achieve adequate control of COPD while on an: Inhaled corticosteroid therapy in combination with a long-acting beta agonist (LABA) and long-acting muscarinic antagonist (LAMA) OR LABA-LAMA therapy if inhaled corticosteroids are contraindicated. Bullous Pemphigoid: The member has a diagnosis of bullous pemphigoid confirmed by biopsy or serologic tests AND Prescribed by or in consultation with a dermatologist, allergist, or immunologist AND member has had previous treatment with, intolerance or contraindication to one high potency topical corticosteroid (e.g. augmented betamethasone dipropionate 0.05%, clobetasol cream/ointment 0.05%, triamcinolone acetonide 0.5%) OR oral corticosteroid (e.g. prednisone). Allergic Fungal Rhinosinusitis (AFRS): Member has a diagnosis of allergic fungal rhinosinusitis (AFRS) AND Member has a history of sino-nasal surgery.
Part B Prerequisite	0

ELELYSO - VIAL (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	NA
Required Medical Information	Gaucher Disease. The member has a confirmed diagnosis of Type 1 Gaucher disease.
Age Restriction	NA
Prescriber Restriction	Licensed Practitioner.
Coverage Duration	Plan Year Duration.
Other Criteria	NA
Part B Prerequisite	0

ELIGARD - SYRINGE (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ELIGARD (3 MONTH) - SYRINGE (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ELIGARD (4 MONTH) - SYRINGE (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ELIGARD (6 MONTH) - SYRINGE (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

eltrombopag olamine - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Initial: 6 Months Duration. Continuation: Plan year duration.
Other Criteria	
Part B Prerequisite	0

EMGALITY PEN - PEN INJECTOR (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

EMGALITY SYRINGE - SYRINGE (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

EMPLICITI - VIAL (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members with disease progression while on Empliciti (elotuzumab)
Required Medical Information	Multiple Myeloma:The member has a diagnosis of multiple myeloma AND One of the following scenarios apply: The member has disease progression after receiving one to three prior lines of therapy AND Empliciti (elotuzumab) will be given in combination with lenalidomide (Revlimid) and dexamethasone OR in combination with bortezomib (Velcade) and dexamethasone OR The member has disease progression after receiving at least two prior therapies, including lenalidomide and a proteasome inhibitor AND Empliciti (elotuzumab) will be given in combination with pomalidomide (Pomalyst) and dexamethasone (Omission of corticosteroid from regimen is allowed if intolerance/contraindication).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

EMSAM - PATCH, TRANSDERMAL 24 HOURS

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Major Depressive Disorder: The member is an adult with a clinical diagnosis of major depressive disorder (MDD). The member has tried or cannot use a generic SSRI (e.g. citalopram, fluoxetine, paroxetine, or sertraline), generic SNRI (e.g. venlafaxine or duloxetine), a generic bupropion product (75mg/100mg IR, 100mg/150mg/200mg SR, or 150mg/300mg XL) OR mirtazapine.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ENBREL - SYRINGE (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	Must be 18 years of age for Ankylosing Spondylitis, Psoriatic arthritis, or RA. Must be 4 years of age for Plaque Psoriasis. Must be 2 years of age for Polyarticular Juvenile Idiopathic Arthritis and Juvenile Psoriatic Arthritis.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ENBREL MINI - CARTRIDGE (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	Must be 18 years of age for Ankylosing Spondylitis, Psoriatic arthritis, or RA. Must be 4 years of age for Plaque Psoriasis. Must be 2 years of age for Polyarticular Juvenile Idiopathic Arthritis and Juvenile Psoriatic Arthritis.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ENBREL SURECLICK - PEN INJECTOR (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	Must be 18 years of age for Ankylosing Spondylitis, Psoriatic arthritis, or RA. Must be 4 years of age for Plaque Psoriasis. Must be 2 years of age for Polyarticular Juvenile Idiopathic Arthritis and Juvenile Psoriatic Arthritis.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ENSACOVE - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

ENTYVIO PEN - PEN INJECTOR (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Ulcerative Colitis: The member has a diagnosis of moderately to severely active ulcerative colitis AND The member has tried or cannot use TWO of the following: preferred adalimumab product [i.e., adalimumab-adbm, adalimumab-adaz], preferred ustekinumab product (i.e., Otulfi, Yesintek), Skyrizi, Rinvoq, or Tremfya OR the member has completed induction doses with intravenous Entyvio. Crohns Disease: The member has a diagnosis of moderately to severely active Crohns disease AND The member has tried or cannot use TWO of the following: preferred adalimumab product [i.e., adalimumab-adbm, adalimumab-adaz], preferred ustekinumab product (i.e., Otulfi, Yesintek), Skyrizi, Rinvoq, or Tremfya OR the member has completed induction doses with intravenous Entyvio.
Age Restriction	18 years of age or older
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ENVARSUS XR - TABLET, EXTENDED RELEASE 24 HR

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Member must have had a kidney transplant AND Must be using Envarsus XR for prophylaxis of organ rejection AND Must be using in combination with other immunosuppressants.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

EPIDIOLEX - SOLUTION, ORAL

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Dravet Syndrome: The member has a diagnosis of seizures associated with Dravet Syndrome AND The member has been unable to achieve seizure control with at least one other drug used for the treatment of Dravet Syndrome (e.g. clobazam, valproic acid, topiramate). Lennox-Gastaut Syndrome: The member has a diagnosis of seizures associated with Lennox-Gastaut syndrome AND Epidiolex (cannabidiol) is prescribed by or in consultation with a specialist (i.e. neurologist, epileptologist) AND The member has been unable to achieve seizure control with at least one other drug used for the treatment of Lennox-Gastaut syndrome (e.g., valproic acid, lamotrigine, topiramate). Tuberous Sclerosis Complex: The member has a diagnosis of seizures associated with Tuberous Sclerosis Complex.
Age Restriction	
Prescriber Restriction	Epidiolex (cannabidiol) is prescribed by or in consultation with a specialist (i.e. neurologist, epileptologist).
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

EPRONTIA - SOLUTION, ORAL

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Migraine Prophylaxis: Member is using for prophylaxis of migraine headache AND has had previous treatment with or intolerance to immediate release topiramate tablet or capsule AND has had previous treatment, intolerance, or contraindication with one of the following: a seratonin and norepinephrine reuptake inhibitor or a tricyclic antidepressant. Epilepsy Adjunctive Therapy: Member has a diagnosis of partial-onset (focal) seizures, primary generalized tonic-clonic seizures, or seizures associated with Lennox-Gastaut syndrome (LGS) AND Member will use Eprontia (topiramate) in combination with at least one other drug for controlling seizures AND Member has tried or cannot use immediate-release topiramate tablet or capsule. Epilepsy Monotherapy: Member has a diagnosis of partial-onset (focal) seizures or primary generalized tonic-clonic seizures AND Member has tried or cannot use immediate release topiramate tablet or capsule.
Age Restriction	Migraine Prophylaxis: Member must be 12 years of age or older. Epilepsy indications: Member must be 2 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

ERBITUX - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members with colorectal cancer (CRC) in which RAS mutation status is unknown [Applies to Colorectal Cancer [Wild-Type RAS; Wild-Type KRAS/NRAS]and Colorectal Cancer [KRAS G12C-mutated]. Member has had disease progression on Vectibix or Erbutux. [All Indications]
Required Medical Information	Colorectal Cancer [Wild-Type RAS; Wild-Type KRAS/NRAS]. The member has a diagnosis of advanced or metastatic colorectal cancer (mCRC) AND The member has colorectal cancer that is documented as wild-type, non-mutated KRAS and NRAS [wild-type RAS] AND Erbitux (cetuximab) will be used as one of the following: as monotherapy in members intolerant to irinotecan or who have experienced disease progression following therapy with both irinotecan and oxaliplatin based therapy OR in combination with irinotecan-based therapy OR in combination with fluorouracil based therapy (CAPEOX, FOLFOX, FOLFIRI). Head and Neck Cancer. The member has a diagnosis of locally or regionally advanced squamous cell head and neck cancer with concomitant XRT (radiation) OR The member has recurrent (including locoregional), unresectable, or metastatic squamous cell head and neck cancer and one of the following applies: Erbitux (cetuximab) will be given as monotherapy or Erbitux (cetuximab) will be given in conjunction with a platinum agent OR Erbitux (cetuximab) will be given in combination with platinum-based therapy with 5-FU (fluorouracil). Colorectal Cancer [BRAF V600E-mutated]: The member has documented BRAF V600E metastatic colorectal cancer (mCRC) AND One of the following scenarios applies: The member has documented BRAF V600E metastatic colorectal cancer and progressive disease on prior therapy and Erbitux (cetuximab) will be given in combination with Braftovi (encorafenib) OR The member has documented BRAF V600E metastatic colorectal cancer and Erbitux (cetuximab) will be given in combination with Braftovi (encorafenib) and FOLFOX (fluorouracil, leucovorin, and oxaliplatin) regimen.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration

ERBITUX - VIAL (ML)

Other Criteria	Colorectal Cancer [KRAS G12C-mutated]. The member has a diagnosis of locally advanced or metastatic colorectal cancer (mCRC) AND The colorectal cancer (CRC) has a documented KRAS G12C mutation, as determined by an FDA-approved test AND The member has received prior systemic treatment with chemotherapy [fluoropyrimidine-based chemotherapy (e.g., 5-fluorouracil, capecitabine), oxaliplatin, irinotecan] AND Erbitux (cetuximab) will be given in combination with Krazati (adagrasib).
Part B Prerequisite	0

ERIVEDGE - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Erivedge (vismodegib) therapy is not considered medically necessary for members with the following concomitant conditions: Members that have experienced disease progression while on Erivedge (vismodegib).
Required Medical Information	Advanced Basal Cell Carcinoma. The member has a diagnosis of metastatic basal cell carcinoma OR The member has a diagnosis of locally advanced basal cell carcinoma AND one of the following applies: The member has disease that has recurred following surgery OR the member is not a candidate for surgery.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

ERLEADA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on Erleada (apalutamide).
Required Medical Information	Prostate Cancer (non-metastatic castration resistant): The member has a diagnosis of non-metastatic castration resistant prostate cancer AND the member will use Erleada (apalutamide) in combination with androgen deprivation therapy (e.g. previous bilateral orchioectomy or luteinizing hormone-releasing hormone (LHRH) analog [i.e., LHRH agonist/ antagonist]). Prostate Cancer (metastatic castration-sensitive): The member has a diagnosis of metastatic castration-sensitive prostate cancer AND the member will use Erleada (apalutamide) in combination with androgen deprivation therapy (e.g. previous bilateral orchioectomy or LHRH analog [i.e., LHRH agonist/ antagonist]).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

erlotinib - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pancreatic Cancer: The member has a diagnosis of unresectable, locally advanced or metastatic pancreatic cancer. AND erlotinib is being used in combination with gemcitabine. Non-small cell lung cancer. The member has a diagnosis of metastatic NSCLC AND all of the following apply: The member has known documented activated EGFR mutation (such as E19del in exon 19 or L858R in exon 21). Renal Cell Carcinoma: Diagnosis of relapsed or unresectable stage IV renal cell carcinoma with non clear histology.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 months duration
Other Criteria	
Part B Prerequisite	0

eslicarbazepine - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ETCAMAH - TABLET

PA Criteria	Criteria Details
Off-Label Uses	Pending CMS Review
Exclusion Criteria	Pending CMS Review
Required Medical Information	Pending CMS Review
Age Restriction	Pending CMS Review
Prescriber Restriction	Pending CMS Review
Coverage Duration	Pending CMS Review
Other Criteria	Pending CMS Review
Part B Prerequisite	Pending CMS Review

EULEXIN - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	The member has a diagnosis of prostate cancer AND will be using Eulexin (flutamide) alone or in combination therapy with a luteinizing hormone-releasing hormone (LHRH) analog.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

EVDI - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	Pending CMS Review
Exclusion Criteria	Pending CMS Review
Required Medical Information	Pending CMS Review
Age Restriction	Pending CMS Review
Prescriber Restriction	Pending CMS Review
Coverage Duration	Pending CMS Review
Other Criteria	Pending CMS Review
Part B Prerequisite	Pending CMS Review

everolimus (antineoplastic) - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on everolimus.
Required Medical Information	Advanced Renal Cell Carcinoma (RCC). The member has a diagnosis of advanced /metastatic renal cell carcinoma (stage IV). Subependymal Giant Cell Astrocytoma (SEGA) with TSC [Adults and Pediatrics]. The member has a diagnosis of Subependymal Giant Cell Astrocytoma (SEGA) associated with tuberous sclerosis complex AND The member requires therapeutic intervention but is not a candidate for curative surgical resection. Neuroendocrine Tumors: The member has disease that is unresectable, locally advanced or metastatic and one of the following applies: The member has a diagnosis of progressive neuroendocrine tumors of pancreatic origin (PNET) OR The member has a diagnosis of progressive, well differentiated, non-functional neuroendocrine tumors of gastrointestinal or lung.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	Angiomyolipoma and Tuberous Sclerosis Complex (TSC). The member has a diagnosis of renal angiomyolipoma and tuberous sclerosis complex AND Immediate surgery is not required. Metastatic Breast Cancer. The member has a diagnosis of hormone receptor (HR) positive and human epidermal growth factor receptor 2 (HER2) negative advanced or metastatic disease AND the member has been treated with endocrine therapy (e.g. letrozole, anastrozole) AND The member will use Everolimus in combination with exemestane or fulvestrant. Tuberous sclerosis complex (TSC)- associated partial onset seizures [Adults and Pediatrics]: The member has diagnosis of TSC- associated partial onset seizures AND Everolimus tablets for oral suspension is being used as adjunctive therapy.
Part B Prerequisite	0

EXXUA - TABLET, EXTENDED RELEASE 24 HR

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Prolonged QTc interval greater than 450 msec at baseline. Congenital long QT syndrome. Severe hepatic impairment.
Required Medical Information	Major Depressive Disorder: The member must be utilizing Exxua (gepirone) for the treatment of major depressive disorder (MDD) AND The member has had prior therapy, contraindication, or intolerance to any two antidepressants from any of the following categories: generic SSRI (e.g., citalopram, fluoxetine, paroxetine, or sertraline), generic SNRI (e.g., venlafaxine or duloxetine), bupropion OR mirtazapine.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

FANAPT - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Schizophrenia: The member must be utilizing Fanapt for treatment of schizophrenia. The member has tried or cannot use at least 2 of the following: risperidone, olanzapine, quetiapine, ziprasidone, aripiprazole, or lurasidone. Bipolar I Disorder (manic or mixed episode): The member must be utilizing Fanapt for the treatment of bipolar I disorder (manic or mixed episode). The member has tried or cannot use at least 2 of the following: risperidone, olanzapine, quetiapine, ziprasidone, or aripiprazole.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

FANAPT TITRATION PACK A - TABLET, DOSE PACK

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Schizophrenia: The member must be utilizing Fanapt for treatment of schizophrenia. The member has tried or cannot use at least 2 of the following: risperidone, olanzapine, quetiapine, ziprasidone, aripiprazole, or lurasidone. Bipolar I Disorder (manic or mixed episode): The member must be utilizing Fanapt for the treatment of bipolar I disorder (manic or mixed episode). The member has tried or cannot use at least 2 of the following: risperidone, olanzapine, quetiapine, ziprasidone, or aripiprazole.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

FANAPT TITRATION PACK B - TABLET, DOSE PACK

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Schizophrenia: The member must be utilizing Fanapt for treatment of schizophrenia. The member has tried or cannot use at least 2 of the following: risperidone, olanzapine, quetiapine, ziprasidone, aripiprazole, or lurasidone. Bipolar I Disorder (manic or mixed episode): The member must be utilizing Fanapt for the treatment of bipolar I disorder (manic or mixed episode). The member has tried or cannot use at least 2 of the following: risperidone, olanzapine, quetiapine, ziprasidone, or aripiprazole.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

FANAPT TITRATION PACK C - TABLET, DOSE PACK

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Schizophrenia: The member must be utilizing Fanapt for treatment of schizophrenia. The member has tried or cannot use at least 2 of the following: risperidone, olanzapine, quetiapine, ziprasidone, aripiprazole, or lurasidone. Bipolar I Disorder (manic or mixed episode): The member must be utilizing Fanapt for the treatment of bipolar I disorder (manic or mixed episode). The member has tried or cannot use at least 2 of the following: risperidone, olanzapine, quetiapine, ziprasidone, or aripiprazole.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

FARXIGA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Pending CMS Review
Required Medical Information	Type 2 Diabetes Mellitus. The member has a diagnosis of type 2 diabetes mellitus AND the member is using Farxiga (dapagliflozin) for one of the following: to improve glycemic control OR to reduce the risk of hospitalization for heart failure and either established cardiovascular disease or multiple cardiovascular risk factors AND If the member is using for glycemic control, the provider attests member will use Farxiga (dapagliflozin) in combination with diet and exercise AND the member has tried or cannot use generic dapagliflozin tablet. Heart Failure. The member has a diagnosis of heart failure AND the member is using Farxiga (dapagliflozin) to reduce the risk of cardiovascular death, hospitalization for heart failure, and urgent heart failure visit AND the member has tried or cannot use generic dapagliflozin tablet. Chronic Kidney Disease at Risk of Progression. The member has been diagnosed with chronic kidney disease at risk of progression AND the member is using Farxiga (dapagliflozin) to reduce the risk of sustained eGFR decline, end stage kidney disease, cardiovascular death, and hospitalization for heart failure AND the member has tried or cannot use generic dapagliflozin tablet.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

FASENRA PEN - AUTO-INJECTOR (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

felbamate - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Partial-Onset (Focal) Seizures: Member has a diagnosis of partial-onset (focal) seizures AND The provider attests that the members epilepsy is so severe that the risk of aplastic anemia is deemed acceptable in light of the benefits conferred by its use AND Member has normal baseline serum transaminases (AST/ALT) AND Member has been unable to achieve seizure control with at least TWO other drugs for controlling partial-onset seizures (e.g., levetiracetam, lamotrigine, carbamazepine, zonisamide, divalproex, gabapentin or topiramate). Lennox-Gastaut Syndrome: The member has a diagnosis of seizures associated with Lennox-Gastaut syndrome AND Felbamate (Felbate) is prescribed by or in consultation with a specialist (i.e. neurologist, epileptologist) AND The provider attests that the member's epilepsy is so severe that the risk of aplastic anemia is deemed acceptable in light of the benefits conferred by its use AND Member has normal baseline serum transaminases (AST/ALT) AND The member has been unable to achieve seizure control with at least two other drugs used for the treatment of Lennox-Gastaut syndrome (e.g. valproic acid, lamotrigine, rufinamide, topiramate).
Age Restriction	
Prescriber Restriction	Felbamate (Felbatol) is prescribed by or in consultation with a specialist (i.e. neurologist, epileptologist).
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

FETZIMA - CAPSULE,EXTENDED RELEASE 24 HR DOSE PACK

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Major depressive disorder: The member must be utilizing Fetzima for treatment of major depressive disorder. For new starts only: The member has tried or cannot use a serotonin and norepinephrine reuptake inhibitor (SNRI) (e.g., duloxetine, venlafaxine, or desvenlafaxine succinate ER) AND a generic bupropion product (75mg/100mg IR, 100mg/150mg/200mg SR, or 150mg/300mg XL) or mirtazapine.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

FILSPARI - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	FSGS: Member is 8 years of age and older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

fingolimod - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	The member has a diagnosis of a relapsing form of multiple sclerosis, to include relapsing-remitting or active secondary progressive disease OR the member has a diagnosis of a clinically isolated syndrome (CIS).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

FINTEPLA - SOLUTION, ORAL

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Dravet Syndrome: The member has a diagnosis of seizures associated with Dravet Syndrome AND Fintepla (fenfluramine) is prescribed by or in consultation with a specialist (i.e. neurologist, epileptologist) AND The member has been unable to achieve seizure control with at least one other drug used for the treatment of Dravet Syndrome (e.g. clobazam, valproic acid, topiramate). Lennox-Gastaut Syndrome: The member has a diagnosis of seizures associated with Lennox-Gastaut syndrome AND Fintepla (fenfluramine) is prescribed by or in consultation with a specialist (i.e. neurologist, epileptologist) AND The member has been unable to achieve seizure control with at least two other drugs used for the treatment of Lennox-Gastaut syndrome (e.g. valproic acid, lamotrigine, rufinamide, topiramate).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

FIRDAPSE - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	History of seizures (not to be inferred from pharmacy claims)
Required Medical Information	Lambert-Eaton Myasthenic Syndrome (LEMS). The member has a confirmed diagnosis of Lambert-Eaton Myasthenic Syndrome (LEMS) by a specialist (e.g. neurologist) AND The diagnosis is supported by results from a clinical evaluation (e.g. electromyography or the presence of autoantibodies directed against VGCC {voltagegated calcium channels}).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

FIRMAGON - VIAL (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

FIRMAGON KIT W DILUENT SYRINGE - VIAL (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

FORTEO - PEN INJECTOR (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

FOTIVDA - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member experiences disease progression on Fotivda (tivozanib).
Required Medical Information	Relapsed or refractory advanced renal cell carcinoma: The member has a diagnosis of relapsed or refractory advanced renal cell carcinoma AND The member has received two prior systemic therapies (e.g., immuno-oncology checkpoint inhibitors, cabozantinib, axitinib) AND Fotivda (tivozanib) is given as a single agent for subsequent therapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

FRUZAQLA - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member has had disease progression on Fruzaqla (fruquintinib).
Required Medical Information	Metastatic Colorectal Cancer (mCRC). Member has a diagnosis of metastatic colorectal cancer (mCRC) AND Member has been previously treated with the following: fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy AND anti-VEGF therapy (e.g., bevacizumab product) AND anti-EGFR therapy [if verified KRAS/NRAS wild-type (normal) and medically appropriate].
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

GAMUNEX-C - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	Member is younger than 13 years of age for treatment of AIDS/HIV.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	Pending CMS Review
Part B Prerequisite	0

GAVRETO - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member experiences disease progression on RET inhibitors (e.g., pralsetinib, selpercatinib).
Required Medical Information	Non-small cell lung cancer: The member has a diagnosis of metastatic non-small lung cancer AND the disease is documented as RET fusion positive AND Gavreto (pralsetinib) is being used as monotherapy. Thyroid cancer: The member has a diagnosis of metastatic or advanced thyroid cancer AND The disease is documented RET fusion positive AND The disease no longer responds to radioactive iodine treatment (if radioactive iodine is appropriate) AND Gavreto (pralsetinib) is being used as monotherapy for systemic therapy.
Age Restriction	Thyroid Cancer: The member is 12 years of age and older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 Months Duration
Other Criteria	
Part B Prerequisite	0

GAZYVA - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	For Follicular Lymphoma: The length of maintenance therapy exceeds 2 years for low-grade non-Hodgkins lymphoma (e.g. follicular lymphoma, marginal zone lymphoma).
Required Medical Information	Chronic Lymphocytic Leukemia: The member must have a diagnosis of chronic lymphocytic leukemia AND Gazyva (obinutuzumab) will be given as part of a supported combination regimen or as monotherapy. Follicular Lymphoma: The member must have a diagnosis of follicular lymphoma AND One of the following: For first line therapy: Gazyva (obinutuzumab) will be given as part of a supported combination regimen with or without maintenance monotherapy OR For subsequent line of therapy: Gazyva (obinutuzumab) will be given as part of a supported combination regimen or as monotherapy. Lupus Nephritis: The member must have a diagnosis of active lupus nephritis AND The member must be utillizing Gazyva in combination with standard therapy (e.g., corticosteroids with mycophenolate or cyclophosphamide).
Age Restriction	Lupus Nephritis: The member is 18 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

gefitinib - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Non-small cell lung cancer (NSCLC): The member has a diagnosis of metastatic or recurrent non-small cell lung cancer (NSCLC) AND the following applies: The member has a documented epidermal growth factor receptor (EGFR) exon 19 deletion or exon 21 (L858R) substitution mutation AND The member is using Iressa (gefitinib) as monotherapy (without concomitant chemotherapy).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Six month duration
Other Criteria	
Part B Prerequisite	0

GILOTRIF - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Non-small cell lung cancer (NSCLC): The member has a diagnosis of metastatic non-small cell lung cancer (NSCLC) AND the following apply: The member has a documented non-resistant epidermal growth factor receptor (EGFR) mutation (sensitizing EGFR mutation e.g., exon 19 deletion, L861Q, S768I, G719X, L858R) AND The member is using Gilotrif (afatinib) as monotherapy (without concomitant chemotherapy) OR squamous cell histology after disease progression on platinum containing chemotherapy and is using Gilotrif (afatinib) as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Six month duration
Other Criteria	
Part B Prerequisite	0

glatiramer - SYRINGE (ML)

PA Criteria	Criteria Details
Off-Label Uses	NA
Exclusion Criteria	NA
Required Medical Information	The member has a diagnosis of a relapsing form of multiple sclerosis, to include relapsing-remitting or active secondary progressive disease OR the member has a diagnosis of a clinically isolated syndrome (CIS).
Age Restriction	NA
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	NA
Part B Prerequisite	0

glatopa - SYRINGE (ML)

PA Criteria	Criteria Details
Off-Label Uses	NA
Exclusion Criteria	NA
Required Medical Information	The member has a diagnosis of a relapsing form of multiple sclerosis, to include relapsing-remitting or active secondary progressive disease OR the member has a diagnosis of a clinically isolated syndrome (CIS).
Age Restriction	NA
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	NA
Part B Prerequisite	0

glutamine (sickle cell) - POWDER IN PACKET (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

GOMEKLI - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Neurofibromatosis type 1 (NF1): The member has symptomatic plexiform neurofibromas (PN) [neurofibromatosis type 1 (NF1)] AND The disease is not amenable to complete resection.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

HAEGARDA - VIAL (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	The member is being treated by a specialist in hereditary angioedema (i.e. allergist and/or immunologist).
Coverage Duration	Plan Year Duration
Other Criteria	Pending CMS Review
Part B Prerequisite	0

HERNEXEOS - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Non-Small Cell Lung Cancer (NSCLC): The member has a diagnosis of unresectable or metastatic, non-squamous, non-small cell lung cancer AND The disease is documented positive for HER2 (ERBB2) tyrosine kinase domain activating mutations AND Hernexeos (zongertinib) will be given as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

HYRNUO - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Non-Small Cell Lung Cancer (NSCLC): The member has a diagnosis of locally advanced or metastatic non-squamous non-small cell lung cancer (NSCLC) AND The disease is documented positive for HER2 (ERBB2) tyrosine kinase domain (TKD) activating mutations AND The member has received prior systemic therapy (e.g., platinum-based chemotherapy) AND Hyrnuo (sevabertinib) will be given as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

IBRANCE - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

IBTROZI - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	ROS1 Positive NSCLC: The member has diagnosis of locally advanced or metastatic non-small cell lung cancer (NSCLC) AND The member has disease which is ROS-1 rearrangement(s) positive AND The member will be using Ibtrozi (taletrectinib) as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

icatibant - SYRINGE (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	The member is being treated by a specialist in hereditary angioedema (i.e. allergist and/or immunologist).
Coverage Duration	Plan Year Duration
Other Criteria	Pending CMS Review
Part B Prerequisite	0

ICLUSIG - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression while on Iclusig (ponatinib).
Required Medical Information	Chronic Myeloid Leukemia (chronic phase): The member has a diagnosis of chronic phase chronic myeloid leukemia (CML) AND one of the following apply: The member has an intolerance, resistance, or contraindication to at least two available tyrosine kinase inhibitors indicated for the treatment of CML OR The member has a documented T315I mutation. Chronic Myeloid Leukemia (accelerated or blast phase): The member has a diagnosis of accelerated or blast phase chronic myeloid leukemia (CML) AND one of the following apply: There are no other kinase inhibitors indicated OR the member has a documented T315I mutation. Acute Lymphoblastic Leukemia: The member has a diagnosis of Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL) AND one of the following apply: There are no other kinase inhibitors indicated OR The member has a documented T315I mutation OR Iclusig (ponatinib) will be used in combination with chemotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

IDHIFA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member has experienced disease progression while on or following Idhifa(enasidenib).
Required Medical Information	Acute Myeloid Leukemia, Relapsed/Refractory: The member has a diagnosis of acute myeloid leukemia (AML) AND The member has relapsed or refractory disease AND The member has a documented IDH2 mutation AND One of the following applies: The member will be using Idhifa (enasidenib) as monotherapy OR the member will be using Idhifa (enasidenib) as a component of repeating the initial successful induction regimen, if late relapse (relapse occurring later than 12 months). Acute Myeloid Leukemia, Newly diagnosed: The member has a diagnosis of acute myeloid leukemia (AML) AND the member has newly diagnosed disease AND the member is not a candidate for intensive induction therapy due to comorbidities AND the member has a documented IDH2 mutation AND the member will be using Idhifa (enasidenib) as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

imatinib - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Patients that have experienced disease progression while on imatinib.
Required Medical Information	One of the following applies: The member has a diagnosis of Ph+ CML that is newly diagnosed in the chronic phase OR The member has a diagnosis of Ph+ CML that is in accelerated phase or blast crisis OR The member has a diagnosis of Ph+ CML that is in chronic phase after failure of interferon-alpha therapy. Acute lymphoid leukemia (ALL).The member has a diagnosis of Ph+ ALL that is relapsed, refractory, or newly diagnosed and imatinib is being added to consolidation or induction therapy OR the member has a diagnosis of PH+ALL and receiving maintenance therapy. The member has a diagnosis of Kit (CD117)-positive GIST. The member has a diagnosis of Dermatofibrosacrome protuberans (DFSP) that is adjuvant (positive surgical margins following excision) unresectable, recurrent, and/or metastatic. The member has a diagnosis of chronic eosinophilic leukemia or hypereosinophilic syndrome. The member has a diagnosis of MDS or chronic MPD that is associated with platelet-derived growth factor receptor (PDGFR) gene rearrangement.(ex. Chronic myelomonocyte leukemia, atypical chronic myeloid leukemia, juvenile myelomonocyte leukemia).The member has a diagnosis of aggressive systemic mastocytosis. The member must not harbor the D816v mutation of C-kit. Melanoma. The member has a diagnosis of unresectable melanoma with activating mutation of C-kit. Imatinib will be used as single agent in subsequent therapy.
Age Restriction	[Pediatric] Chronic myeloid leukemia (CML) and [Pediatric] Acute Lymphoid Luekemia (ALL)- The member is at least one year of age.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	Pediatric indications: The patient has a diagnosis of Philadelphia chromosome positive (Ph+) CML that is newly diagnosed in chronic phase OR The patient has a diagnosis of Ph+ CML that is in chronic phase with disease recurrence after stem cell transplant OR The patient has a diagnosis of Ph+ CML that is in chronic phase after failure of interferon-alpha therapy.Acute Lymphoid Luekemia (ALL). The member is newly diagnosed with Ph+ ALL AND the member will be using imatinib in combination with chemotherapy.
Part B Prerequisite	0

IMBRUVICA - SUSPENSION, ORAL (FINAL DOSE FORM)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression while on or following Imbruvica (ibrutinib).
Required Medical Information	Chronic Lymphocytic Leukemia (CLL)/Small Lymphocytic Lymphoma (SLL): The member has a diagnosis of Chronic Lymphocytic Leukemia (CLL)/Small Lymphocytic Lymphoma (SLL). Waldenstroms Macroglobulinemia: The member has a diagnosis of Waldenstroms macroglobulinemia AND The member is using Imbruvica (ibrutinib) as monotherapy or in combination with a rituximab product. Marginal Zone Lymphoma: The member has a diagnosis of marginal zone lymphoma AND The member is using Imbruvica (ibrutinib) as second line or subsequent for refractory or progressive disease AND The member is using Imbruvica (ibrutinib) as monotherapy. Chronic Graft Versus Host Disease: The member has a diagnosis of chronic graft versus host disease (cGVHD) AND The member has been unable to achieve treatment goals with at least one prior line of systemic therapy (e.g. corticosteroids).
Age Restriction	cGVHD: Member age is 1 year or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

IMFINZI - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	All indications except Bladder Cancer and GE/GEJC: Disease progression while on or following anti-PD-1/PD-L1 therapy [e.g., Opdivo (nivolumab), Keytruda (pembrolizumab), Tecentriq (atezolizumab), Imfinzi (durvalumab)]. For unresectable stage III NSCLC, member has not exceeded a maximum of twelve (12) months of therapy. For perioperative NSCLC, for adjuvant treatment, and GE/GEJC member has not exceeded a maximum of 12 cycles after surgery.
Required Medical Information	Non-Small Cell Lung Cancer (NSCLC) (unresectable): Member has diagnosis of unresectable stage III non-small cell lung cancer (NSCLC) AND Imfinzi (durvalumab) will be used as consolidation therapy AND Member has not experienced disease progression following completion of concurrent platinum-based chemotherapy and radiation AND Imfinzi (durvalumab) will be used as monotherapy. Non-Small Cell Lung Cancer (NSCLC) (metastatic): Member has a diagnosis of metastatic non-small cell lung cancer (NSCLC) AND Member has no documented EGFR or ALK genomic tumor aberrations AND Member will be using Imfinzi (durvalumab) in combination with Imjudo (tremelimumab-actl) and platinum-based chemotherapy as first-line therapy. Small Cell Lung Cancer: The member has a diagnosis of extensive-stage small cell lung cancer AND Imfinzi (durvalumab) will be given in combination with etoposide and carboplatin as first line therapy followed by maintenance therapy with Imfinzi (durvalumab) as a single agent. Biliary Tract Cancer: the member has a diagnosis of locally advanced or metastatic biliary tract cancer AND the member will be using Imfinzi (durvalumab) in combination with gemcitabine and cisplatin. Hepatocellular Carcinoma: The member has a diagnosis of locally advanced unresectable and/or metastatic hepatocellular carcinoma AND The member will be using Imfinzi (durvalumab) in combination with Imjudo (tremelimumab-actl) as first-line therapy only. Endometrial cancer: The member has a diagnosis of primary advanced or recurrent endometrial cancer AND the disease is mismatch repair deficient (dMMR) AND Imfinzi (durvalumab) will be used in combination with carboplatin and paclitaxel followed by Imfinzi (durvalumab) as a single agent.
Age Restriction	
Prescriber Restriction	Licensed Practitioner.
Coverage Duration	6 months Duration.

Other Criteria	Non-small cell lung cancer (perioperative): The member has a diagnosis of non-small cell lung cancer AND The disease is resectable (tumors greater than or equal to 4 cm and/or node positive) AND The disease does not have known epidermal growth factor receptor (EGFR) mutations or anaplastic lymphoma kinase (ALK) rearrangements AND Imfinzi (durvalumab) will be used in combination with chemotherapy as neoadjuvant treatment followed by Imfinzi (durvalumab) as monotherapy as adjuvant treatment after surgery. Small Cell Lung Cancer (Limited stage): The member has a diagnosis of limited-stage small cell lung cancer AND The member's disease has not progressed following concurrent platinum-based chemotherapy and radiation therapy AND Imfinzi (durvalumab) will be given as monotherapy. Bladder Cancer (muscle invasive): The member has a diagnosis of muscle invasive bladder cancer (MIBC) AND The member will be using Imfinzi (durvalumab) in combination with gemcitabine and cisplatin as neoadjuvant treatment, followed by a single agent Imfinzi (durvalumab) as adjuvant treatment following radical cystectomy. Gastric or gastroesophageal junction adenocarcinoma (GE/GEJC): The member has a diagnosis of resectable gastric or gastroesophageal junction adenocarcinoma AND Imfinzi (durvalumab) will be used in combination with fluorouracil, leucovorin, oxaliplatin and docetaxel (FLOT) as neoadjuvant and adjuvant treatment, followed by Imfinzi (durvalumab) monotherapy as part of remaining adjuvant treatment. Bladder Cancer (non-muscle invasive): The member has a diagnosis of high-risk non-muscle invasive bladder cancer (High risk criteria may include one of the following: T1 tumor, grade 3/high grade tumor, carcinoma in situ or multiple and recurrent and large tumors.) AND Imfinzi (durvalumab) will be used in combination with Bacillus Calmette-Guerin (BCG) vaccine AND The member is BCG naive defined as no prior intravesical BCG vaccine or greater than 2 years since last exposure.
Part B Prerequisite	0

IMJUDO - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Disease progression on Imjudo (tremelimumab-actl)
Required Medical Information	Hepatocellular carcinoma (HCC): The member has diagnosis of unresectable hepatocellular carcinoma AND The member will be given Imjudo (tremelimumab-actl) in combination with Imfinzi (durvalumab). Non-small cell lung cancer (NSCLC): The member has a diagnosis of metastatic, advanced, recurrent non-small cell lung cancer AND NSCLC does not express sensitizing genomic tumor aberrations (e.g., EGFR, ALK) AND The member will be given Imjudo (tremelimumab-actl) in combination with Imfinzi (durvalumab) and platinum based chemotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Six month duration
Other Criteria	
Part B Prerequisite	0

IMKELDI - SOLUTION, ORAL

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Patients that have experienced disease progression while on imatinib.
Required Medical Information	One of the following applies: The member has a diagnosis of Ph+ CML that is newly diagnosed in the chronic phase OR The member has a diagnosis of Ph+ CML that is in accelerated phase or blast crisis OR The member has a diagnosis of Ph+ CML that is in chronic phase after failure of interferon-alpha therapy. Acute lymphoid leukemia (ALL). The member has a diagnosis of Ph+ ALL that is relapsed, refractory, or newly diagnosed and imatinib is being added to consolidation or induction therapy OR the member has a diagnosis of PH+ALL and receiving maintenance therapy. The member has a diagnosis of Kit (CD117)-positive GIST. The member has a diagnosis of Dermatofibrosacrome protuberans (DFSP) that is adjuvant (positive surgical margins following excision) unresectable, recurrent, and/or metastatic. The member has a diagnosis of chronic eosinophilic leukemia or hypereosinophilic syndrome. The member has a diagnosis of MDS or chronic MPD that is associated with platelet-derived growth factor receptor (PDGFR) gene rearrangement.(ex. Chronic myelomonocyte leukemia, atypical chronic myeloid leukemia, juvenile myelomonocyte leukemia).The member has a diagnosis of aggressive systemic mastocytosis. The member must not harbor the D816v mutation of C-kit. Melanoma. The member has a diagnosis of unresectable melanoma with activating mutation of C-kit. Imatinib will be used as single agent in subsequent therapy.
Age Restriction	[Pediatric] Chronic myeloid leukemia (CML) and [Pediatric] Acute Lymphoid Luekemia (ALL) - The member is at least one year of age.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	Pediatric indications: The patient has a diagnosis of Philadelphia chromosome positive (Ph+) CML that is newly diagnosed in chronic phase OR The patient has a diagnosis of Ph+ CML that is in chronic phase with disease recurrence after stem cell transplant OR The patient has a diagnosis of Ph+ CML that is in chronic phase after failure of interferon-alpha therapy.Acute Lymphoid Luekemia (ALL). The member is diagnosed with Ph+ ALL AND the member will be using imatinib in combination with chemotherapy. For Imkeldi requests (all indications): Members must have an intolerance or contraindication with generic imatinib.

Part B Prerequisite	0

INCRELEX - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The bone epiphyses are closed.
Required Medical Information	Member has a diagnosis of GH gene deletion with development of neutralizing antibodies to GH OR The patient has a diagnosis of severe primary IGF-1 deficiency defined by:height standard deviation score below or equal -3.0 and basal IGF-1 standard deviation score below or equal -3.0 and normal or elevated growth hormone.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

INLURIYO - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Estrogen Receptor (ER) Positive Breast Cancer: The member has a diagnosis of ER-positive, HER2-negative advanced or metastatic breast cancer AND The breast cancer has a documented ESR1-mutation AND The member has experienced progressive disease following at least one prior line of endocrine therapy (e.g., aromatase inhibitor, CDK 4/6 inhibitor).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

INLYTA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on Inlyta (axitinib).
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

INQOVI - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression on hypomethylators (e.g. azacitidine, decitabine).
Required Medical Information	Myelodysplastic Syndromes - Chronic Myelomonocytic Leukemia: The member has a diagnosis of myelodysplastic syndromes (MDS), including previously treated and untreated, de novo or secondary MDS OR chronic myelomonocytic leukemia (CMML) AND the member will be using Inqovi (decitabine and cedazuridine) as a single agent.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

INREBIC - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on Inrebic (fedratinib).
Required Medical Information	Myelofibrosis: The member has a diagnosis of primary myelofibrosis or secondary myelofibrosis (i.e. post-polycythemia vera or post-essential thrombocythemia) AND The member has one of the following risk categories, as defined by accepted risk stratification tools for myelofibrosis (e.g. International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or DIPSS-PLUS): Intermediate-2 risk disease OR High-risk disease AND the member will be using Inrebic (fedratinib) as monotherapy AND The member has a medical reason as to why Jakafi (ruxolitinib) cannot be used.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ipratropium bromide - HFA AEROSOL WITH ADAPTER (GRAM)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Chronic Obstructive Pulmonary Disease (COPD): Diagnosis of Chronic Obstructive Pulmonary Disease (COPD), including chronic bronchitis and emphysema. The member has had previous treatment, contraindication, or intolerance to Spiriva (i.e. Spiriva Handihaler, Spiriva Respimat).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ITOVEBI - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression on a PIK3CA inhibitor (e.g., alpelisib).
Required Medical Information	Metastatic Breast Cancer - PIK3CA-mutated: The member has a diagnosis of advanced or metastatic hormone receptor (HR)-positive and human epidermal growth factor receptor 2 (HER2)-negative breast cancer AND The disease is documented as PIK3CA-mutated, as detected by an FDA-test AND The member has endocrine resistant disease [i.e., disease progression on or after adjuvant endocrine therapy (e.g., anastrozole, tamoxifen)] AND Itovebi (inavolisib) will be given in combination with Ibrance (palbociclib) and fulvestrant.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

IWILFIN - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member has had prior progression on Iwilfin (eflornithine). Member has been on therapy for more than two years.
Required Medical Information	High-Risk Neuroblastoma (HRNB): Member has diagnosis of high-risk neuroblastoma AND Member has had at least a partial response (or better) to prior multiagent, multimodality therapy including anti-GD2 immunotherapy (e.g., dinutuximab).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

JAKAFI - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on Jakafi (ruxolitinib).
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

JAKAFI XR - TABLET, EXTENDED RELEASE 24 HR

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on Jakafi (ruxolitinib).
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

JAYPIRCA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Mantle cell lymphoma: The member has a diagnosis of mantle cell lymphoma AND the member has relasped or refractory disease AND the member has received at least two prior lines of systemic therapy, including a BTK inhibitor AND the member will be using Jaypirca (pirtobrutinib) as monotherapy. Chronic Lymphocytic Leukemia (CLL)/Small Lymphocytic Lymphoma (SLL): The member has a diagnosis of chronic lymphocytic leukemia (CLL) or small lymphocytic lymphoma (SLL) AND the member has previously received a covalent BTK inhibitor (e.g., Imbruvica (ibrutinib), Calquence (acalabrutinib) or Brukinsa (zanubrutinib)) AND the member will be using Jaypirca (pirtobrutinib) as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

JEMPERLI - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Disease progression while on or following prior anti-PD-1/PD-L1 therapy (e.g.,nivolumab, pembrolizumab).
Required Medical Information	Endometrial cancer (2nd line or beyond): The member has diagnosis of recurrent or advanced endometrial cancer AND The member has documented dMMR endometrial cancer AND The member has progressed on prior platinum containing regimen AND Jemperli (dostarlimab-gxly) is administered as monotherapy as subsequent therapy. Solid tumors (dMMR): The member has a diagnosis of unresectable or metastatic documented mismatch repair deficient (d-MMR) solid tumors AND the member has disease that has progressed on prior therapy with no alternative treatments AND Jemperli (dostarlimab-gxly) is administered as monotherapy. Endometrial cancer (frontline therapy): The member has diagnosis of primary advanced or recurrent endometrial cancer AND Jemperli (dostarlimab-gxly) will be used in combination with carboplatin and paclitaxel, followed by maintenance therapy as a single agent.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 months duration. Endometrial cancer (frontline therapy) only: Plan Year Duration.
Other Criteria	
Part B Prerequisite	0

JIDEYTRO - TABLET

PA Criteria	Criteria Details
Off-Label Uses	Pending CMS Review
Exclusion Criteria	Pending CMS Review
Required Medical Information	Pending CMS Review
Age Restriction	Pending CMS Review
Prescriber Restriction	Pending CMS Review
Coverage Duration	Pending CMS Review
Other Criteria	Pending CMS Review
Part B Prerequisite	Pending CMS Review

JYLAMVO - SOLUTION, ORAL

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Acute Lymphoblastic Leukemia (ALL): The member has a diagnosis of acute lymphoblastic leukemia AND The member will be using Jylamvo (methotrexate) as part of combination chemotherapy maintenance regimen AND The member has had previous treatment, intolerance or documented difficulty swallowing generic methotrexate tablets. Mycosis Fungoides: The member has a diagnosis of mycosis fungoides (cutaneous T-cell lymphoma) AND The member will be using Jylamvo (methotrexate) as a single agent or as part of combination chemotherapy regimen AND The member has had previous treatment, intolerance or documented difficulty swallowing generic methotrexate tablets. Non-Hodgkin lymphomas: The member has a diagnosis of non-Hodgkin lymphoma AND The member will be using Jylamvo (methotrexate) as part of a metronomic combination chemotherapy regimen AND The member has had previous treatment, intolerance or documented difficulty swallowing generic methotrexate tablets. Rheumatoid Arthritis: The member has a diagnosis of Rheumatoid Arthritis AND The member has had previous treatment, intolerance or documented difficulty swallowing generic methotrexate tablets. Psoriasis: The member has a diagnosis of severe psoriasis AND The member has had previous treatment, intolerance or documented difficulty swallowing generic methotrexate tablets. Polyarticular Juvenile Idiopathic Arthritis: The member has a diagnosis of polyarticular juvenile idiopathic arthritis AND The member has had previous treatment, intolerance or documented difficulty swallowing generic methotrexate tablets. B vs D coverage determination may also be required.
Age Restriction	Mycosis Fungoides, Non-Hodgkin lymphomas, Rheumatoid Arthritis, and Psoriasis: The member must be 18 years or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

KERENDIA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Chronic kidney disease associated with type 2 diabetes: The member has a diagnosis of chronic kidney disease (CKD) associated with type 2 diabetes (T2D) AND The member is currently receiving, unless contraindicated or intolerant, the maximally tolerated dose of: Either an angiotensin-converting enzyme inhibitor (e.g. Lisinopril) OR an angiotensin receptor blocker (e.g. losartan). Heart Failure: The member has a diagnosis of heart failure AND has a left ventricular ejection fraction greater than or equal to 40%.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

KESIMPTA PEN - PEN INJECTOR (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	The member has a diagnosis of a relapsing form of multiple sclerosis, to include relapsing-remitting or active secondary progressive disease OR the member has a diagnosis of a clinically isolated syndrome (CIS).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

KEYTRUDA - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Disease progression while on or following prior anti-PD-1/PD-L1 therapy (e.g., nivolumab, atezolizumab) (applicable to all indications except Head and Neck Squamous Cell Carcinoma perioperative). Member requiring urgent cytoreductive therapy (applicable to PMBCL only). Member not to exceed one year of total adjuvant treatment (applicable to melanoma, NSCLC, and renal cell carcinoma only). Total treatment duration of Keytruda (pembrolizumab) is more than 12 months (Head and Neck Squamous Cell Carcinoma perioperative only).
Required Medical Information	Melanoma: unresectable or metastatic melanoma OR melanoma OR stage IIB, IIC, or III melanoma AND as monotherapy for adjuvant treatment after complete resection with involvement of lymph node(s). NSCLC-1st Line: metastatic NSCLC AND 1 of the following applies: disease with PD-L1 expression [TPS greater than or equal to 1%] with no EGFR or ALK genomic tumor aberrations and as 1st line AND tumor expresses PD-L1 as determined by an FDA-approved test AND used as monotherapy OR nonsquamous histology with no EGFR or ALK genomic tumor aberrations and in combo with pemetrexed and carboplatin or cisplatin as 1st line therapy followed by Keytruda maintenance in combo with pemetrexed OR squamous histology and used in combo with carboplatin and paclitaxel or Abraxane as 1st line followed by Keytruda maintenance OR stage III NSCLC and not candidate for surgical resection or definitive chemoradiation AND PD-L1 expression with no EGFR or ALK genomic tumor aberrations and as 1st line AND Tumor expresses PD-L1 as determined by an FDA-approved test AND as monotherapy. NSCLC-Subsequent: metastatic NSCLC AND progression on or following chemo and EGFR inhibitor, if EGFR mutation positive or ALK inhibitor, if ALK positive AND Tumor expresses PD-L1 as determined by an FDA-approved test AND as monotherapy. Head-Neck Cancer: recurrent or metastatic non-nasopharyngeal head and neck squamous cell carcinoma AND 1 of following: disease progression on platinum-containing chemo and as monotherapy OR in combo with platinum and 5-FU for 1st line treatment OR monotherapy in 1st line and disease expresses CPS score greater than or equal to 1 as detected by an FDA-approved test. Hodgkin's Lymphoma-Adult: as monotherapy for refractory or relapsed disease. Hodgkin's Lymphoma-Peds: monotherapy and 1 of following: Refractory disease OR Relapsed after 2 or more lines of prior therapy. CSCC: recurrent or metastatic CSCC AND disease is not amenable to curative surgery or radiation AND used as a monotherapy.
Age Restriction	Stage IIB, IIC, or III melanoma - member is 12 years of age or older.

KEYTRUDA - VIAL (ML)

Prescriber Restriction	Licensed Practitioner
Coverage Duration	NSCLC (Neoadjuvant therapy): 3 months duration. All other indications: 6 months duration.

KEYTRUDA - VIAL (ML)

Other Criteria	MSI-High/d-MMR Solid tumors: unresectable or metastatic documented microsatellite instability-high or mismatch repair deficient solid tumors (excluding pediatric patients with MSI-H central nervous system cancers) AND 1 of the following: disease that progressed on prior therapy with no alternative treatments and as monotherapy OR colorectal cancer AND 1 of the following: Keytruda as monotherapy and as subsequent therapy after progression on fluoropyrimidine, oxaliplatin, and irinotecan or 1st line as monotherapy in unresectable or metastatic colorectal cancer. Urothelial Cancer: locally advanced or metastatic urothelial cancer AND 1 of the following: initial therapy in members ineligible to receive platinum containing chemo OR as subsequent monotherapy after disease progression within 12 months of neoadjuvant or adjuvant chemo OR in combination with Padcev (enfortumab vedotin-ejfv) in the front-line setting. Cervical Cancer: recurrent or metastatic cervical cancer AND disease progression on or after chemo AND disease expresses CPS score greater than or equal to 1 as determined by an FDA approved test AND as monotherapy OR persistent, recurrent, or metastatic cervical cancer AND cancer express CPS score of greater than or equal to 1 as determined by an FDA approved test and will be used with chemo, with or without bevacizumab, as 1st line therapy OR diagnosis of FIGO 2014 Stage III-IVA cervical cancer AND will be used with chemo, followed by monotherapy. Primary Mediastinal Large B-Cell Lymphoma [Adults and pediatrics]: relapsed or refractory disease after 2 or more prior lines of treatment AND as monotherapy. Merkel cell carcinoma-Adult and pediatric: recurrent locally advanced or metastatic merkel cell carcinoma AND as monotherapy. HCC: has prior therapy with a 1st line therapy (e.g., sorafenib) AND as monotherapy. RCC: advanced or metastatic RCC AND in combo with Inlyta or Lenvima as 1st line therapy. Esophageal Cancer: recurrent locally advanced or metastatic squamous cell carcinoma of the esophagus AND disease expresses PD-L1 as determined by an FDA approved test AND given as subsequent therapy as a single agent. Endometrial cancer: metastatic or recurrent endometrial cancer AND not MSI-H or pMMR as determined by an FDA approved test AND not candidate for surgery or radiation AND has experienced disease progression on prior systemic therapy AND given in combo with Lenvima as subsequent therapy. NMIBC with carcinoma in situ AND ineligible for or has elected not to undergo cystectomy AND has BCG-unresponsive disease, defined as persistent or recurrent high-grade bladder cancer within 6 months of intravesical BCG therapy AND as monotherapy. TMB-H Solid tumors: unresectable or metastatic solid tumors with documented TMB-H [greater than or equal to 10 mutations/megabase] (excluding pediatric patients with TMB-H central nervous system cancers) AND disease progressed on prior therapy with no alternative treatments AND given as monotherapy.
Part B Prerequisite	0

KEYTRUDA QLEX - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Disease progression while on or following prior anti-PD-1/PD-L1 therapy (e.g., nivolumab, atezolizumab) (applicable to all indications except Head and Neck Squamous Cell Carcinoma perioperative). Member not to exceed one year of total adjuvant treatment (applicable to melanoma, NSCLC, and renal cell carcinoma only). Total treatment duration of Keytruda (pembrolizumab) is more than 12 months (Head and Neck Squamous Cell Carcinoma perioperative only).
Required Medical Information	Melanoma: unresectable or metastatic melanoma OR melanoma OR stage IIB, IIC, or III melanoma AND as monotherapy for adjuvant treatment after complete resection with involvement of lymph node(s). NSCLC-1st Line: metastatic NSCLC AND 1 of the following applies: disease with PD-L1 expression [TPS greater than or equal to 1%] with no EGFR or ALK genomic tumor aberrations and as 1st line AND tumor expresses PD-L1 as determined by an FDA-approved test AND used as monotherapy OR nonsquamous histology with no EGFR or ALK genomic tumor aberrations and in combo with pemetrexed and carboplatin or cisplatin as 1st line therapy followed by pembrolizumab product maintenance in combo with pemetrexed OR squamous histology and used in combo with carboplatin and paclitaxel or Abraxane as 1st line followed by pembrolizumab product maintenance OR stage III NSCLC and not candidate for surgical resection or definitive chemoradiation AND PD-L1 expression with no EGFR or ALK genomic tumor aberrations and as 1st line AND Tumor expresses PD-L1 as determined by an FDA-approved test AND as monotherapy. NSCLC-Subsequent: metastatic NSCLC AND progression on or following chemo and EGFR inhibitor, if EGFR mutation positive or ALK inhibitor, if ALK positive AND Tumor expresses PD-L1 as determined by an FDA-approved test AND as monotherapy. Head-Neck Cancer: recurrent or metastatic non-nasopharyngeal head and neck squamous cell carcinoma AND 1 of following: disease progression on platinum-containing chemo and as monotherapy OR in combo with platinum and 5-FU for 1st line treatment OR monotherapy in 1st line and disease expresses CPS score greater than or equal to 1 as detected by an FDA-approved test. CSCC: recurrent or metastatic CSCC AND disease is not amenable to curative surgery or radiation AND used as a monotherapy.
Age Restriction	Stage IIB, IIC, or III melanoma - member is 12 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	NSCLC (Neoadjuvant therapy): 3 months duration. All other indications: 6 months duration.

KEYTRUDA QLEX - VIAL (ML)

Other Criteria	MSI-High/d-MMR Solid tumors: unresectable or metastatic documented microsatellite instability-high or mismatch repair deficient solid tumors (excluding pediatric patients with MSI-H central nervous system cancers) AND 1 of the following: disease that progressed on prior therapy with no alternative treatments and as monotherapy OR colorectal cancer AND 1 of the following: Keytruda as monotherapy and as subsequent therapy after progression on fluoropyrimidine, oxaliplatin, and irinotecan or 1st line as monotherapy in unresectable or metastatic colorectal cancer. Urothelial Cancer: locally advanced or metastatic urothelial cancer AND 1 of the following: initial therapy in members ineligible to receive platinum containing chemo OR as subsequent monotherapy after disease progression within 12 months of neoadjuvant or adjuvant chemo OR in combination with Padcev (enfortumab vedotin-ejfv) in the front-line setting. Cervical Cancer: recurrent or metastatic cervical cancer AND disease progression on or after chemo AND disease expresses CPS score greater than or equal to 1 as determined by an FDA approved test AND as monotherapy OR persistent, recurrent, or metastatic cervical cancer AND cancer express CPS score of greater than or equal to 1 as determined by an FDA approved test and will be used with chemo, with or without bevacizumab, as 1st line therapy OR diagnosis of FIGO 2014 Stage III-IVA cervical cancer AND will be used with chemo, followed by monotherapy. Primary Mediastinal Large B-Cell Lymphoma [Adults and pediatrics]: relapsed or refractory disease after 2 or more prior lines of treatment AND as monotherapy. Merkel cell carcinoma-Adult and pediatric: recurrent locally advanced or metastatic merkel cell carcinoma AND as monotherapy. HCC: has prior therapy with a 1st line therapy (e.g., sorafenib) AND as monotherapy. RCC: advanced or metastatic RCC AND in combo with Inlyta or Lenvima as 1st line therapy. Esophageal Cancer: recurrent locally advanced or metastatic squamous cell carcinoma of the esophagus AND disease expresses PD-L1 as determined by an FDA approved test AND given as subsequent therapy as a single agent. Endometrial cancer: metastatic or recurrent endometrial cancer AND not MSI-H or pMMR as determined by an FDA approved test AND not candidate for surgery or radiation AND has experienced disease progression on prior systemic therapy AND given in combo with Lenvima as subsequent therapy. NMIBC with carcinoma in situ AND ineligible for or has elected not to undergo cystectomy AND has BCG-unresponsive disease, defined as persistent or recurrent high-grade bladder cancer within 6 months of intravesical BCG therapy AND as monotherapy. TMB-H Solid tumors: unresectable or metastatic solid tumors with documented TMB-H [greater than or equal to 10 mutations/megabase] (excluding pediatric patients with TMB-H central nervous system cancers) AND disease progressed on prior therapy with no alternative treatments AND given as monotherapy.
Part B Prerequisite	0

KISQALI - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Early Breast Cancer - Combination with Aromatase Inhibitor: Member exceeds three years of total Kisqali (ribociclib) based treatment.
Required Medical Information	Breast Cancer-Combination with Aromatase Inhibitors. The member has a diagnosis of advanced or metastatic hormone receptor (HR)-positive and human epidermal growth factor receptor 2 (Her2neu)-negative breast cancer AND the member will be using Kisqali (ribociclib) in combination with an aromatase inhibitor (e.g., letrozole) as first line endocrine therapy. Breast Cancer- Combination with fulvestrant. The member has a diagnosis of advanced or metastatic hormone receptor-positive and human epidermal growth factor receptor 2-negative breast cancer AND the member will be using Kisqali (ribociclib) in combination with fulvestrant. Early Breast Cancer-Combination with Aromatase Inhibitor: The member has a diagnosis of hormone receptor (HR) - positive and human epidermal growth factor receptor 2 (HER2) - negative, stage II or III early breast cancer at high risk of recurrence AND the member will be using Kisqali (ribociclib) in combination with an aromatase inhibitor (e.g., letrozole) as adjuvant treatment.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

KOMZIFTI - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression while on or following a menin inhibitor (e.g., Komzifti (ziftomenib), Revuforj (revumenib)).
Required Medical Information	Acute myeloid leukemia (AML): The member has a diagnosis of relapsed or refractory AML AND The disease is confirmed to have a susceptible nucleophosmin 1 (NPM1) mutation AND The member has no other satisfactory treatment alternatives.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

KOSELUGO - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member experienced disease progression on Koselugo (selumetinib)
Required Medical Information	Neurofibromatosis type 1: The member has a diagnosis of neurofibromatosis type 1 which is symptomatic, inoperable plexiform neurofibromas and Koselugo (selumetinib) is given as a monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

KRAZATI - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member experienced disease progression on KRAS G12C inhibitor (e.g., sotorasib, adagrasib).
Required Medical Information	Non-small cell lung cancer (NSCLC): The member has a diagnosis of locally advanced or metastatic NSCLC AND the NSCLC has documented KRAS G12C mutation AND the member has experienced disease progression on one prior therapy AND Krazati (adagrasib) will be given as monotherapy. Colorectal Cancer (CRC): The member has a diagnosis of locally advanced or metastatic colorectal cancer (CRC) AND the CRC has a documented KRAS G12C mutation AND the member has received prior systemic treatment with chemotherapy [fluoropyrimidine-based chemotherapy (e.g., 5-fluorouracil, capecitabine), oxaliplatin, irinotecan] AND Krazati (adagrasib) will be given in combination with Erbitux (cetuximab).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 months duration
Other Criteria	
Part B Prerequisite	0

lanreotide - SYRINGE (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Diagnosis of acromegaly. The member has a diagnosis of acromegaly. The member has had an inadequate response to or cannot be treated with traditional therapies including surgury and/or radiation therapy. Gastroenteropancreatic Neuroendocrine Tumors (GEP-NETs): The member has a diagnosis of unresectable, well- or moderately-differentiated, locally advanced, or metastatic gastroenteropancreatic neuroendocrine tumors. Carcinoid Syndrome: The member has a diagnosis of carcinoid syndrome with symptoms of flushing and/or diarrhea.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

lapatinib - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on Tykerb (lapatinib).
Required Medical Information	Breast Cancer. The member has a diagnosis of HER2(human epidermal growth factor receptor2) positive advanced or metastatic breast cancer AND The member had prior therapy, contraindication, or intolerance with an anthracycline (e.g. doxorubicin) and a taxane (e.g. paclitaxel) OR The member has a diagnosis of HER2 positive metastatic breast cancer AND Used as first line treatment in combination with an aromatase inhibitor (Femara (letrozole), Arimidex (anastrozole) or Aromasin (exemestane)) for hormone receptor positive disease.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 month duration
Other Criteria	
Part B Prerequisite	0

LAZCLUZE - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

lenalidomide - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on lenalidomide.
Required Medical Information	Myelodysplastic Syndromes (MDS) with 5Q deletion. The member has a diagnosis of low- or intermediate-1 risk myelodysplastic syndrome (MDS) AND the member has a documented deletion 5q chromosomal abnormality AND the member has one of the following: Symptomatic Anemia (e.g. chronic fatigue, malaise) OR Transfusion-dependent anemia OR Anemia that is not controlled with erythroid stimulating agent. Myelodysplastic Syndromes (MDS) without 5Q deletion (non-5Q deletion). The member has a diagnosis of low- or intermediate-1 risk myelodysplastic syndrome (MDS) AND the member has no documented 5q deletion abnormality AND the member has one of the following: Symptomatic Anemia (e.g. chronic fatigue, malaise) OR Transfusion-dependent anemia OR Anemia that is not controlled with erythroid stimulating agent. Multiple Myeloma. Diagnosis of active Multiple Myeloma or Systemic Light Chain Amyloidosis. Primary induction OR for relapsed/refractory disease, lenalidomide therapy should be utilized in conjunction with dexamethasone if no contraindication. As maintenance therapy in patients with multiple myeloma following autologous hematopoietic stem cell transplantation. Chronic Lymphoid Leukemia. Diagnosis of relapsed or refractory Chronic Lymphocytic Leukemia (CLL).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	Hodgkin Lymphoma: The member has a diagnosis of classical Hodgkin lymphoma AND The member has relapsed or refractory disease AND The member will be using for third line or subsequent systemic therapy. Non-Hodgkin Lymphoma: The member has one of the following diagnoses: AIDS-related B-cell lymphoma, diffuse large B-cell lymphoma, mucosa-associated lymphoid tissue (MALT) lymphoma [either gastric or nongastric], marginal zone lymphoma, or mantle cell lymphoma AND The member has relapsed or refractory disease OR The member has a diagnosis of follicular lymphoma AND The member will use as first line therapy or for relapsed or refractory disease.
Part B Prerequisite	0

LENVIMA - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on Lenvima (lenvatinib).
Required Medical Information	Thyroid Cancer: The member has a diagnosis of locally recurrent or metastatic, progressive differentiated thyroid cancer (i.e. papillary carcinoma, follicular carcinoma or oncocytic carcinoma) AND The disease no longer responds to radioactive iodine treatment (if radioactive iodine is appropriate) AND Lenvima (lenvatinib) will be used as a single agent. Renal Cell Carcinoma- second line therapy: The member has a diagnosis of advanced renal cell carcinoma AND the member is using in combination with everolimus AND The member has a medical reason as to why Cabometyx (cabozantinib) cannot be initiated or continued as second line therapy. Hepatocelluar Carcinoma: The member has a diagnosis of unresectable carcinoma AND Lenvima (lenvatinib) will be given as a single agent. Endometrial cancer: The member has a diagnosis of advanced, metastatic or recurrent endometrial cancer AND The disease is not MSI-H or The disease is documented mismatch repair proficient (pMMR) as determined by an FDA-approved test AND The member is not a candidate for curative surgery or radiation AND The member has experienced disease progression on prior or following systemic therapy in any setting AND Lenvima (levantinib) will be given in combination with a pembrolizumab product as subsequent therapy. Renal cell carcinoma- first line therapy: The member has a diagnosis of advanced renal cell carcinoma AND Lenvima (levatinib) will be given in combination with a pembrolizumab product (Keytruda, Keytruda Qlex) as first line therapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

leuprolide acetate (3 month) - VIAL (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Prostate Cancer: Diagnosis of advanced prostate cancer or at risk for disease recurrence. Endometriosis: Diagnosis of endometriosis. Fibroids (Uterine leiomyomata). The patient must have a diagnosis of anemia due to uterine leiomymota. Central Precocious Puberty: Pediatric member with a diagnosis of central precocious puberty (idiopathic or neurogenic). Invasive Breast Cancer: The patient has a diagnosis of hormone responsive (ER and/or PR +) invasive breast cancer AND The patient must be pre or perimenopausal. Recurrent Ovarian Cancer: Patient has recurrent ovarian cancer (epithelial cell cancer, fallopian tube cancer, primary peritoneal cancer or ovarian stromal tumor).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration for all except for Endometriosis: 6 months and Fibroids: 3 months
Other Criteria	
Part B Prerequisite	0

levoleucovorin calcium - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members with pernicious anemia or megaloblastic anemia secondary to the lack of vitamin B12.
Required Medical Information	Osteosarcoma: The member is being treated with high dose methotrexate for osteosarcoma. The patient has been treated with leucovorin calcium and has experienced documented side effects either due to lack of leucovorin calcium efficacy OR due to leucovorin calcium formulation necessitating a change in therapy. Impaired methotrexate elimination or inadvertent over dosage of folic acid antagonists. The patient has been treated with methotrexate or other folic acid antagonist and is currently exhibiting signs of toxicity likely due to aforementioned therapy. The patient has been treated with leucovorin calcium and has experienced documented side effects either due to lack of leucovorin calcium efficacy or due to leucovorin calcium formulation necessitating a change in therapy. Advanced Metastatic Colorectal Cancer. The member has advanced metastatic colorectal cancer. The member is receiving palliative treatment with combination chemotherapy with 5-fluorouracil AND The member has been treated with leucovorin calcium and has experienced documented side effects either due to lack of leucovorin calcium efficacy or has experienced documented side effects due to leucovorin calcium formulation necessitating a change in therapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

LIBTAYO - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Cutaneous squamous cell carcinoma, Basal cell carcinoma (BCC), and Non-small cell lung cancer (NSCLC): Member has experienced disease progression on or after prior PD-1/PD-L1 inhibitor (e.g., Keytruda).
Required Medical Information	Cutaneous squamous cell carcinoma. The member has a diagnosis of locally advanced or metastatic cutaneous squamous cell carcinoma (CSCC) AND the disease is not amenable to curative surgery or radiation AND Libtayo (cemiplimab-rwlc) is being used as a monotherapy. Cutaneous squamous cell carcinoma (adjuvant): The member has a diagnosis of cutaneous squamous cell carcinoma at high risk of recurrence AND Libtayo (cemiplimab-rwlc) will be used as monotherapy after surgery and radiation. Basal cell carcinoma (BCC): The member has locally advanced BCC or metastatic BCC AND the disease has been treated with prior hedgehog pathway inhibitor OR treatment with a hedgehog pathway inhibitor is inappropriate AND Libtayo (cemiplimab-rwlc) is given monotherapy. Non-small cell lung cancer (NSCLC): The member has a diagnosis of NSCLC without tumor aberrations (e.g., EGFR, ALK, ROS-1) AND The disease is locally advanced (not amenable to surgery or definitive chemoradiation) or metastatic AND The tumor expresses documented high PD-L1 expression [Tumor Proportion Score (TPS) greater than or equal to 50%] AND Libtayo (cemiplimab-rwlc) is given as monotherapy OR Libtayo (cemiplimab-rwlc) is given in combination with platinum based chemotherapy as first line therapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Six month duration
Other Criteria	
Part B Prerequisite	0

LIFYORLI - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Platinum-Resistant Epithelial Ovarian, Fallopian Tube or Primary Peritoneal Cancer: Member has a diagnosis of epithelial ovarian, fallopian tube or primary peritoneal cancer AND Member has platinum-resistant disease AND Member will be using in combination with nab-paclitaxel AND Member has received at least one but no more than three prior systemic treatment regimens AND At least one of the prior systemic treatment regimens included bevacizumab product.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

LIVTENCITY - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	The member must have a diagnosis of cytomegalovirus (CMV) post-transplant AND The member must be refractory to treatment with ganciclovir, valganciclovir, cidofovir or foscarnet.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

LONSURF - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression while on Lonsurf.
Required Medical Information	Metastatic Colorectal Cancer: The member has a diagnosis of metastatic colorectal cancer AND The member is using Lonsurf as monotherapy or in combination with bevacizumab product AND The member has experienced disease progression, intolerance, or contraindication with the following therapies: fluoropyrimidine-based chemotherapy (e.g., 5-fluorouracil, capecitabine), oxaliplatin-based chemotherapy, irinotecan-based chemotherapy, and if appropriate, an anti-VEGF therapy (e.g., bevacizumab product) AND If the member is RAS wild-type: the member has experienced disease progression, intolerance, or contraindication with anti-EGFR therapy (e.g. cetuximab or panitumumab). Gastric cancer. The member has recurrent locally advanced or metastatic gastric or gastroesophageal junction adenocarcinoma AND The member has experienced disease progression on or after two lines of therapy including fluoropyrimidine (e.g., 5-fluorouracil), platinum (e.g., cisplatin), either taxane (e.g., paclitaxel) or irinotecan and if appropriate, HER2/neu-targeted therapy (e.g., trastuzumab) AND Lonsurf will be given subsequent therapy as a single agent.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Six months duration
Other Criteria	
Part B Prerequisite	0

LOQTORZI - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member has experienced disease progression on or after prior PD-1/PD-L1 inhibitor (e.g., Keytruda).
Required Medical Information	Nasopharyngeal carcinoma (NPC): The member has a diagnosis of metastatic or recurrent, locally advanced NPC AND One of the following applies: Loqtorzi (toripalimab-tpzi) will be used in first-line setting in combination with cisplatin and gemcitabine, followed by monotherapy OR Loqtorzi (toripalimab-tpzi) will be used as a single agent for disease progression on or after a platinum-containing regimen.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 months duration
Other Criteria	
Part B Prerequisite	0

LORBRENA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Non-small cell lung cancer [ALK-Positive]. The member has a diagnosis of recurrent, advanced, or metastatic non-small cell lung cancer (NSCLC) AND The disease is documented as anaplastic lymphoma kinase (ALK) positive AND Lorbrena (lorlatinib) will be given as a single agent AND Lorbrena (lorlatinib) will be given in the metastatic setting: as first line therapy OR as subsequent therapy after disease progression with a previous ALK inhibitor (e.g., alectinib, brigatinib). Non- small cell lung cancer [ROS1-Positive]: The member has a diagnosis of recurrent, advanced, or metastatic NSCLC AND The disease is documented as ROS1-positive AND Lorbrena (lorlatinib) will be given as subsequent therapy after disease progression with a previous ROS1 inhibitor [e.g., Augtyro (repotrectinib), Ibtrozi (taletrectinib), Rozlytrek (entrectinib), or Xalkori (crizotinib)] AND Lorbrena (lorlatinib) will be given as a single agent.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

LUMAKRAS - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member has experienced disease progression on a KRAS G12C inhibitor (e.g., sotorasib, adagrasib). [All indications]
Required Medical Information	Non-small cell lung cancer (NSCLC): The member has a diagnosis of locally advanced or metastatic NSCLC AND The NSCLC has documented KRAS G12C mutation AND The member has experienced disease progression on one prior therapy AND Lumakras (sotorasib) will be given as monotherapy as subsequent treatment. Colorectal Cancer [KRAS G12C-mutated]:The member has a diagnosis of metastatic colorectal cancer (mCRC) AND The disease is documented as KRAS G12C-mutated, as determined by an FDA-approved test AND The member has received prior systemic treatment with chemotherapy [fluoropyrimidine-based chemotherapy (e.g., 5-fluorouracil, capecitabine), oxaliplatin, irinotecan] AND Lumakras (sotorasib) will be given in combination with Vectibix (panitumumab).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 Months Duration
Other Criteria	
Part B Prerequisite	0

LUMIGAN - DROPS

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Member must have a previous treatment, intolerance or contraindication to an AB-rated generic equivalent product.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

LUPRON DEPOT - SYRINGE KIT (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Prostate Cancer: Diagnosis of advanced prostate cancer or at risk for disease recurrence. Endometriosis: Diagnosis of endometriosis. Fibroids (Uterine leiomyomata). The patient must have a diagnosis of anemia due to uterine leiomymota. Central Precocious Puberty: Pediatric member with a diagnosis of central precocious puberty (idiopathic or neurogenic). Invasive Breast Cancer: The patient has a diagnosis of hormone responsive (ER and/or PR +) invasive breast cancer AND The patient must be pre or perimenopausal. Recurrent Ovarian Cancer: Patient has recurrent ovarian cancer (epithelial cell cancer, fallopian tube cancer, primary peritoneal cancer or ovarian stromal tumor).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration for all except for Endometriosis: 6 months and Fibroids: 3 months
Other Criteria	
Part B Prerequisite	0

LUPRON DEPOT (3 MONTH) - SYRINGE KIT (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Prostate Cancer: Diagnosis of advanced prostate cancer or at risk for disease recurrence. Endometriosis: Diagnosis of endometriosis. Fibroids (Uterine leiomyomata). The patient must have a diagnosis of anemia due to uterine leiomymota. Central Precocious Puberty: Pediatric member with a diagnosis of central precocious puberty (idiopathic or neurogenic). Invasive Breast Cancer: The patient has a diagnosis of hormone responsive (ER and/or PR +) invasive breast cancer AND The patient must be pre or perimenopausal. Recurrent Ovarian Cancer: Patient has recurrent ovarian cancer (epithelial cell cancer, fallopian tube cancer, primary peritoneal cancer or ovarian stromal tumor).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration for all except for Endometriosis: 6 months and Fibroids: 3 months
Other Criteria	
Part B Prerequisite	0

LUTRATE DEPOT (3 MONTH) - VIAL (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Prostate Cancer: Diagnosis of advanced prostate cancer or at risk for disease recurrence. Endometriosis: Diagnosis of endometriosis. Fibroids (Uterine leiomyomata). The patient must have a diagnosis of anemia due to uterine leiomymota. Central Precocious Puberty: Pediatric member with a diagnosis of central precocious puberty (idiopathic or neurogenic). Invasive Breast Cancer: The patient has a diagnosis of hormone responsive (ER and/or PR +) invasive breast cancer AND The patient must be pre or perimenopausal. Recurrent Ovarian Cancer: Patient has recurrent ovarian cancer (epithelial cell cancer, fallopian tube cancer, primary peritoneal cancer or ovarian stromal tumor).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration for all except for Endometriosis: 6 months and Fibroids: 3 months
Other Criteria	
Part B Prerequisite	0

LYBALVI - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Use of opioids. Episode of acute opioid withdrawal.
Required Medical Information	Schizophrenia: The member must have a diagnosis of schizophrenia AND The member has tried or cannot use at least one of the following generic atypical antipsychotics: olanzapine, risperidone, quetiapine, ziprasidone, aripiprazole, or lurasidone. Bipolar I Disorder (acute manic or mixed, maintenance monotherapy): The member must have a diagnosis of bipolar I disorder (acute manic or mixed, maintenance monotherapy) AND The member has tried or cannot use at least one of the following generic atypical antipsychotics: olanzapine, risperidone, quetiapine, ziprasidone, or aripiprazole.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

LYNPARZA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on or following PARP inhibitor therapy [e.g. Rubraca (rucaparib), Lynparza (olaparib), Zejula (niraparib)]. Adjuvant setting for High-Risk Early Breast Cancer: member is taking Lynparza (olaparib) total treatment for more than one year.
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	Pending CMS Review
Part B Prerequisite	0

LYTGOBI - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Disease progression on Lytgobi (futibatinib)
Required Medical Information	Cholangiocarcinoma: The member has unresectable locally advanced or metastatic intrahepatic cholangiocarcinoma (iCCA) AND The member has iCCA with documented FGFR2 gene fusions or other rearrangements AND The member has received prior treatment AND Lytgobi (futibatinib) is given as a single agent for subsequent therapy
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Six month duration
Other Criteria	
Part B Prerequisite	0

macitentan - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pulmonary Arterial Hypertension (PAH). The member has a diagnosis of pulmonary arterial hypertension (WHO Group I) confirmed by right heart catheterization.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

MARGENZA - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member experiences disease progression on Margenza (margetuximab-cmkb).
Required Medical Information	Breast cancer: The member has a diagnosis of metastatic breast cancer AND The disease is documented HER2 neu positive AND The member has received two prior anti-HER2 neu based therapies (including trastuzumab products) where one therapy was given in the metastatic setting AND Margenza (margetuximab-cmkb) is given in combination with chemotherapy (gemcitabine, eribulin, vinorelbine, capecitabine) as subsequent therapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

MEKINIST - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Melanoma - Unresectable or metastatic: Members on concomitant Yervoy (ipilimumab), Zelboraf (vemurafenib), Opdivo (nivolumab), Cotellic (cobimetinib), Braftovi (encorafenib), or Mektovi (binimetinib). Members that have experienced disease progression while on Mekinist (trametinib). Members that have experienced disease progression while on prior anti-BRAF/MEK combination therapy [e.g. Cotellic (cobimetinib) with Zelboraf (vemurafenib) or Braftovi (encorafenib) with Mektovi (binimetinib)]. Non-small cell lung cancer, Anaplastic Thyroid Cancer and Metastatic Solid Tumors: Members on concomitant Yervoy (ipilimumab), Zelboraf (vemurafenib), Opdivo (nivolumab), Keytruda (pembrolizumab), Cotellic (cobimetinib), Braftovi (encorafenib), or Mektovi (binimetinib). Members that have experienced disease progression while on prior anti-BRAF/MEK combination therapy [e.g. Cotellic (cobimetinib) with Zelboraf (vemurafenib) or Braftovi (encorafenib) with Mektovi (binimetinib)]. Adjuvant melanoma: member is taking Mekinist (trametinib) total treatment for more than one year. Members on concomitant Yervoy (ipilimumab), Zelboraf (vemurafenib), Opdivo (nivolumab), Keytruda (pembrolizumab), Cotellic (cobimetinib), Braftovi (encorafenib), or Mektovi (binimetinib). Members that have experienced disease progression while on prior anti-BRAF/MEK combination therapy [e.g. Cotellic (cobimetinib) with Zelboraf (vemurafenib) or Braftovi (encorafenib) with Mektovi (binimetinib)]. Low-grade Glioma: Members that have experienced disease progression while on Mekinist (trametinib). Members that have experienced disease progression while on prior anti-BRAF/MEK combination therapy [e.g. Cotellic (cobimetinib) with Zelboraf (vemurafenib) or Braftovi (encorafenib) with Mektovi (binimetinib)].

MEKINIST - TABLET

Required Medical Information	Melanoma - Unresectable or Metastatic: The member has a diagnosis of unresectable or stage IV metastatic melanoma AND The member has a documented BRAF V600 activating mutation AND The member will be using Mekinist as a single-agent (member has not received prior BRAF-inhibitor therapy) OR in combination with Tafinlar (dabrafenib). Non-small cell lung cancer: The member has a diagnosis of recurrent or metastatic non-small cell lung cancer(NSCLC) AND The member has a documented BRAF V600E mutation AND The member will be using Mekinist (trametinib) in combination with Tafinlar (dabrafenib). Melanoma - Adjuvant. The member has a diagnosis of stage III melanoma AND The member has undergone lymph node resection of involved lymph nodes AND The member has a documented BRAF V600 activating mutation AND The member will be using Mekinist (trametinib) in combination with Tafinlar (dabrafenib) for adjuvant treatment. Anaplastic Thyroid Cancer. The member has a diagnosis of locally advanced or metastatic anaplastic thyroid cancer AND The member has a documented V600E mutation AND The member has no satisfactory locoregional treatment options AND The member will be using Tafinlar (dabrafenib) in combination with Mekinist (trametinib). Metastatic Solid Tumors: the member has a diagnosis of unresectable or metastatic solid tumors with documented BRAF V600E mutation AND the member has disease that has progressed on prior therapy and has no satisfactory alternative treatments AND Mekinist is given in combination with Tafinlar. Low-grade Glioma: The member has a diagnosis of low-grade glioma (LGG) AND The member has a documented BRAF V600E mutation AND Mekinist (trametinib) will be used in combination with Tafinlar (dabrafenib).
Age Restriction	Low-grade Glioma only: The member is a pediatric age 1 year of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

MEKTOVI - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members on concomitant Yervoy (ipilimumab), Opdivo (nivolumab), Keytruda (pembrolizumab), Zelboraf (vemurafenib), Cotellic (cobimetinib), Tafinlar (dabrafenib), or Mekinist (trametinib). Members that have experienced disease progression while on prior anti-BRAF/MEK combination therapy [e.g. Cotellic (cobimetinib) with Zelboraf (vemurafenib) or Tafinlar (dabrafenib) with Mekinist (trametinib)].
Required Medical Information	Melanoma - Unresectable or metastatic. The member has a diagnosis of unresectable or stage IV metastatic melanoma AND The member has a documented BRAF V600 activating mutation AND The member will be using Braftovi (encorafenib) in combination with Mektovi (binimetinib). Metastatic non-small cell lung cancer: The member has documented BRAF V600E metastatic non-small cell lung cancer (NSCLC) AND The member will be using Braftovi (encorafenib) in combination with Mektovi (binimetinib).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

memantine - SOLUTION, ORAL

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Diagnosis of Autism or Atypical Autism (PDD)
Required Medical Information	
Age Restriction	An automatic approval if member is greater than 26 years of age.Prior Auth required for age 26 or younger.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

mifepristone - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Hyperglycemia secondary to hypercortisolism. Diagnosis of endogenous Cushing's syndrome. AND Type 2 diabetes mellitus or glucose intolerance. AND Failed surgery or are not candidates for surgery.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

MODEYSO - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Diffuse Midline Glioma: The member has a diagnosis of diffuse midline glioma AND The disease has documented H3 K27M mutation AND The member has experienced disease progression or recurrence following prior therapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

MOUNJARO - PEN INJECTOR (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member is using for weight loss.
Required Medical Information	Diabetes Mellitus. The member has a diagnosis of type 2 diabetes mellitus.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

MVASI - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Malignant Pleural Mesothelioma, Soft Tissue Sarcoma, Recurrent Primary CNS Tumor (including Glioblastoma multiforme), Renal Cell Carcinoma, Non-small cell lung cancer (non-squamous cell histology): The member has experienced disease progression while on or following a bevacizumab product. Metastatic colorectal cancer: A bevacizumab product may not be used in conjunction with Vectibix (panitumumab) or Erbitux (cetuximab).
Required Medical Information	For Avastin (bevacizumab), Alymsys (bevacizumab-maly), Jobevne (bevacizumab-nwgd), and Vegzelma (bevacizumab-adcd) oncology requests: member must have an intolerance or contraindication with Mvasi or Zirabev. Metastatic colorectal cancer: metastatic colorectal cancer AND 1 of the following applies: using a bevacizumab product in combo with fluoropyrimidine (e.g., 5-fluorouracil or capecitabine) based chemo for 1st or 2nd-line therapy OR in combo with fluoropyrimidine-irinotecan or fluoropyrimidine-oxaliplatin-based chemo for 2nd-line therapy in patients who have progressed on 1st-line bevacizumab-containing regimens. Non-small cell lung cancer (non-squamous cell histology). NSCLC with non-squamous cell histology AND the disease is unresectable, locally advanced, recurrent, or metastatic AND A bevacizumab product will be used as part of a supported combination chemotherapy. Hepatocellular carcinoma: unresectable or metastatic HCC AND a bevacizumab product will be used as 1st line therapy in combo with an atezolizumab product.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration

MVASI - VIAL (ML)

Other Criteria	Renal cell carcinoma: Member has RCC and a bevacizumab product will be used in combo with interferon alpha. Recurrent Primary CNS Tumor (including Glioblastoma multiforme): Diagnosis of progressive or recurrent glioblastoma or anaplastic glioma AND a bevacizumab product is being used as a single agent or in combo with irinotecan, carmustine, lomustine or temozolomide AND the member does not have a CNS hemorrhage. Soft Tissue Sarcoma: One of the following apples: Diagnosis of angiosarcoma AND a bevacizumab product is being used as a single agent OR The member has a diagnosis of solitary fibrous tumor and a bevacizumab product is being used in combo with temozolomide. Macular Retinal Edema. Avastin is being used to treat central or branch retinal vein occlusion with macular retinal edema. Cervical Cancer: member has recurrent, or metastatic cervical cancer AND A bevacizumab product will be used as part of a supported combination therapy. Endometrial Cancer: The member has a diagnosis of endometrial cancer AND One of the following applies: The member has stage III-IV disease or recurrent disease AND a bevacizumab product will be used in combination with carboplatin and paclitaxel, followed by maintenance with a bevacizumab product maintenance OR The member has recurrent disease AND a bevacizumab product monotherapy will be used as second-line or subsequent therapy if not previously used. Malignant Pleural Mesothelioma. Diagnosis of unresectable malignant pleural mesothelioma and a bevacizumab product will be used in combo with cisplatin and pemetrexed or carboplatin and pemetrexed followed by maintenance with a bevacizumab product (for responders). Epithelial ovarian, fallopian tube, or primary peritoneal cancer: Diagnosis of epithelial ovarian, fallopian tube or primary peritoneal cancer AND One of the following applies: The disease is recurrent, platinum-resistant AND a bevacizumab product will be used in combination with paclitaxel, pegylated liposomal doxorubicin or topotecan AND the member has not received more than 2 prior chemotherapy regimens OR The disease is recurrent, platinum-sensitive AND a bevacizumab product will be used in combination with carboplatin and paclitaxel, or in combination with carboplatin and gemcitabine, followed by a bevacizumab product monotherapy AND The disease is stage III or IV AND initial surgical resection was completed AND a bevacizumab product will be used in combo with carboplatin and paclitaxel, followed by a bevacizumab product monotherapy OR . Age Related Macular Degeneration (Avastin requests only): Avastin (bevacizumab) is being used to treat age related macular degeneration. Diabetic Macular Edema (Avastin requests only): Avastin (bevacizumab) is being used to treat diabetic macular edema.
Part B Prerequisite	0

NERLYNX - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member has disease progression on Nerlynx (neratinib). Member is taking Nerlynx (neratinib) total treatment for more than one year [applicable only to early stage breast cancer].
Required Medical Information	Early stage Breast Cancer: The member has early stage (i.e. Stage I, II, III) documented HER2 + positive disease AND The member has completed adjuvant therapy with a trastuzumab containing treatment AND Nerlynx (neratinib) is being used for the treatment in extended adjuvant setting. Metastatic Breast Cancer. The member has metastatic or advanced breast cancer and all of the following apply: The member has documented HER2 positive disease and The member has received two or more prior anti-HER2 based regimens in the metastatic setting and Nerlynx (neratinib) is given in combination with capecitabine.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

NEXLETOL - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Hypercholesterolemia: Member has a diagnosis of hypercholesterolemia (including heterozygous familial hypercholesterolemia [HeFH]) and Has tried or cannot use a statin. Cardiovascular Disease Prevention: Member must have an increased risk of major adverse cardiovascular events (e.g acute coronary syndromes, history of myocardial infarction, stable or unstable angina, coronary or other arterial revascularization, stroke, transient ischemic attack or peripheral arterial disease, all of presumed atherosclerotic origin) AND Has tried or cannot use a statin.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

NEXLIZET - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Hypercholesterolemia: Member has a diagnosis of hypercholesterolemia (including heterozygous familial hypercholesterolemia [HeFH]) and Has tried or cannot use a statin. Cardiovascular Disease Prevention: Member must have an increased risk of major adverse cardiovascular events (e.g acute coronary syndromes, history of myocardial infarction, stable or unstable angina, coronary or other arterial revascularization, stroke, transient ischemic attack or peripheral arterial disease, all of presumed atherosclerotic origin) AND Has tried or cannot use a statin.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

nilotinib hcl - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on any nilotinib product. For ALL and CML: The member has one of the following mutations: T315I, Y253H, E255K/V, or F359V/C/I.
Required Medical Information	Chronic Myelogenous Leukemia (CML). The member has a diagnosis of Philadelphia chromosome positive chronic myeloid leukemia (CML) AND One of the following applies: The member has accelerated or blast phase CML OR The member has a diagnosis of chronic phase CML that has not been previously treated, and one of the following applies: Intermediate- or high-risk score for disease progression OR Low-risk score for disease progression and has contraindication to, intolerance to, or unable to achieve treatment goals with imatinib OR The members has a diagnosis of chronic phase CML that has received previous treatment. Advanced Gastrointestinal Stromal Tumor (GIST). Diagnosis of advanced unresectable GIST.The member has progressive disease or is intolerant to prior therapy with imatinib, sunitinib, or Stivarga. Acute Lymphoblastic Leukemia (ALL). The member has diagnosis of Philadelphia positive acute lymphoblastic leukemia. Pediatric CML: Diagnosis of chronic phase Ph+ chronic myeloid leukemia (CML) OR Diagnosis of accelerated phase Ph+ chronic myeloid leukemia (CML) AND resistance, intolerance, or contraindication to prior TKI therapy.
Age Restriction	Pediatric CML- member is greater than or equal to 1 year of age.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

NINLARO - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members with disease progression on Ninlaro (ixazomib).
Required Medical Information	Multiple Myeloma: second line. The member has a diagnosis of relapsed or refractory multiple myeloma AND Ninlaro (ixazomib) will be used after disease progression on at least one prior therapy AND Ninlaro (ixazomib) will be used in combination with lenalidomide and dexamethasone or cyclophosphamide and dexamethasone (Omission of corticosteroid from regimen is allowed if intolerance/contraindication). Multiple Myeloma: third line or subsequent. The member has a diagnosis of relapsed or refractory multiple myeloma AND Ninlaro (ixazomib) will be used after disease progression on at least two prior therapies AND Ninlaro (ixazomib) will be used in combination with pomalidomide and dexamethasone (Omission of corticosteroid from regimen is allowed if intolerance/contraindication) AND The members has demonstrated disease progression on or within 60 days of completion of the last therapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 Months Duration
Other Criteria	
Part B Prerequisite	0

nintedanib - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

NIVESTYM - SYRINGE (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 Months Duration
Other Criteria	Febrile Neutropenia Prophylaxis, in members with acute myeloid leukemia receiving chemotherapy: The member must have a diagnosis Acute Myeloid Leukemia (AML). The member must be receiving either induction chemotherapy OR consolidation Chemotherapy AND The member is not administering filgrastim product earlier than 24 hours after cytotoxic chemotherapy or within 24 hours before chemotherapy. Febrile Neutropenia Prophylaxis, In non-myeloid malignancies following Hematopoietic Stem Cell Transplant (HCST): The member must have had a Hematopoietic Stem Cell Transplant (HCST) (e.g. bone marrow transplant, peripheral-blood progenitor cell (PBPC) transplant) for a non-myeloid malignancy AND The member is not administering filgrastim product earlier than 24 hours after cytotoxic chemotherapy or within 24 hours before chemotherapy Harvesting of peripheral blood stem cells: The member is using as a component of a hematopoietic stem cell mobilization protocol and one of the following applies: is scheduled for autologous peripheral-blood stem cell (PBSC) transplantation, is storing cells for a possible future autologous transplant, or is donating stem cells for an allogeneic or syngeneic PBSC transplant. Neutropenic disorder, chronic (Severe), Symptomatic: The member must have a diagnosis of congenital, cyclic, or idiopathic neutropenia. Neutropenia in AIDS patients: The member must have a diagnosis of AIDS with neutropenia. Treatment of Aplastic Anemia: The member must have a diagnosis of Aplastic Anemia. Treatment of Agranulocytosis: The member must have a diagnosis of congenital or drug induced agranulocytosis. Hematopoietic Syndrome of Acute radiation syndrome: The member has been acutely exposed to myelosuppressive doses of radiation.
Part B Prerequisite	0

NUBEQA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on Nubeqa (darolutamide).
Required Medical Information	Prostate Cancer (non-metastatic castration-resistant): The member has a diagnosis of non-metastatic castration-resistant prostate cancer AND the member will use Nubeqa (darolutamide) in combination with androgen deprivation therapy (e.g. previous bilateral orchiectomy or luteinizing hormone-releasing hormone (LHRH) analog [i.e., LHRH agonist/ antagonist]). Prostate Cancer (metastatic hormone-sensitive prostate cancer): The member has a diagnosis of metastatic hormone-sensitive prostate cancer AND the member will use Nubeqa (darolutamide) alone or with docetaxel AND the member will use Nubeqa (darolutamide) in combination with androgen deprivation therapy (e.g. previous bilateral orchiectomy or LHRH analog [i.e., LHRH agonist/ antagonist]).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

NUCALA - SYRINGE (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	Chronic Obstructive Pulmonary Disease (COPD) with an eosinophilic phenotype: Member must be 18 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	Chronic Rhinosinusitis with Nasal Polyposis: The member must meet ALL of the following criteria: Diagnosis of Chronic Rhinosinusitis with Nasal Polyposis AND Nucala (mepolizumab) will be used in conjunction with a daily intranasal corticosteroid spray AND member is unable to achieve adequate control of symptoms with maximum tolerated intranasal corticosteroid therapy. Chronic Obstructive Pulmonary Disease (COPD) with an eosinophilic phenotype: Diagnosis of Chronic Obstructive Pulmonary Disease (COPD) AND Member has an eosinophilic phenotype defined as an elevated peripheral blood eosinophil level of the following before therapy is started: greater than or equal to 150 cells/microliter OR greater than or equal to 300 cells/microliter within 12 months prior to therapy initiation. Member has been unable to achieve adequate control of COPD while on an: Inhaled corticosteroid therapy in combination with a long-acting beta agonist (LABA) and long-acting muscarinic antagonist (LAMA) OR LABA-LAMA therapy if inhaled corticosteroids are contraindicated.
Part B Prerequisite	0

NUPLAZID - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Parkinson's Disease Psychosis: The member is using Nuplazid for the treatment of hallucinations and delusions associated with Parkinson's disease (PD) psychosis AND the symptoms of psychosis have appeared after the diagnosis of PD AND psychosis is not related to other causes other than PD.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

octreotide acetate - SYRINGE (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

octreotide,microspheres - VIAL (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ODOMZO - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression while on Odomzo.
Required Medical Information	Basal Cell Carcinoma: The member has a diagnosis of locally advanced basal cell carcinoma AND The member has experienced recurrence or disease progression following surgery or radiation OR has a contraindication to surgery or radiation.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

OFEV - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

OGIVRI - VIAL (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	For Herceptin (trastuzumab), Kanjinti (trastuzumab-anns) , Herzuma (trastuzumab-pkrb), Hercessi (trastuzumab-strf) or Ontruzant (trastuzumab-dkst) requests: member must have an intolerance or contraindication Trazimera (trastuzumab-qyyp) OR Ogivri (trastuzumab-dkst) and meets below criteria: Breast Cancer: The member has a diagnosis of breast cancer and HER2 (human epidermal growth factor receptor2) positive disease (e.g., defined as IHC 3+ or IHC2+/ISH positive). Gastric Cancer: The member has a diagnosis of advanced, gastric cancer or gastroesophageal adenocarcinoma and HER2 positive disease (e.g., defined as IHC 3+ or IHC2+/ISH positive) AND trastuzumab is being used in combination with cisplatin and fluorouracil or capecitabine.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	All Medically accepted uses not listed above: The member is using a trastuzumab product for a medically accepted indication as defined by Medicare rules. For Herceptin (trastuzumab), Kanjinti (trastuzumab-anns), Herzuma (trastuzumab-pkrb), Hercessi (trastuzumab-strf), or Ontruzant (trastuzumab-dttb) requests: member must have an intolerance or contraindication to Trazimera (trastuzumab-qyyp) or Ogivri (trastuzumab-dkst).
Part B Prerequisite	0

OGSIVEO - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member has had disease progression on Ogsiveo (nirogacestat).
Required Medical Information	Desmoid tumors: Member has been diagnosed with progressing desmoid tumor AND Member will be using Ogsiveo (nirogacestat) as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

OJEMDA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member has experienced disease progression while on Ojemda (tovorafenib).
Required Medical Information	Pediatric Low-Grade Glioma (pLGG): The member has a diagnosis of relapsed or refractory pediatric low-grade glioma (LGG) AND The member has a documented BRAF fusion or rearrangement, or documented BRAF V600E mutation.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

OJJAARA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on Ojjaara (momelotinib).
Required Medical Information	Myelofibrosis: The member has a diagnosis of primary myelofibrosis or secondary myelofibrosis (i.e. post-polycythemia vera or post-essential thrombocythemia) AND The member has one of the following risk categories, as defined by accepted risk stratification tools for myelofibrosis (e.g. International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or DIPSS-PLUS): Intermediate-risk disease OR High-risk disease AND The member has anemia* AND The member will be using Ojjaara (momelotinib) as monotherapy AND The member has tried, intolerant to, or has a contraindication to Jakafi (Ruxolitinib) or has hemoglobin less than 8 g/dL. *Anemia is defined as hemoglobin less than 10 g/dL or having transfusion-dependent anemia.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

OMLYCLO - SYRINGE (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Chronic Idiopathic Urticaria. Member has a diagnosis of chronic idiopathic urticaria. Member has remained symptomatic despite at least 2 weeks of H1 antihistamine therapy, unless contraindicated. Member will continue to receive H1 antihistamine therapy while on Xolair, unless contraindicated. Moderate or Severe persistent asthma. The patient has a diagnosis of moderate or severe persistent asthma. The patient has evidence of specific allergic sensitivity confirmed by positive skin test (i.e. prick/puncture test) or blood test (i.e. RAST) for a specific IgE or in vitro reactivity to a perennial aeroallergen. Member must have baseline serum IgE greater than or equal to 30 IU/ml. The patient has inadequately controlled asthma despite the use of Inhaled Corticosteroids.
Age Restriction	The patient is 12 years of age or older for diagnosis of chronic idiopathic urticaria. The patient is 6 years of age or older for moderate to severe persistent asthma. The patient is 18 years of age or older for nasal polyps. The patient is 1 year or older for IgE Mediated Food Allergy.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	Nasal Polyps: The member must meet all of the following criteria: have a diagnosis of nasal polyps (e.g., Chronic Rhinosinusitis with Nasal Polyposis [CRSwNP]) AND Xolair will be used in combination with a daily intranasal corticosteroid spray AND is unable to achieve adequate control of symptoms with maximum tolerated intranasal corticosteroid therapy. IgE Mediated Food Allergy: Member has been diagnosed with IgE mediated food allergy AND Member is using Xolair to reduce allergic reactions (Type I), including anaphylaxis, that may occur with accidental exposure to one of more foods.
Part B Prerequisite	0

OMNITROPE - VIAL (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	Pending CMS Review
Part B Prerequisite	0

ONUREG - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression on hypomethylators (e.g. azacitidine, decitabine).
Required Medical Information	Acute Myeloid Leukemia: The member has a diagnosis of acute myeloid leukemia AND The member is using Onureg (azacitidine) for post-remission therapy AND The member has achieved first complete remission (CR) or complete remission with incomplete blood count recovery (CRi) following intensive induction chemotherapy AND The member is not able to complete or declines intensive curative therapy (e.g. allogeneic hematopoietic stem cell transplant) AND The member will use Onureg (azacitidine) as a single agent.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

OPDIVO - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	All indications except Hepatocellular carcinoma, MSI-H or dMMR: Disease progression while on or following anti-PD-1/PD-L1 therapy (e.g. Opdivo [nivolumab], Keytruda [pembrolizumab], Tecentriq [atezolizumab], Bavencio [avelumab]). Adjuvant melanoma only: member is taking Opdivo (nivolumab) total treatment for more than one year. Neoadjuvant NSCLC only: member is taking Opdivo (nivolumab) total treatment in combination with platinum-doublet chemotherapy for more than 3 cycles. Hepatocellular carcinoma: Disease progression while on or following anti-PD-1/PD-L1 combination therapy (e.g. Imjudo [tremelimumab-actl] with Imfinzi [durvalumab]). MSI-H or dMMR: Disease progression while on or following anti-PD-1/PD-L1 combination therapy (e.g. Opdivo [nivolumab] with Yervoy [ipilimumab].
Required Medical Information	Melanoma: diagnosis of unresectable or metastatic melanoma AND using as monotherapy OR in combo with Yervoy (ipilimumab) followed by Opdivo (nivolumab) or Opdivo Qvantig (nivolumab and hyaluronidase-nvhy) monotherapy. Melanoma-Adjuvant: diagnosis of stage IIB, IIC, III or stage IV melanoma AND undergone complete resection of disease AND using as adjuvant treatment AND using as monotherapy. Non-Small Cell Lung Cancer-subsequent therapy: diagnosis of metastatic squamous or non-squamous NSCLC AND experienced disease progression on or after chemo and EGFR inhibitor, if EGFR mutation positive or ALK inhibitor, if ALK positive AND using as monotherapy. Renal Cell Carcinoma (RCC): diagnosis of advanced RCC AND one of the following applies: will be using as subsequent therapy AND has received prior anti-angiogenic therapy OR Will be using in combo with Yervoy followed by Opdivo or Opdivo Qvantig monotherapy as first line therapy OR Will be using in combo with Cabometyx (cabozantinib) as first line therapy. Classical Hodgkin Lymphoma: diagnosis of classical Hodgkin Lymphoma AND one of the following applies: has relapsed or refractory disease AND using as monotherapy or in combo with a supported combo therapy OR has newly diagnosed disease AND using in combo with doxorubicin, vinblastine and dacarbazine. Non-Small Cell Lung Cancer (NSCLC)-Perioperative: diagnosis of NSCLC AND must not have any known epidermal growth factor receptor (EGFR) mutations or anaplastic lymphoma kinase (ALK) rearrangements AND must have resectable disease (tumors greater than or equal to 4 cm or node positive) AND using in the neoadjuvant setting, in combo with platinum-doublet chemo followed by Opdivo or Opdivo Qvantig as monotherapy. Hepatocellular Carcinoma: The member has a diagnosis of hepatcellular carcinoma AND using in combo with Yervoy followed by Opdivo or Opdivo Qvantig monotherapy.
Age Restriction	

OPDIVO - VIAL (ML)

Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 months duration

OPDIVO - VIAL (ML)

Other Criteria	Non-nasopharyngeal recurrent or metastatic Squamous Cell Carcinoma of the Head and Neck (SCCHN) AND using as monotherapy AND disease progression on or after platinum based therapy. Locally advanced or metastatic urothelial cancer AND using as monotherapy AND 1 of the following apply: used as 2nd or subsequent line-therapy OR Disease progression within 12 months of neoadjuvant or adjuvant chemo OR high risk of recurrence after radical surgical resection of disease. Unresectable or metastatic urothelial carcinoma AND in combo with cisplatin and gemcitabine, followed by monotherapy maintenance. Unresectable or metastatic colorectal cancer with documented Microsatellite Instability-High (MSI-H) or Mistmatch Repair Deficient (dMMR) AND 1 of the following applies: disease progressed following treatment with oxaliplatin-, irinotecan-, or fluoropyrimidine-based therapy and will be using Opdivo or Opdivo Qvantiq as monotherapy OR using in combo with Yervoy, followed by Opdivo or Opdivo Qvantig as monotherapy. Non-small cell lung cancer (NSCLC) 1st Line Therapy: metastatic NSCLC AND 1 of the following applies: Disease with PD-L1 expression greater than or equal to 1% with no EGFR or ALK genomic tumor aberrations and given as 1st line therapy AND Tumor expresses PD-L1 as determined by FDA-approved test AND in combo with Yervoy OR Disease with no EGFR or ALK genomic tumor aberrations and given as 1st line therapy AND in combo with Yervoy AND used in combo with 2 cycles of platinum doublet chemo. Esophageal cancer: unresectable advanced, recurrent, or metastatic squamous cell carcinoma of the esophagus and 1 of the following apply: disease progressed after prior fluoropyrimidine- and platinum-based chemo AND will be given as subsequent monotherapy OR in combo with fluoropyrimidine- and platinum containing chemo for 1st line treatment AND the tumor expressed PD-L1 great than or equal to 1 OR as 1st line treatment Opdivo in combo with Yervoy AND the tumor expressed PD-L1 great than or equal. Unresectable malignant pleural mesothelioma AND 1 of the following scenarios applies: 1st-line treatment AND in combo with Yervoy OR as subsequent treatment, if not administered 1st-line AND as single agent OR in combo with Yervoy. Gastric Cancer, Gastroesophageal Junction Cancer, and Esophageal Adenocarcinoma (advanced or metastatic): given with a regimen containing fluoropyrimidine- and platinum-based chemo AND the tumor expressed PD-L1 great than or equal. Esophageal or Gastroesophageal Junction cancer (residual disease post-surgery and preoperative chemoradiation): received neoadjuvant chemoradiation AND complete surgical resection of esophageal or gastroesophageal junction AND has residual pathologic disease AND given as subsequent monotherapy. NSCLC- Neoadjuvant Therapy: early stage NSCLC AND has resectable disease (tumors greater than 4 cm or node positive) AND using in neoadjuvant setting, in combo with platinum-doublet chemo.
Part B Prerequisite	0

OPDIVO QVANTIG - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	All indications except Hepatocellular carcinoma, MSI-H or dMMR: Disease progression while on or following anti-PD-1/PD-L1 therapy (e.g. Opdivo [nivolumab], Keytruda [pembrolizumab], Tecentriq [atezolizumab], Bavencio [avelumab]). Adjuvant melanoma only: member is taking Opdivo Qvantig (nivolumab and hyaluronidase-nvhy) total treatment for more than one year. Neoadjuvant NSCLC only: member is taking Opdivo Qvantig (nivolumab and hyaluronidase-nvhy) total treatment in combination with platinum-doublet chemotherapy for more than 3 cycles. Hepatocellular carcinoma: Disease progression while on or following anti-PD-1/PD-L1 combination therapy (e.g. Imjudo [tremelimumab-actl] with Imfinzi [durvalumab]). MSI-H or dMMR: Disease progression while on or following anti-PD-1/PD-L1 combination therapy (e.g. Opdivo [nivolumab] with Yervoy [ipilimumab].
Required Medical Information	Melanoma: diagnosis of unresectable or metastatic melanoma AND using as monotherapy OR Opdivo (nivolumab) in combo with Yervoy (ipilimumab) followed by Opdivo Qvantig (nivolumab and hyaluronidase-nvhy) monotherapy. Melanoma-Adjuvant: diagnosis of stage IIB, IIC, III or stage IV melanoma AND undergone complete resection of disease AND using as adjuvant treatment AND using as monotherapy. Non-Small Cell Lung Cancer-subsequent therapy: diagnosis of metastatic squamous or non-squamous NSCLC AND experienced disease progression on or after chemo and EGFR inhibitor, if EGFR mutation positive or ALK inhibitor, if ALK positive AND using as monotherapy. Renal Cell Carcinoma (RCC): diagnosis of advanced RCC AND one of the following applies: will be using as subsequent therapy AND has received prior anti-angiogenic therapy OR Opdivo (nivolumab)using in combo with Yervoy followed by Opdivo Qvantig monotherapy as first line therapy OR Will be using in combo with Cabometyx (cabozantinib) as first line therapy. Non-Small Cell Lung Cancer (NSCLC)-Perioperative: diagnosis of NSCLC AND must not have any known epidermal growth factor receptor (EGFR) mutations or anaplastic lymphoma kinase (ALK) rearrangements AND must have resectable disease (tumors greater than or equal to 4 cm or node positive) AND using in the neoadjuvant setting, in combo with platinum-doublet chemo followed by Opdivo or Opdivo Qvantig as monotherapy. Hepatocellular Carcinoma: The member has a diagnosis of hepatcellular carcinoma AND Opdivo (nivolumab) using in combo with Yervoy, followed by Opdivo Qvantig monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner

OPDIVO QVANTIG - VIAL (ML)

Coverage Duration	6 months duration
Other Criteria	Non-nasopharyngeal recurrent or metastatic Squamous Cell Carcinoma of the Head and Neck (SCCHN) AND using as monotherapy AND disease progression on or after platinum based therapy. Locally advanced or metastatic urothelial cancer AND using as monotherapy AND 1 of the following apply: used as 2nd or subsequent line-therapy OR Disease progression within 12 months of neoadjuvant or adjuvant chemo OR high risk of recurrence after radical surgical resection of disease. Unresectable or metastatic urothelial carcinoma AND in combo with cisplatin and gemcitabine, followed by monotherapy maintenance. Unresectable or metastatic colorectal cancer with documented Microsatellite Instability-High (MSI-H) or Mistmatch Repair Deficient (dMMR) AND 1 of the following applies: disease progressed following treatment with oxaliplatin-, irinotecan-, or fluoropyrimidine-based therapy and will be as monotherapy OR Opdivo (nivolumab) using in combo with Yervoy, followed by Opdivo Qvantig as monotherapy. Non-small cell lung cancer (NSCLC) 1st Line Therapy: metastatic NSCLC AND 1 of the following applies: Disease with PD-L1 expression greater than or equal to 1% with no EGFR or ALK genomic tumor aberrations and given as 1st line therapy AND Tumor expresses PD-L1 as determined by FDA-approved test OR Disease with no EGFR or ALK genomic tumor aberrations and given as 1st line therapy AND used in combo with 2 cycles of platinum doublet chemo. Esophageal cancer: unresectable advanced, recurrent, or metastatic squamous cell carcinoma of the esophagus and 1 of the following apply: disease progressed after prior fluoropyrimidine- and platinum-based chemo AND will be given as subsequent monotherapy OR in combo with fluoropyrimidine- and platinum containing chemo for 1st line treatment AND the tumor expressed PD-L1 great than or equal to 1. Gastric Cancer, Gastroesophageal Junction Cancer, and Esophageal Adenocarcinoma (advanced or metastatic): given with a regimen containing fluoropyrimidine- and platinum-based chemo AND the tumor expressed PD-L1 great than or equal. Esophageal or Gastroesophageal Junction cancer (residual disease post-surgery and preoperative chemoradiation): received neoadjuvant chemoradiation AND complete surgical resection of esophageal or gastroesophageal junction AND has residual pathologic disease AND given as subsequent monotherapy. NSCLC- Neoadjuvant Therapy: early stage NSCLC AND has resectable disease (tumors greater than 4 cm or node positive) AND using in neoadjuvant setting, in combo with platinum-doublet chemo.
Part B Prerequisite	0

OPDUALAG - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Disease progression while on or following prior anti-PD-1/PD-L1 combination therapy (e.g., nivolumab with ipilimumab). Members on concomitant Zelboraf (vemurafenib), Tafinlar (dabrafenib), Mekinist (trametinib) or Cotellic (cobimetinib) therapy. Safety and efficacy have not been established.
Required Medical Information	Melanoma: Unresectable or metastatic melanoma: The member must have a diagnosis of unresectable or metastatic melanoma AND Opdualag is administered as monotherapy.
Age Restriction	The member must be 12 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 months duration
Other Criteria	
Part B Prerequisite	0

OPIPZA - FILM, MEDICATED (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Schizophrenia: The member has clinically diagnosed schizophrenia AND The member has tried or cannot use at least two of the following generic atypical antipsychotics: risperidone, olanzapine, quetiapine, ziprasidone, aripiprazole, or lurasidone. Major Depressive Disorder (MDD): The member has clinically diagnosed major depressive disorder (MDD) AND The member has tried or cannot use generic aripiprazole in combination with at least one product considered antidepressant therapy (ADT) AND Opipza will be used as adjunctive or add-on treatment to ADT and not as monotherapy. Irritability associated with Autistic Disorder or Tourette's Disorder: The member has one of the following clinically diagnosed conditions: Irritability associated with autistic disorder OR Tourette's disorder.
Age Restriction	Schizophrenia: The member is 13 years of age or older. MDD: The member is 18 years of age or older. Irritability associated with Autistic Disorder or Tourette's Disorder: The member is 6 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

ORGOVYX - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Pediatric members less than 18 years old.
Required Medical Information	Prostate Cancer: The member has a diagnosis of advanced prostate cancer or has a high risk of disease recurrence.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ORSERDU - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Estrogen Receptor (ER)- positive Breast Cancer: The member has ER-positive, HER2-negative advanced or metastatic breast cancer AND the breast cancer has documented ESR1-mutation as determined by FDA approved test AND the member has progressive disease following at least one prior line endocrine therapy (e.g., fulvestrant, CDK 4/6 inhibitor) AND Orserdu (elacestrant) is given as single agent as subsequent therapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

OTEZLA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	Member is 18 years of age or older for treatment of oral ulcers associated with Behcet's Disease and mild plaque psoriasis. Member is 6 years of age or older for treatment of moderate to severe plaque psoriasis and psoriatic arthritis.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

OTEZLA STARTER - TABLET, DOSE PACK

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	Member is 18 years of age or older for treatment of oral ulcers associated with Behcet's Disease and mild plaque psoriasis. Member is 6 years of age or older for treatment of moderate to severe plaque psoriasis and psoriatic arthritis.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

OTEZLA XR - TABLET, EXTENDED RELEASE 24 HR

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	Member is 18 years of age or older for treatment of oral ulcers associated with Behcet's Disease and mild plaque psoriasis. Member is 6 years of age or older for treatment of moderate to severe plaque psoriasis and psoriatic arthritis.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

OTEZLA XR INITIATION - TABLET AND TABLET ER DOSE PACK

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	Member is 18 years of age or older for treatment of oral ulcers associated with Behcet's Disease and mild plaque psoriasis. Member is 6 years of age or older for treatment of moderate to severe plaque psoriasis and psoriatic arthritis.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

OTULFI - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	Moderate to severe chronic plaque psoriasis and psoriatic arthritis: The member must be 6 years of age or older. For Ulcerative Colitis: The member must be 18 years of age or older. For Crohn's Disease: The member must be 2 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

OZEMPIC - PEN INJECTOR (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member is using for weight loss.
Required Medical Information	Diabetes Mellitus. The member has a diagnosis of type 2 diabetes mellitus.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

PANRETIN - GEL (GRAM)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	AIDS-related Kaposi's sarcoma: The member has a diagnosis of AIDS-related Kaposi's sarcoma AND systemic therapy is not required.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

pazopanib - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on previous pazopanib therapy.
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

PEGASYS - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Chronic Hepatitis B - Adults: The member must have a diagnosis of chronic hepatitis B AND The member must have compensated liver disease AND The member must have evidence of viral replication AND The member must have evidence of liver inflammation AND The member has tried or cannot use both of the following: tenofovir disoproxil fumarate and entecavir. Chronic Hepatitis B - Pediatrics: The member must have a diagnosis of chronic hepatitis B AND The member must be non-cirrhotic AND The member must be HBeAg-positive AND The member must have evidence of viral replication AND The member must have elevations in serum alanine aminotransferase (ALT) AND the member has tried or cannot use both of the following: tenofovir disoproxil fumarate and entecavir. Chronic Hepatitis C - Adults: The member must have a diagnosis of chronic hepatitis C AND The member has compensated liver disease AND Pegasys (peginterferon alpha-2a) will be taken in combination with at least 1 other medication indicated for the treatment of chronic Hepatitis C AND the member cannot use generic sofosbuvir/velpatasvir.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Pending CMS Review
Other Criteria	Chronic Hepatitis C - Pediatrics: The member must have a diagnosis of chronic hepatitis C AND The member has compensated liver disease AND Pegasys (peginterferon alpha-2a) will be taken in combination with ribavirin AND the member cannot use generic sofosbuvir/velpatasvir.
Part B Prerequisite	0

PEMAZYRE - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member experienced disease progression on Pemazyre (pemigatinib)
Required Medical Information	Cholangiocarcinoma: The member has unresectable locally advanced or metastatic cholangiocarcinoma and the disease is fibroblast growth factor receptor 2 (FGFR2) fusion or other rearrangement as detected by an FDA-approved test and the member has received prior treatment AND Pemazyre (pemigatinib) is given as a single agent for subsequent therapy. Relapsed or refractory myeloid/lymphoid neoplasms: the member has a diagnosis of relapsed or refractory myeloid/lymphoid neoplasms (MLNs) AND MLNs documented as fibroblast growth factor receptor 1 (FGFR1) rearrangement.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Six months duration
Other Criteria	
Part B Prerequisite	0

perampanel - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Partial-Onset (Focal) Seizures: The member has a diagnosis of partial-onset (focal) seizures AND Member has been unable to achieve seizure control with at least TWO other drugs for controlling partial-onset seizures (e.g., levetiracetam, lamotrigine, carbamazepine, zonisamide, divalproex, gabapentin or topiramate). Adjunctive treatment of generalized tonic-clonic seizures: The member has a diagnosis of generalized tonic-clonic seizures AND Member will use Fycompa (perampanel) in combination with at least one other drug for controlling seizures AND Member has been unable to achieve seizure control with at least TWO other drugs for controlling generalized tonic-clonic seizures (e.g. lamotrigine, topiramate, carbamazepine, gabapentin, divalproex).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

PERJETA - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member exceeds a total treatment of 52 weeks or 18 treatment cycles (applicable to neoadjuvant and/or adjuvant treatment).
Required Medical Information	Metastatic Breast Cancer. The member has a diagnosis of metastatic breast cancer and HER2 (human epidermal growth factor receptor2) positive disease (e.g., defined as IHC 3+ or IHC2+/ISH positive) AND one of the following applies: will be receiving Perjeta (pertuzumab) in combination with trastuzumab product and docetaxel or paclitaxel and has not received prior anti-HER2 therapy or chemotherapy for metastatic disease OR The member will be receiving Perjeta (pertuzumab) in combination with a trastuzumab product with or without cytotoxic therapy as second or subsequent line therapy in members previously treated with chemotherapy and a trastuzumab product in the absence of prior Perjeta (pertuzumab). Early Stage Breast Cancer. The member has a diagnosis of locally advanced, inflammatory, or early stage breast cancer (either greater than 2 cm in diameter or node positive) and HER2 positive disease (e.g., defined as IHC 3+ or IHC2+/ISH positive) AND Perjeta (pertuzumab) will be used as neoadjuvant treatment as part of a complete treatment regimen and one of the following applies: in combination with trastuzumab product and docetaxel or paclitaxel (after completion of combination of doxorubicin plus cyclophosphamide regimen) or in combination with TCH (docetaxel, carboplatin, and trastuzumab product) OR The member has a diagnosis of early stage HER2 (e.g., defined as IHC 3+ or IHC2+/ISH positive) positive breast cancer at high risk of recurrence (e.g., node positive disease, hormone receptor negative, T2 non-metastatic disease) AND Perjeta (pertuzumab) will be used as adjuvant therapy and one of the following applies: combination with trastuzumab product and paclitaxel or docetaxel (following doxorubicin plus cyclophosphamide regimen) or docetaxel plus carboplatin.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

pimecrolimus - CREAM (GRAM)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Must have a diagnosis of atopic dermatitis and have had previous treatment with one of the following topical generic products: triamcinolone 0.025%, 0.1%, 0.5%, mometasone, betamethasone dipropionate.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

PIQRAY - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members have severe hypersensitivity to Piqray (alpelisib). Members has experienced disease progression on PIK3CA inhibitors (e.g., alpelisib).
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

pirfenidone - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

pomalidomide - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression while on pomalidomide.
Required Medical Information	For brand Pomalyst requests: Member must have an intolerance or contraindication to generic pomalidomide AND member must meet clinical criteria: Multiple Myeloma: The member has a diagnosis of Multiple Myeloma AND One of the following situations apply: The member has received one prior line of therapy and will be using pomalidomide in combination with one of the following regimens: Bortezomib and dexamethasone, Carfilzomib and dexamethasone, or Darzalex (daratumumab), carfilzomib and dexamethasone OR The member has received at least one prior line of therapy, which included an immunomodulatory agent (e.g., lenalidomide or thalidomide) and a proteasome inhibitor (e.g., bortezomib or carfilzomib) AND The member demonstrated disease progression on or within 60 days of completion of the last therapy regimen and will be using pomalidomide in combination with one of the following regimens: Ninlaro (ixazomib) and dexamethasone, Dexamethasone, Xpovio (selinexor) and dexamethasone, Cyclophosphamide and dexamethasone, Single agent pomalidomide OR The member will be using pomalidomide in combination with one of the following regimens: Darzalex (daratumumab) and dexamethasone, Empliciti (elotuzumab) and dexamethasone, Sarclisa (isatuximab) and dexamethasone. Omission of corticosteroid from regimen is allowed if intolerance/contraindication. Kaposi Sarcoma: The member has a diagnosis of AIDS-related Kaposi sarcoma after failure of highly active antiretroviral therapy OR The member has a diagnosis of Kaposi sarcoma that is HIV-negative.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

POMALYST - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression while on pomalidomide.
Required Medical Information	For brand Pomalyst requests: Member must have an intolerance or contraindication to generic pomalidomide AND member must meet clinical criteria: Multiple Myeloma: The member has a diagnosis of Multiple Myeloma AND One of the following situations apply: The member has received one prior line of therapy and will be using pomalidomide in combination with one of the following regimens: Bortezomib and dexamethasone, Carfilzomib and dexamethasone, or Darzalex (daratumumab), carfilzomib and dexamethasone OR The member has received at least one prior line of therapy, which included an immunomodulatory agent (e.g., lenalidomide or thalidomide) and a proteasome inhibitor (e.g., bortezomib or carfilzomib) AND The member demonstrated disease progression on or within 60 days of completion of the last therapy regimen and will be using pomalidomide in combination with one of the following regimens: Ninlaro (ixazomib) and dexamethasone, Dexamethasone, Xpovio (selinexor) and dexamethasone, Cyclophosphamide and dexamethasone, Single agent pomalidomide OR The member will be using pomalidomide in combination with one of the following regimens: Darzalex (daratumumab) and dexamethasone, Empliciti (elotuzumab) and dexamethasone, Sarclisa (isatuximab) and dexamethasone. Omission of corticosteroid from regimen is allowed if intolerance/contraindication. Kaposi Sarcoma: The member has a diagnosis of AIDS-related Kaposi sarcoma after failure of highly active antiretroviral therapy OR The member has a diagnosis of Kaposi sarcoma that is HIV-negative.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

posaconazole - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Prophylaxis against Invasive Aspergillus and Candida Infections. The member must be using it for prophylaxis against invasive Aspergillus or Candida infections, and The member must be severely immunocompromised (such as hematopoietic stem cell transplant recipient with graft-vs-host disease, or neutropenic patients with acute myelogenous leukemia (AML) or myelodysplastic syndromes (MDS). Treatment of invasive Aspergillus or fungal infections caused by Fusarium and/or Zygomycetes. The member must have documentation for treatment of invasive Aspergillus or fungal infections caused by Fusarium and/or Zygomycetes, and The member must have documented resistant strains of or clinically refractory to standard antifungal agents (e.g. voriconazole, itraconazole) or those who can not receive other antifungal agents due to potential toxicities, intolerance, or contraindications. Treatment of Esophageal Candidiasis. The member must have a diagnosis for esophageal candidiasis and The member has a documented inadequate response/refractory or intolerant to itraconazole and fluconazole.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

POTELIGEO - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	NA
Exclusion Criteria	Member has experienced disease progression while on or following Poteligeo (mogamulizumab-kpkc).
Required Medical Information	Mycosis fungoides or Sézary syndrome: The member has a diagnosis of mycosis fungoides or Sézary syndrome AND The member has relapsed or refractory disease AND The member will be using Poteligeo (mogamulizumab-kpkc) as the sole systemic therapy.
Age Restriction	NA
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 Months Duration
Other Criteria	NA
Part B Prerequisite	0

PREMARIN - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Member must have a previous treatment, intolerance or contraindication to an AB-rated generic equivalent product.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

PREVYMIS - PELLETS IN PACKET (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	Prophylaxis of CMV Infection and Disease in CMV Seropositive Recipients [R+] of an Allogeneic Hematopoietic Stem Cell Transplant (HSCT): The member must be at least 6 months of age or older. Prophylaxis of CMV disease in kidney transplant recipients at high risk (Donor CMV seropositive/Recipient CMV seronegative [D+/R-]): The member must be at least 12 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	12 Months Duration
Other Criteria	
Part B Prerequisite	0

QINLOCK - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Gastrointestinal Stromal Tumor (GIST): The member has a diagnosis of advanced GIST AND The member has received prior therapy with three or more kinase inhibitors, including imatinib AND Qinlock (ripretinib) is being used as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 months duration
Other Criteria	
Part B Prerequisite	0

quinine sulfate - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Prolongation of QT interval. Myasthenia gravis. Optic neuritis.
Required Medical Information	Plasmodium Falciparum Malaria: Diagnosis of uncomplicated chloroquine-resistant Plasmodium falciparum malaria.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

QULIPTA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Migraine Prevention: Will be using Qulipta (atogepant) for the preventive treatment of migraines.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

RADICAVA - INTRAVENOUS SOLUTION, PIGGYBACK (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

RADICAVA ORS - SUSPENSION, ORAL (FINAL DOSE FORM)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

RADICAVA ORS STARTER KIT SUSP - SUSPENSION, ORAL (FINAL DOSE FORM)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

RASONQUE - TABLET

PA Criteria	Criteria Details
Off-Label Uses	Pending CMS Review
Exclusion Criteria	Pending CMS Review
Required Medical Information	Pending CMS Review
Age Restriction	Pending CMS Review
Prescriber Restriction	Pending CMS Review
Coverage Duration	Pending CMS Review
Other Criteria	Pending CMS Review
Part B Prerequisite	Pending CMS Review

REPATHA PUSHTRONEX - WEARABLE INJECTOR

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Hypercholesterolemia: Member has a diagnosis of hypercholesterolemia (Note: includes heterozygous familial hypercholesterolemia [HeFH], homozygous familial hypercholesterolemia [HoFH], pure hypercholesterolemia, and mixed hyperlipidemia) AND Has tried or cannot use a high-intensity statin. Cardiovascular Event Prevention: The member must have an increased risk of major adverse cardiovascular events (e.g. acute coronary syndromes, history of myocardial infarction, stable or unstable angina, coronary or other arterial revascularization, stroke, transient ischemic attack or peripheral arterial disease, all of presumed atherosclerotic origin) AND Has tried or cannot use a high-intensity statin.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

REPATHA SURECLICK - PEN INJECTOR (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Hypercholesterolemia: Member has a diagnosis of hypercholesterolemia (Note: includes heterozygous familial hypercholesterolemia [HeFH], homozygous familial hypercholesterolemia [HoFH], pure hypercholesterolemia, and mixed hyperlipidemia) AND Has tried or cannot use a high-intensity statin. Cardiovascular Event Prevention: The member must have an increased risk of major adverse cardiovascular events (e.g. acute coronary syndromes, history of myocardial infarction, stable or unstable angina, coronary or other arterial revascularization, stroke, transient ischemic attack or peripheral arterial disease, all of presumed atherosclerotic origin) AND Has tried or cannot use a high-intensity statin.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

REPATHA SYRINGE - SYRINGE (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Hypercholesterolemia: Member has a diagnosis of hypercholesterolemia (Note: includes heterozygous familial hypercholesterolemia [HeFH], homozygous familial hypercholesterolemia [HoFH], pure hypercholesterolemia, and mixed hyperlipidemia) AND Has tried or cannot use a high-intensity statin. Cardiovascular Event Prevention: The member must have an increased risk of major adverse cardiovascular events (e.g. acute coronary syndromes, history of myocardial infarction, stable or unstable angina, coronary or other arterial revascularization, stroke, transient ischemic attack or peripheral arterial disease, all of presumed atherosclerotic origin) AND Has tried or cannot use a high-intensity statin.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

RETACRIT - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Initial: Plan year duration. Reauth: Plan year duration.
Other Criteria	Pending CMS Review
Part B Prerequisite	0

RETEVMO - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member experiences disease progression on Retevmo
Required Medical Information	Non-small cell lung cancer. The member has a diagnosis of locally advanced or metastatic non-small cell lung cancer AND The disease is documented RET fusion positive AND Retevmo is being used as monotherapy. Medullary Thyroid cancer. The member has a diagnosis of metastatic or advanced medullary thyroid cancer AND The disease is documented RET mutant AND Retevmo is being used as a single agent for systemic therapy. Thyroid cancer. The member has a diagnosis of metastatic or advanced thyroid cancer AND The disease is documented RET fusion positive AND The disease is radioactive iodine refractory (if radioactive iodine is appropriate) AND Retevmo is being used as a single agent for systemic therapy. RET fusion-positive Solid Tumors: the member has locally advanced or metastatic solid tumors AND the solid tumors have documented rearranged during transfection (RET) gene fusion positive AND the member has progressed on or following prior systemic treatment or has no satisfactory alternative treatment options AND Retevmo (selpercatinib) is being administered as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 months duration
Other Criteria	
Part B Prerequisite	0

REVCOVI - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Adenosine deaminase severe combined immune deficiency (ADA-SCID): The member has a diagnosis of adenosine deaminase severe combined immune deficiency.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

REVUFORJ - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression while on or following a menin inhibitor (e.g., Revuforj (revumenib), Komzifti (ziftomenib)).
Required Medical Information	Acute Leukemia: The member has a diagnosis of acute leukemia AND the member has relapsed or refractory disease AND One of the following: the member has documented lysine methyltransferase 2A (KMT2A) mutation as determined by an FDA authorized test OR the member has acute myeloid leukemia with a susceptible nucleophosmin 1 (NPM1) mutation and there are no satisfactory alternative treatment options AND there is a medical reason why Komzifti (ziftomenib) cannot be initiated or continued.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

REXULTI - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Major depressive disorder: The member has a clinical diagnosis of major depressive disorder (MDD) AND The member has tried or cannot use at least one generic oral atypical antipsychotic therapy AND Rexulti (brexpiprazole) will be used with antidepressant therapy and not as monotherapy. Schizophrenia: The member must have clinically diagnosed schizophrenia AND The member has tried or cannot use at least one generic oral atypical antipsychotic therapy. Agitation Associated with Dementia Due to Alzheimer's Disease: The member must have clinically diagnosed agitation associated with dementia due to Alzheimer's disease AND Rexulti (brexpiprazole) will not be used as an as needed (prn) treatment for agitation.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

REZDIFFRA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	The member is at least 18 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

REZLIDHIA - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member has experienced disease progression while on or following Rezlidhia (olutasidenib).
Required Medical Information	Acute Myeloid Leukemia - Relapsed/Refractory: The member has a diagnosis of acute myeloid leukemia (AML) AND The member has relapsed or refractory disease AND The member has a documented IDH1 mutation AND The member will be using Rezlidhia (olutasidenib) as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

RIABNI - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Chronic Lymphocytic Leukemia: High dose CLL therapies (doses greater than 500mg/m²). The length of maintenance therapy exceeds 2 years for chronic lymphocytic leukemia. Non-Hodgkin's Lymphoma (CD-20 positive/B-cell): The member will be using rituximab as maintenance therapy for diffuse large B cell lymphoma (DLBCL). The length of maintenance therapy exceeds 2 years for low-grade non-Hodgkins lymphoma (e.g. follicular lymphoma, marginal zone lymphoma).
Required Medical Information	For requests for Rituxan or Truxima: member must have intolerance or contraindication with Ruxience or Riabni and meet the criteria below. Chronic Lymphocytic Leukemia. The member has a diagnosis of CLL. Non-Hodgkin's Lymphoma (CD-20 positive/B-Cell). The member has a diagnosis of CD-20 positive/B-cell Non-Hodgkin's lymphoma. Hodgkin's Disease (Hodgkin's Lymphoma). The member has a diagnosis of Hodgkin's Disease. The member will be using rituximab product for primary treatment or for relapsed or progressive disease AND disease has confirmed CD20 positivity. Rheumatoid Arthritis. The member has a diagnosis of moderately- to severely-active rheumatoid arthritis. The member has had previous treatment with, contraindication, or intolerance with two of the following: a preferred infliximab product (i.e., Remicade, Avsola, infliximab), Simponi Aria, or Tyenne. The member must be on concomitant treatment with methotrexate during rituximab product therapy, unless contraindicated or intolerant to methotrexate.
Age Restriction	Wegener's Granulomatosis (WG) and Microscopic Polyangiitis (MPA): The member is 2 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration

Other Criteria	Waldenstram's macroglobulinemia. The member must have a diagnosis of Waldenstram's macroglobulinemia. Post-transplant lymphoproliferative disorder. The member has a diagnosis of Post-transplant Lymphoproliferative disease. Immune or Idiopathic Thrombocytopenic Purpura. The member must have a diagnosis of refractory primary immune or idiopathic thrombocytopenic purpura. Member has not had a splenectomy, but has had an insufficient response or is intolerant to corticosteroids, AND immunoglobulins (IVIG), OR has had a splenectomy with an inadequate response or is intolerant to procedure AND had an insufficient response or is intolerant to post-splenectomy corticosteroids. Refractory response is characterized as EITHER: Platelet count less than 25,000/microliter OR Active bleeding due to inadequate platelet function. The member must have had an inadequate response to post-splenectomy corticosteroid therapy for four consecutive weeks within the last three months. Diagnosis of Wegener’s Granulomatosis OR Microscopic Polyangiitis. Must be taking rituximab product in combination with glucocorticoids. Pemphigus Vulgaris (PV). The member must have a diagnosis of moderate to severe Pemphigus Vulgaris.
Part B Prerequisite	0

RINVOQ - TABLET, EXTENDED RELEASE 24 HR

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Rheumatoid Arthritis: The member has a diagnosis of moderate to severely active rheumatoid arthritis AND the member has tried or cannot use at least one TNF blocker (e.g., adalimumab-adbm, adalimumab-adaz, Enbrel). Psoriatic Arthritis: The member has a diagnosis of active psoriatic arthritis AND the member has tried or cannot use at least one TNF blocker (e.g., adalimumab-adbm, adalimumab-adaz, Enbrel). Atopic Dermatitis: The member has a diagnosis of moderate to severe atopic dermatitis AND the member has tried or cannot use at least one other systemic therapy. Ulcerative Colitis: the member has a diagnosis of moderately to severely active ulcerative colitis AND the member has tried or cannot use at least one TNF blocker (e.g., adalimumab-adbm, adalimumab-adaz) or at least one approved systemic therapy if a TNF blocker cannot be used. Ankylosing Spondylitis: the member has a diagnosis of active ankylosing spondylitis AND the member has tried or cannot use at least one TNF blocker (e.g., adalimumab-adbm, adalimumab-adaz, Enbrel). Non-radiographic Axial Spondylarthritis: the member has a diagnosis of non-radiographic axial spondylarthritis with signs of inflammation AND the member has tried or cannot use at least one TNF blocker (e.g., adalimumab-adbm, adalimumab-adaz, Enbrel). Crohn's Disease: the member has a diagnosis of moderately to severely active crohn's disease AND the member has tried or cannot use at least one TNF blocker (e.g., adalimumab-adbm, adalimumab-adaz) or at least one approved systemic therapy if a TNF blocker cannot be used.
Age Restriction	RA, UC, ankylosing spondylitis, Non-radiographic axial spondylarthritis, Giant Cell Arteritis and Crohn's Disease: The member is 18 years of age or older. Atopic Dermatitis: the member is 12 years of age or older. Psoriatic arthritis and polyarticular juvenile idiopathic arthritis: the member is 2 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	Polyarticular Juvenile Idiopathic Arthritis: The member has a diagnosis of active polyarticular juvenile idiopathic arthrits AND the member has tried or cannot use at least one TNF blocker (e.g. adalimumab-adbm, adalimumab-adaz, Enbrel). Giant Cell Arteritis: the member has a diagnosis of giant cell arteritis.

Part B Prerequisite	0

RINVOQ LQ - SOLUTION, ORAL

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Psoriatic Arthritis: The member has a diagnosis of active psoriatic arthritis AND the member has tried or cannot use at least one TNF blocker (e.g., adalimumab-adbm, adalimumab-adaz, Enbrel). Polyarticular Juvenile Idiopathic Arthritis: The member has a diagnosis of active polyarticular juvenile idiopathic arthritis AND the member has tried or cannot use at least one TNF blocker (e.g., adalimumab-adbm, adalimumab-adaz, Enbrel).
Age Restriction	The member is 2 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

riociguat - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Chronic Thromboembolic Pulmonary Hypertension (CTEPH). The member must have a diagnosis of Chronic Thromboembolic Pulmonary Hypertension (CTEPH) (WHO Group 4) AND The member must have CTEPH classified as inoperable or persistent/recurrent after surgical treatment (i.e. pulmonary endarterectomy). Pulmonary Arterial Hypertension (PAH). The member must have a diagnosis of pulmonary arterial hypertension (WHO Group 1) confirmed by right heart catheterization.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ROMVIMZA - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Symptomatic Tenosynovial Giant Cell Tumor: Member has symptomatic tenosynovial giant cell tumor (TGCT) AND Surgical resection will potentially cause worsening functional limitation or severe morbidity.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

ROZLYTREK - PELLETS IN PACKET (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Non-Small Cell Lung Cancer (NSCLC): The member has a diagnosis of metastatic or recurrent non-small cell lung cancer (NSCLC) AND the member has disease which is ROS1-positive. Solid Tumors: the member has a diagnosis of solid tumors which are metastatic OR The member is not a candidate for surgical resection AND The member has a documented neurotrophic tyrosine receptor kinase (NTRK) gene fusion without a known acquired resistance mutation AND The member's disease has progressed following treatment or does not have satisfactory alternative therapy options.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

RUBRACA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on or following PARP inhibitor therapy [e.g., Rubraca (rucaparib), Lynparza (olaparib), Zejula(niraparib)].
Required Medical Information	Epithelial Ovarian Cancer, Fallopian Tube, or Peritoneal Cancer Maintenance Therapy: The member has a diagnosis of epithelial ovarian cancer, fallopian tube cancer, or primary peritoneal cancer AND The member has deleterious BRCA mutation (germline and/or somatic) AND The member is in complete or partial response to their last platinum regimen AND The member will utilize Rubraca (rucaparib) as monotherapy* AND *Discontinue Bevacizumab product before initiating maintenance therapy with Rubraca. Metastatic Castration-Resistant Prostate Cancer: The member has a diagnosis metastatic castration-resistant prostate cancer (mCRPC) AND The member has documented deleterious BRCA mutation (germline and/or somatic) AND The member has had prior treatment with androgen receptor-directed therapy (e.g. abiraterone, Xtandi, Erleada, or Nubeqa) AND The member will use Rubraca (rucaparib) in combination with androgen deprivation therapy (e.g. previous bilateral orchiectomy or luteinizing hormone-releasing hormone (LHRH) analog [i.e., LHRH agonist/ antagonist]).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

rufinamide - SUSPENSION, ORAL (FINAL DOSE FORM)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Patients with familial short QT syndrome.
Required Medical Information	Lennox-Gastaut Syndrome: The member has a diagnosis of seizures associated with Lennox-Gastaut Syndrome (LGS) AND Banzel (rufinamide) will be used in combination with at least one other drug for controlling seizures AND The member has been unable to achieve seizure control with at least one other drug used for the adjunctive treatment of Lennox-Gastaut syndrome (e.g., lamotrigine, topiramate).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

RUXIENCE - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Chronic Lymphocytic Leukemia: High dose CLL therapies (doses greater than 500mg/m²). The length of maintenance therapy exceeds 2 years for chronic lymphocytic leukemia. Non-Hodgkin's Lymphoma (CD-20 positive/B-cell): The member will be using rituximab as maintenance therapy for diffuse large B cell lymphoma (DLBCL). The length of maintenance therapy exceeds 2 years for low-grade non-Hodgkins lymphoma (e.g. follicular lymphoma, marginal zone lymphoma).
Required Medical Information	For requests for Rituxan or Truxima: member must have intolerance or contraindication with Ruxience or Riabni and meet the criteria below. Chronic Lymphocytic Leukemia. The member has a diagnosis of CLL. Non-Hodgkin's Lymphoma (CD-20 positive/B-Cell). The member has a diagnosis of CD-20 positive/B-cell Non-Hodgkin's lymphoma. Hodgkin's Disease (Hodgkin's Lymphoma). The member has a diagnosis of Hodgkin's Disease. The member will be using rituximab product for primary treatment or for relapsed or progressive disease AND disease has confirmed CD20 positivity. Rheumatoid Arthritis. The member has a diagnosis of moderately- to severely-active rheumatoid arthritis. The member has had previous treatment with, contraindication, or intolerance with two of the following: a preferred infliximab product (i.e., Remicade, Avsola, infliximab), Simponi Aria, or Tyenne. The member must be on concomitant treatment with methotrexate during rituximab product therapy, unless contraindicated or intolerant to methotrexate.
Age Restriction	Wegener's Granulomatosis (WG) and Microscopic Polyangiitis (MPA): The member is 2 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration

Other Criteria	Waldenstram's macroglobulinemia. The member must have a diagnosis of Waldenstram's macroglobulinemia. Post-transplant lymphoproliferative disorder. The member has a diagnosis of Post-transplant Lymphoproliferative disease. Immune or Idiopathic Thrombocytopenic Purpura. The member must have a diagnosis of refractory primary immune or idiopathic thrombocytopenic purpura. Member has not had a splenectomy, but has had an insufficient response or is intolerant to corticosteroids, AND immunoglobulins (IVIG), OR has had a splenectomy with an inadequate response or is intolerant to procedure AND had an insufficient response or is intolerant to post-splenectomy corticosteroids. Refractory response is characterized as EITHER: Platelet count less than 25,000/microliter OR Active bleeding due to inadequate platelet function. The member must have had an inadequate response to post-splenectomy corticosteroid therapy for four consecutive weeks within the last three months. Diagnosis of Wegener’s Granulomatosis OR Microscopic Polyangiitis. Must be taking rituximab product in combination with glucocorticoids. Pemphigus Vulgaris (PV). The member must have a diagnosis of moderate to severe Pemphigus Vulgaris.
Part B Prerequisite	0

RYBELSUS - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member is using for weight loss.
Required Medical Information	Diabetes Mellitus. The member has a diagnosis of type 2 diabetes mellitus.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

RYBREVANT - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members who have disease progression on an amivantamab product (does not apply to NSCLC EGFR Exon 19 Deletion or Exon 21 L858R Mutated [Lazcluze (lazertinib) Combination]).
Required Medical Information	Non-small cell lung cancer (NSCLC) [Monotherapy]: The member has a diagnosis of locally advanced or metastatic non-small cell lung cancer (NSCLC) AND The NSCLC has documented epidermal growth factor receptor (EGFR) exon 20 insertion mutations (e.g. as detected by a FDA-approved test) AND the member has documented disease progression on prior platinum-based chemotherapy AND The amivantamab product (i.e., Rybrevant, Rybrevant Faspro) will be given as monotherapy. Non-small cell lung cancer (NSCLC) [Combination Therapy]: The member has a diagnosis of locally advanced or metastatic non-small cell lung cancer (NSCLC) AND The NSCLC has documented epidermal growth factor receptor (EGFR) exon 20 insertion mutations (e.g. as detected by a FDA-approved test) AND The amivantamab product will be given in combination with carboplatin and pemetrexed in the first line setting. NSCLC EGFR Exon 19 Deletion or Exon 21 L858R Mutated [Lazcluze (lazertinib) Combination]: The member has a diagnosis of locally advanced or metastatic non-small cell lung cancer (NSCLC) AND The NSCLC has documented epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R substitution mutations AND The amivantamab product will be given in combination with Lazcluze (lazertinib) in the first-line setting or as subsequent therapy (if not previously given). NSCLC EGFR Exon 19 Deletion or Exon 21 L858R Mutated [Other Combination Therapy]: The member has a diagnosis of locally advanced or metastatic non-small cell lung cancer (NSCLC) AND The NSCLC has documented epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R substitution mutations AND The member has experienced disease progression on or after an EGFR tyrosine kinase inhibitor [e.g. Tagrisso (osimeritinib)] AND The amivantamab product will be given in combination with carboplatin and pemetrexed as subsequent line therapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	

Part B Prerequisite	0

RYBREVANT FASPRO - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members who have disease progression on an amivantamab product (does not apply to NSCLC EGFR Exon 19 Deletion or Exon 21 L858R Mutated [Lazcluze (lazertinib) Combination]).
Required Medical Information	Non-small cell lung cancer (NSCLC) [Monotherapy]: The member has a diagnosis of locally advanced or metastatic non-small cell lung cancer (NSCLC) AND The NSCLC has documented epidermal growth factor receptor (EGFR) exon 20 insertion mutations (e.g. as detected by a FDA-approved test) AND the member has documented disease progression on prior platinum-based chemotherapy AND The amivantamab product (i.e., Rybrevant, Rybrevant Faspro) will be given as monotherapy. Non-small cell lung cancer (NSCLC) [Combination Therapy]: The member has a diagnosis of locally advanced or metastatic non-small cell lung cancer (NSCLC) AND The NSCLC has documented epidermal growth factor receptor (EGFR) exon 20 insertion mutations (e.g. as detected by a FDA-approved test) AND The amivantamab product will be given in combination with carboplatin and pemetrexed in the first line setting. NSCLC EGFR Exon 19 Deletion or Exon 21 L858R Mutated [Lazcluze (lazertinib) Combination]: The member has a diagnosis of locally advanced or metastatic non-small cell lung cancer (NSCLC) AND The NSCLC has documented epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R substitution mutations AND The amivantamab product will be given in combination with Lazcluze (lazertinib) in the first-line setting or as subsequent therapy (if not previously given). NSCLC EGFR Exon 19 Deletion or Exon 21 L858R Mutated [Other Combination Therapy]: The member has a diagnosis of locally advanced or metastatic non-small cell lung cancer (NSCLC) AND The NSCLC has documented epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R substitution mutations AND The member has experienced disease progression on or after an EGFR tyrosine kinase inhibitor [e.g. Tagrisso (osimeritinib)] AND The amivantamab product will be given in combination with carboplatin and pemetrexed as subsequent line therapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	

Part B Prerequisite	0

RYDAPT - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	AML newly diagnosed: Members that are using Rydapt(midostaurin) for post-consolidation therapy. AML relapsed/refractory: Members that have experienced disease progression while on or following Rydapt (midostaurin), Members that are using Rydapt(midostaurin) for post-consolidation therapy. Systemic Mastocytosis: Members that have experienced disease progression while on or following Rydapt (midostaurin).
Required Medical Information	Acute Myeloid Leukemia-Newly diagnosed: The member has newly diagnosed acute myeloid leukemia (AML) AND The member has documented FLT3 mutation-positive disease AND The member will be using Rydapt (midostaurin) in combination with standard cytarabine and daunorubicin induction and cytarabine consolidation chemotherapy. Systemic Mastocytosis: The member has a diagnosis of aggressive systemic mastocytosis (ASM), systemic mastocytosis with associated hematologic neoplasm (SM-AHN), or mast cell leukemia (MCL). Acute Myeloid Leukemia - Relapsed/Refractory: The member has relapsed or refractory acute myeloid leukemia (AML) AND The member has documented FLT3 mutation-positive disease AND The member will be using Rydapt (midostaurin) as a component of repeating the initial successful induction regimen, if late relapse (relapse occurring later than 12 months).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

SANTYL - OINTMENT (GRAM)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Wound Debridement: Member has a chronic dermal ulcer (e.g. pressure ulcer, venous ulcer, diabetic ulcer) or severely burned areas AND Member will be using Santyl (collagenase) for wound debridement.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

sapropterin - TABLET, SOLUBLE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Phenylketonuria (PKU): The member has a diagnosis of PKU.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

SARCLISA - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression while on or following an anti-CD38 inhibitor (e.g. daratumumab, isatuximab-irfc).
Required Medical Information	Multiple myeloma (third line). The member has a diagnosis of multiple myeloma AND The member will be using isatuximab-irfc in combination with Pomalyst (pomalidomide) and dexamethasone (Omission of corticosteroid from regimen is allowed if intolerance/contraindication) AND The member has received at least two prior therapies including Revlimid (lenalidomide) and a proteasome inhibitor (e.g. bortezomib, carfilzomib). Multiple Myeloma (relapsed or refractory): The member has a diagnosis of relapsed or refractory multiple myeloma AND The member will be using isatuximab-irfc in combination with carfilzomib and dexamethasone (Omission of corticosteroid from regimen is allowed if intolerance/contraindication) AND The member has received at least one prior therapy. Newly Diagnosed Multiple Myeloma: The member has newly diagnosed multiple myeloma AND The member is ineligible for autologous stem cell transplant AND The member will be using isatuximab-irfc in combination with bortezomib, lenalidomide, and dexamethasone.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

SARCLISA ESCENA - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression while on or following an anti-CD38 inhibitor (e.g. daratumumab, isatuximab-irfc).
Required Medical Information	Multiple myeloma (third line). The member has a diagnosis of multiple myeloma AND The member will be using isatuximab-irfc in combination with Pomalyst (pomalidomide) and dexamethasone (Omission of corticosteroid from regimen is allowed if intolerance/contraindication) AND The member has received at least two prior therapies including Revlimid (lenalidomide) and a proteasome inhibitor (e.g. bortezomib, carfilzomib). Multiple Myeloma (relapsed or refractory): The member has a diagnosis of relapsed or refractory multiple myeloma AND The member will be using isatuximab-irfc in combination with carfilzomib and dexamethasone (Omission of corticosteroid from regimen is allowed if intolerance/contraindication) AND The member has received at least one prior therapy. Newly Diagnosed Multiple Myeloma: The member has newly diagnosed multiple myeloma AND The member is ineligible for autologous stem cell transplant AND The member will be using isatuximab-irfc in combination with bortezomib, lenalidomide, and dexamethasone.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 Months Duration
Other Criteria	
Part B Prerequisite	0

SCEMBLIX - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression while on Scemblix (asciminib).
Required Medical Information	Chronic Myeloid Leukemia (chronic phase): The member has a diagnosis of Philadelphia chromosome-positive chronic myeloid leukemia (Ph+ CML) in chronic phase AND One of the following applies: The member has newly diagnosed disease OR The member has previously treated disease OR The member has T315I mutation.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

SECUADO - PATCH, TRANSDERMAL 24 HOURS

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Schizophrenia: The member has a diagnosis of schizophrenia. The member has tried or cannot use at least two of the following: risperidone, olanzapine, quetiapine, ziprasidone, aripiprazole, lurasidone.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

SIGNIFOR - AMPUL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Cushings disease: Diagnosis of Cushings disease AND Pituitary surgery is not an option or has not been curative AND No severe hepatic impairment (Child-Pugh C).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

SIGNIFOR LAR - VIAL (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Acromegaly: The member has a diagnosis of acromegaly AND the member has had an inadequate response to pituitary surgery or pituitary surgery is not an option AND the member has tried or cannot use generic lanreotide OR octreotide AND the member does not have severe hepatic impairment (Child-Pugh C). Cushing's disease. The member must have a diagnosis of Cushing's disease AND the member has had an inadequate response to pituitary surgery or pituitary surgery is not an option AND the member does not have severe hepatic impairment (Child-Pugh C).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

sildenafil (pulm.hypertension) - TABLET

PA Criteria	Criteria Details
Off-Label Uses	NA
Exclusion Criteria	NA
Required Medical Information	Pulmonary Arterial Hypertension(PAH): The member must have a diagnosis of pulmonary arterial hypertension (WHO Group I) confirmed by right heart catheterization.
Age Restriction	NA
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	NA
Part B Prerequisite	0

SIRTURO - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Multidrug-resistant tuberculosis (MDR-TB). The member must have a diagnosis of pulmonary multidrug-resistant tuberculosis (MDR-TB) confirmed by drug susceptibility testing (DST). Bedaquiline will be used as part of a multidrug regimen.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	24 weeks duration
Other Criteria	
Part B Prerequisite	0

SKYRIZI - SYRINGE (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Plaque Psoriasis: The member has a diagnosis of moderate to severe chronic plaque psoriasis AND The member has tried at least one conventional oral systemic treatment (e.g. methotrexate, cyclosporine), or cannot use any conventional oral systemic treatments. Psoriatic Arthritis: The member has a diagnosis of active psoriatic arthritis. Moderately to severely active Crohns disease: member has a diagnosis of moderately to severely active Crohns disease. Moderately to severely active ulcerative colitis: The member must have a diagnosis of moderately to severely active ulcerative colitis.
Age Restriction	Pending CMS Review
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

sodium oxybate - SOLUTION, ORAL

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Narcolepsy with cataplexy: The member has a diagnosis of narcolepsy with cataplexy. Narcolepsy with excessive daytime sleepiness: The member has a diagnosis of narcolepsy AND The member has excessive daytime sleepiness (EDS) associated with narcolepsy AND The member has tried or cannot use at least one CNS stimulant (e.g., methylphenidate or amphetamine salt combination immediate release) or modafinil.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	The member will be using no more than one of the following products at any given time: Xyrem (sodium oxybate), Xywav (calcium magnesium, potassium, and sodium oxybates), Lumryz (sodium oxybate), Wakix (pitolisant), or Sunosi (solriamfetol).
Part B Prerequisite	0

sofosbuvir-velpatasvir - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Chronic Hepatitis C Virus Genotypes: The member must have a diagnosis of chronic hepatitis C (HCV).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Pending CMS Review
Other Criteria	
Part B Prerequisite	0

SOMAVERT - VIAL (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

sorafenib - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on sorafenib. [Applicable to all indications except Gastrointestinal stromal tumor (GIST)]
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

STIVARGA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression while on Stivarga (regorafenib). [Applicable to all indications except Gastrointestinal stromal tumor (GIST)]
Required Medical Information	Metastatic Colorectal Cancer. The member has a diagnosis of metastatic colorectal cancer AND The member is using Stivarga (regorafenib) as monotherapy AND The member has experienced disease progression or cannot use ALL of the following therapies: fluoropyrimidine (regimens include 5-FU/capecitabine),oxaliplatin-based chemotherapy, irinotecan-based chemotherapy, and anti-VEGF therapy (e.g., bevacizumab, ziv-aflibercept) AND If the member is RAS wild-type, has experienced disease progression or cannot use an anti-EGFR therapy (e.g., cetuximab, panitumumab). Gastrointestinal Stromal Tumor. The member has a diagnosis of locally advanced, unresectable or metastatic gastrointestinal stromal tumor AND The member has experienced disease progression or cannot take imatinib mesylate and sunitinib malate. Hepatocellular Cancers: The member has a diagnosis of hepatocellular carcinoma AND Stivarga (regorafenib) is being given as monotherapy AND The member has experienced progression after first line therapy (e.g., sorafenib). Soft Tissue sarcoma. Diagnosis of advanced or metastatic soft tissue sarcoma (e.g., angiosarcoma, non-adipocytic sarcoma, pleomorphic rhabdomyosarcoma) AND Stivarga (regorafenib) is being given as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 Month Duration
Other Criteria	
Part B Prerequisite	0

STRENSIQ - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Hypophosphatasia (HPP): The member must have a diagnosis of perinatal-onset, infantile-onset, or juvenile-onset hypophosphatasia based on the following criteria: one of the following: clinical signs and symptoms supporting onset of hypophatasia prior to 18 years of age OR radiographic evidence supporting hypophosphatasia diagnoses prior to 18 years of age AND one of the following: documented gene mutation of tissue-nonspecific alkaline phosphatase (TNSALP) OR Low total serum alkaline phosphatase (ALP) activity determined by the gender- and age-specific reference range, AND Elevated urine concentration of phosphoethanolamine (PEA) determined by age-specific reference range, OR Elevated serum pyridoxal 5'-phosphate (PLP) level (normal range 5 – 50 mcg/L), OR Documented gene mutation of tissue-nonspecific alkaline phosphatase (TNSALP).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

sunitinib malate - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on sunitinib. [Applicable to all indications except Gastrointestinal stromal tumor (GIST)]. Member not to exceed a total treatment of 54 weeks [applicable to adjuvant therapy for renal cell carcinoma].
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

SYMPAZAN - FILM, MEDICATED (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Lennox-Gastaut Syndrome: The member has a diagnosis of seizures associated with Lennox-Gastaut Syndrome (LGS) AND Sympazan (clobazam film) will be used in combination with at least one other drug for controlling seizures AND the member has had prior therapy with AND has a documented contraindication (e.g., dysphagia) to BOTH generic clobazam tablet AND oral suspension formulation.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

TABRECTA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member experiences disease progression on Tabrecta (capmatinib).
Required Medical Information	Non-Small Lung Cell Cancer (NSCLC): The member has a diagnosis of metastatic NSCLC AND the disease is documented MET exon 14 skipping positive.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

tadalafil - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Benign Prostatic Hyperplasia (BPH). The member has a diagnosis of benign prostatic hyperplasia (BPH). The member has failed previous treatment or has a contraindication/intolerance to an alpha blocker (i.e. terazosin, doxazosin, tamsulosin, alfuzosin) AND a 5-alpha reductase inhibitor (i.e. finasteride, dutasteride).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

tadalafil (pulm. hypertension) - TABLET

PA Criteria	Criteria Details
Off-Label Uses	NA
Exclusion Criteria	NA
Required Medical Information	Pulmonary Arterial Hypertension (PAH). The member must have a diagnosis of pulmonary arterial hypertension (WHO Group I) confirmed by right heart catheterization.
Age Restriction	NA
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	NA
Part B Prerequisite	0

TAFINLAR - TABLET FOR SUSPENSION

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Melanoma - Unresectable or metastatic: Concomitant Yervoy (ipilimumab), Opdivo (nivolumab), Zelboraf (vemurafenib), Cotellic (cobimetinib), Braftovi (encorafenib), or Mektovi (binimetinib). Experienced disease progression on Tafinlar (dabrafenib). Experienced disease progression on prior anti-BRAF/MEK combination therapy [e.g. Cotellic (cobimetinib) with Zelboraf (vemurafenib) or Braftovi (encorafenib) with Mektovi (binimetinib)]. Anaplastic Thyroid Cancer, Metastatic Solid Tumors: Concomitant Yervoy (ipilimumab), Opdivo (nivolumab), Keytruda (pembrolizumab), Zelboraf (vemurafenib), Cotellic (cobimetinib), Braftovi (encorafenib), or Mektovi (binimetinib). Experienced disease progression on Tafinlar (dabrafenib). Experienced disease progression on prior anti-BRAF/MEK combination therapy [e.g. Cotellic (cobimetinib) with Zelboraf (vemurafenib) or Braftovi (encorafenib) with Mektovi (binimetinib)]. NSCLC: Concomitant Yervoy (ipilimumab), Opdivo (nivolumab), Keytruda (pembrolizumab), Zelboraf (vemurafenib), Cotellic (cobimetinib), Braftovi (encorafenib), or Mektovi (binimetinib). Experienced disease progression on prior anti-BRAF/MEK combination therapy [e.g. Cotellic (cobimetinib) with Zelboraf (vemurafenib) or Braftovi (encorafenib) with Mektovi (binimetinib)]. Adjuvant melanoma: Taking Tafinlar (dabrafenib) total treatment for more than 1 year. Concomitant Yervoy (ipilimumab), Opdivo (nivolumab), Keytruda (pembrolizumab), Zelboraf (vemurafenib), Cotellic (cobimetinib), Braftovi (encorafenib), or Mektovi (binimetinib). Experienced disease progression on prior anti-BRAF/MEK combination therapy [e.g. Cotellic (cobimetinib) with Zelboraf (vemurafenib) or Braftovi (encorafenib) with Mektovi (binimetinib)]. LGG: Experienced disease progression on Tafinlar (dabrafenib). Experienced disease progression on prior anti-BRAF/MEK combination therapy [e.g. Cotellic (cobimetinib) with Zelboraf (vemurafenib) or Braftovi (encorafenib) with Mektovi (binimetinib)].
Required Medical Information	Pending CMS Review
Age Restriction	Low-grade Glioma only: The member is a pediatric age 1 year of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration

Other Criteria	
Part B Prerequisite	0

TAGRISSO - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members who have disease progression on Tagrisso (osimertinib) (applicable to adjuvant therapy in NSCLC and NSCLC locally advanced, unresectable stage III). Total treatment exceeds three years (applicable to adjuvant therapy in NSCLC).
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

TALZENNA - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members have experienced disease progression while on or following PARP inhibitor therapy (eg, olaparib).
Required Medical Information	Breast Cancer. Member has a diagnosis of locally advanced or metastatic, HER-2 negative breast cancer AND Member has documented deleterious germline or suspected germline BRCA mutated disease AND if member has hormone receptor positive disease then is endocrine refractory AND Talzenna (talazoparib) will be used as monotherapy. Metastatic Castration-Resistant Prostate Cancer (mCRPC). Member has a diagnosis of Metastatic Castration-Resistant Prostate Cancer (mCRPC) AND Member has a documented (HRR) gene-mutated disease AND Member will use Talzenna (talazoparib) in combination with androgen deprivation therapy (e.g. previous bilateral orchiectomy or luteinizing hormone-releasing hormone (LHRH) analog [i.e., LHRH agonist/ antagonist]) AND Talzenna (talazoparib) is given in combination with Xtandi (enzalutamide).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

TARPEYO - CAPSULE,DELAYED RELEASE (ENTERIC COATED)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Primary immunoglobulin A nephropathy (IgAN): Member has a diagnosis of primary immunoglobulin A nephropathy (IgAN) confirmed by biopsy AND Tarpeyo is being used to reduce the loss of kidney function in those who are at risk for disease progression
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

TAZVERIK - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member experiences disease progression on Tazverik
Required Medical Information	Epithelioid Sarcoma: The member has a diagnosis of metastatic or locally advanced epithelioid sarcoma not eligible for complete resection AND Tazverik will be given as monotherapy. Follicular lymphoma: The member has a diagnosis of relapsed/refractory follicular lymphoma AND one of the following applies: The member has a documented EZH2 mutation by an FDA approved test and the member has received at least two prior therapies and the member will be using Tazverik (tazemetostat) as monotherapy OR The member has no satisfactory alternative treatment options and The member will be using Tazverik (tazemetostat) as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

TECENTRIQ - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	All indications except Bladder Cancer: Disease progression while on or following anti-PD-1/PD-L1 therapy (e.g. Opdivo [nivolumab], Keytruda [pembrolizumab], Tecentriq [atezolizumab], Bavencio [avelumab]). Adjuvant setting for Non-Small Cell Lung Cancer: member is taking Tecentriq (atezolizumab) total treatment for more than one year.
Required Medical Information	Non-Small Cell Lung Cancer (advanced or metastatic frontline): The member has a diagnosis of advanced or metastatic NSCLC AND member has disease with no EGFR or ALK genomic tumor aberrations and one of the following scenarios is applied: The member has non-squamous cell histology AND Tecentriq will be given as a component of one of the two combo regimens: in combination with carboplatin and paclitaxel and Bevacizumab Product as first line therapy followed by maintenance therapy with combination an atezolizumab product and Bevacizumab Product OR in combo with Abraxane (nabpaclitaxel) and carboplatin as first line therapy. OR Disease has high PD-L1 expression [PD-L1 stained greater than or equal to 50% of tumor cells OR PD-L1 stained tumor-infiltrating immune cells covering greater than or equal to 10% of the tumor area] AND PD-L1 tumor expression is determined by an FDA-approved test AND will be given as first-line therapy AND The member will be using as monotherapy. Non-Small Cell Lung Cancer (metastatic - subsequent): The member has a diagnosis of metastatic non-small cell lung carcinoma AND The member has experienced disease progression on or after chemotherapy and EGFR inhibitor or ALK inhibitor (post confirmed EGFR or ALK genomic tumor aberration positivity) AND atezolizumab product will be used as monotherapy. Alveolar Soft Part Sarcoma (ASPS): The member has a diagnosis of unresectable or metastatic alveolar soft part sarcoma AND The member will be using an atezolizumab product as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 months duration

TECENTRIQ - VIAL (ML)

Other Criteria	Small Cell Lung Cancer: The member has a diagnosis of extensive-stage small cell lung cancer AND atezolizumab product will be given in combination with etoposide and carboplatin as first line therapy followed by maintenance therapy with atezolizumab product, either as monotherapy or in combination with Zepzelca (lurbinectedin). Heptatocellular Carcinoma: The member has a diagnosis of unresectable or metastatic hepatocellular carcinoma AND atezolizumab product will be used as first line therapy in combination with bevacizumab product. Melanoma: The member has a diagnosis of unresectable or metastatic melanoma AND The member has a documented BRAF V600 activating mutation AND the member will use an atezolizumab product in combination with Cotellic (cobimetinib) and Zelboraf (vemurafenib). Non-Small Cell Lung Cancer (Adjuvant): The member must have a diagnosis of Stage II to IIIA non-small cell lung cancer AND The disease has expression of PD-L1 on greater than or equal to 1% of tumor cells as determined by an FDA-approved test AND The member is post complete surgical resection and adjuvant platinum-based chemotherapy AND The member will be using an atezolizumab product as monotherapy in the adjuvant setting. Bladder cancer: The member has a diagnosis of muscle invasive bladder cancer AND After cycsteomy, the member has circulating tumor DNA molecular residual disease as determined by an FDA approved test AND An atezolizumab product will be used as a single agent as part of adjuvant treatment.
Part B Prerequisite	0

TECENTRIQ HYBREZA - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	All indications except Bladder Cancer: Disease progression while on or following anti-PD-1/PD-L1 therapy (e.g. Opdivo [nivolumab], Keytruda [pembrolizumab], Tecentriq [atezolizumab], Bavencio [avelumab]). Adjuvant setting for Non-Small Cell Lung Cancer: member is taking Tecentriq (atezolizumab) total treatment for more than one year.
Required Medical Information	Non-Small Cell Lung Cancer (advanced or metastatic frontline): The member has a diagnosis of advanced or metastatic NSCLC AND member has disease with no EGFR or ALK genomic tumor aberrations and one of the following scenarios is applied: The member has non-squamous cell histology AND Tecentriq will be given as a component of one of the two combo regimens: in combination with carboplatin and paclitaxel and Bevacizumab Product as first line therapy followed by maintenance therapy with combination an atezolizumab product and Bevacizumab Product OR in combo with Abraxane (nabpaclitaxel) and carboplatin as first line therapy. OR Disease has high PD-L1 expression [PD-L1 stained greater than or equal to 50% of tumor cells OR PD-L1 stained tumor-infiltrating immune cells covering greater than or equal to 10% of the tumor area] AND PD-L1 tumor expression is determined by an FDA-approved test AND will be given as first-line therapy AND The member will be using as monotherapy. Non-Small Cell Lung Cancer (metastatic - subsequent): The member has a diagnosis of metastatic non-small cell lung carcinoma AND The member has experienced disease progression on or after chemotherapy and EGFR inhibitor or ALK inhibitor (post confirmed EGFR or ALK genomic tumor aberration positivity) AND atezolizumab product will be used as monotherapy. Alveolar Soft Part Sarcoma (ASPS): The member has a diagnosis of unresectable or metastatic alveolar soft part sarcoma AND The member will be using an atezolizumab product as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 months duration

Other Criteria	Small Cell Lung Cancer: The member has a diagnosis of extensive-stage small cell lung cancer AND atezolizumab product will be given in combination with etoposide and carboplatin as first line therapy followed by maintenance therapy with atezolizumab product, either as monotherapy or in combination with Zepzelca (lurbinectedin). Heptatocellular Carcinoma: The member has a diagnosis of unresectable or metastatic hepatocellular carcinoma AND atezolizumab product will be used as first line therapy in combination with bevacizumab product. Melanoma: The member has a diagnosis of unresectable or metastatic melanoma AND The member has a documented BRAF V600 activating mutation AND the member will use an atezolizumab product in combination with Cotellic (cobimetinib) and Zelboraf (vemurafenib). Non-Small Cell Lung Cancer (Adjuvant): The member must have a diagnosis of Stage II to IIIA non-small cell lung cancer AND The disease has expression of PD-L1 on greater than or equal to 1% of tumor cells as determined by an FDA-approved test AND The member is post complete surgical resection and adjuvant platinum-based chemotherapy AND The member will be using an atezolizumab product as monotherapy in the adjuvant setting. Bladder cancer: The member has a diagnosis of muscle invasive bladder cancer AND After cycsteomy, the member has circulating tumor DNA molecular residual disease as determined by an FDA approved test AND An atezolizumab product will be used as a single agent as part of adjuvant treatment.
Part B Prerequisite	0

TEPMETKO - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member experiences disease progression on MET inhibitor (e.g., Tabrecta, Tepmetko).
Required Medical Information	Non-small cell lung cancer (NSCLC). The member has a diagnosis of metastatic non-small cell lung cancer (NSCLC) AND The disease is documented MET exon 14 skipping positive.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

teriflunomide - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	The member has a diagnosis of a relapsing form of multiple sclerosis, to include relapsing-remitting or active secondary progressive disease, OR the member has a diagnosis of clinically isolated syndrome (CIS).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

teriparatide - PEN INJECTOR (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

testosterone - GEL IN METERED-DOSE PUMP

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Member has one of the following diagnoses: Metastatic Breast Cancer (testosterone enanthate only) OR Delay Puberty in Males (testosterone enanthate only) OR Primary hypogonadism: testicular failure due to cryptorchidism, bilateral torsion, orchitis, vanishing testis syndrome, orchiectomy, Klinefelter's syndrome, chemotherapy, or toxic damage from alcohol or heavy metals OR Hypogonadotropic hypogonadism: idiopathic gonadotropin or luteinizing hormone-releasing hormone (LHRH) deficiency or pituitary-hypothalamic injury from tumors, trauma, or radiation AND Member has had one of the following: Documentation of two morning serum testosterone levels (total or free) that are less than the reference range for the lab, taken at separate times, prior to treatment OR Documentation of a serum testosterone level (total or free) that is less than or within the reference range for the lab, when already on treatment.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

testosterone cypionate - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Member has one of the following diagnoses: Metastatic Breast Cancer (testosterone enanthate only) OR Delay Puberty in Males (testosterone enanthate only) OR Primary hypogonadism: testicular failure due to cryptorchidism, bilateral torsion, orchitis, vanishing testis syndrome, orchiectomy, Klinefelter's syndrome, chemotherapy, or toxic damage from alcohol or heavy metals OR Hypogonadotropic hypogonadism: idiopathic gonadotropin or luteinizing hormone-releasing hormone (LHRH) deficiency or pituitary-hypothalamic injury from tumors, trauma, or radiation AND Member has had one of the following: Documentation of two morning serum testosterone levels (total or free) that are less than the reference range for the lab, taken at separate times, prior to treatment OR Documentation of a serum testosterone level (total or free) that is less than or within the reference range for the lab, when already on treatment.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

testosterone enanthate - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Member has one of the following diagnoses: Metastatic Breast Cancer (testosterone enanthate only) OR Delay Puberty in Males (testosterone enanthate only) OR Primary hypogonadism: testicular failure due to cryptorchidism, bilateral torsion, orchitis, vanishing testis syndrome, orchiectomy, Klinefelter's syndrome, chemotherapy, or toxic damage from alcohol or heavy metals OR Hypogonadotropic hypogonadism: idiopathic gonadotropin or luteinizing hormone-releasing hormone (LHRH) deficiency or pituitary-hypothalamic injury from tumors, trauma, or radiation AND Member has had one of the following: Documentation of two morning serum testosterone levels (total or free) that are less than the reference range for the lab, taken at separate times, prior to treatment OR Documentation of a serum testosterone level (total or free) that is less than or within the reference range for the lab, when already on treatment.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

tetrabenazine - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Diagnosis of chorea associated with Huntington's disease.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

TEVIMBRA - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Esophageal squamous cell carcinoma: The member has a diagnosis of unresectable or metastatic esophageal squamous cell carcinoma AND One of the following: Tevimbra (tislelizumab-jsgr) will be used as monotherapy AND the member has received prior systemic chemotherapy that did not include a PD-L1 inhibitor OR The member will be using Tevimbra (tislelizumab-jsgr) as first line therapy AND the tumor expresses PD-L1 (Combined Positive Score (CPS) greater than or equal to 1) AND tevimbra (tislelizumab-jsgr) will be used in combination with platinum-containing chemotherapy, followed by Tevimbra (tislelizumab-jsgr) monotherapy. Gastric or gastroesophageal junction adenocarcinoma: The member has a diagnosis of unresectable or metastatic gastric or gastroesophageal junction adenocarcinoma AND The tumor is HER2-negative and PD-L1 positive (CPS greater than or equal to 1) AND Tevimbra (tislelizumab-jsgr) will be used as first line therapy AND Tevimbra (tislelizumab-jsgr) will be used in combination with platinum and fluoropyrimidine-based chemotherapy, followed by Tevimbra (tislelizumab-jsgr) monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 months duration
Other Criteria	
Part B Prerequisite	0

THALOMID - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on thalidomide.
Required Medical Information	Erythema Nodosum Leprosum (ENL). The member is currently having acute cutaneous manifestations of moderate to severe erythema nodosum leprosum (ENL) OR Thalomid (thalidomide) is prescribed for maintenance therapy for prevention and suppression of the cutaneous manifestations of (ENL) recurrence. Multiple Myeloma. The member has a diagnosis of Multiple Myeloma.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

TIBSOVO - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member has experienced disease progression while on or following Tibsovo (ivosidenib)
Required Medical Information	Acute Myeloid Leukemia- Relapsed/Refractory: The member has a diagnosis of acute myeloid leukemia (AML) AND The member has relapsed or refractory disease AND The member has a documented IDH1 mutation AND one of the following applies: The member will be using Tibsovo (ivosedinib) as monotherapy OR the member will be using Tibsovo as a component of repeating the initial successful induction regimen, if late relapse (relapse occurring later than 12 months). Acute Myeloid Leukemia Newly diagnosed: The member has a diagnosis of acute myeloid leukemia (AML) AND the member has newly diagnosed disease AND one of the following applies: the member is not a candidate for intensive induction therapy due to comorbidities OR the member declines intensive induction therapy. The member has a documented IDH1 mutation AND the member will be using Tibsovo as monotherapy or in combination with azactidine. Cholangiocarcinoma: The member has locally advanced or metastatic cholangiocarcinoma AND the disease has documented isocitrate dehydrogenate-1 (IDH1) mutation AND Tibsovo (ivosedinib) will be a subsequent therapy and used as monotherapy. Myelodysplastic syndromes (MDS): The member has a diagnosis of myelodysplastic syndromes AND The member has relapsed or refractory disease AND The member has a documented IDH1 mutation AND The member will be using Tibsovo (ivosidenib) as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

TOBI PODHALER - CAPSULE, WITH INHALATION DEVICE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Patients colonized with Burkholderia cepacia.
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	1

tolvaptan (polycys kidney dis) - TABLET, SEQUENTIAL

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Autosomal Dominant Polycystic Kidney Disease. Has a diagnosis of autosomal dominant polycystic kidney disease (ADPKD) AND Provider attests member is at risk of rapidly progressive disease AND Must have CKD stage 2 - 4 AND Medication is being used to slow kidney function decline.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

topiramate - SOLUTION, ORAL

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Migraine Prophylaxis: Member is using for prophylaxis of migraine headache AND has had previous treatment with or intolerance to immediate release topiramate tablet or capsule AND has had previous treatment, intolerance, or contraindication with one of the following: a seratonin and norepinephrine reuptake inhibitor or a tricyclic antidepressant. Epilepsy Adjunctive Therapy: Member has a diagnosis of partial-onset (focal) seizures, primary generalized tonic-clonic seizures, or seizures associated with Lennox-Gastaut syndrome (LGS) AND Member will use Eprontia (topiramate) in combination with at least one other drug for controlling seizures AND Member has tried or cannot use immediate-release topiramate tablet or capsule. Epilepsy Monotherapy: Member has a diagnosis of partial-onset (focal) seizures or primary generalized tonic-clonic seizures AND Member has tried or cannot use immediate release topiramate tablet or capsule.
Age Restriction	Migraine Prophylaxis: Member must be 12 years of age or older. Epilepsy indications: Member must be 2 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

torpenz - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member has experienced disease progression while on everolimus.
Required Medical Information	Subependymal Giant Cell Astrocytoma (SEGA) with TSC [Adult and Pediatrics]: Member must have an intolerance or contraindication to generic everolimus AND The member has a diagnosis of SEGA associated with tuberous sclerosis complex (TSC) AND The member requires therapeutic intervention but is not a candidate for curative surgical resection. Angiomyolipoma and Tuberous Sclerosis Complex (TSC): Member must have an intolerance or contraindication to generic everolimus AND The member has a diagnosis of renal angiomyolipoma and tuberous sclerosis complex (TSC) AND Immediate surgery is not required. Metastatic Breast Cancer: Member must have an intolerance or contraindication to generic everolimus AND The member has a diagnosis of hormone receptor (HR) positive and human epidermal growth factor receptor 2 (HER2) negative metastatic disease AND The member has been treated with endocrine therapy (e.g., letrozole, anastrozole) AND The member will use everolimus in combination with exemestane or fulvestrant.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

tramadol - TABLET, EXTENDED RELEASE 24 HR

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Severe and Persistent pain: Has persistent, non-acute pain that requires around-the-clock treatment AND Has pain severe enough to require an opioid pain medication AND previous trials of alternative treatments (e.g., nonopioid analgesics) have been inadequate AND Has tried or cannot take tramadol 50mg tablet.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

TRAZIMERA - VIAL (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	For Herceptin (trastuzumab), Kanjinti (trastuzumab-anns) , Herzuma (trastuzumab-pkrb), Hercessi (trastuzumab-strf) or Ontruzant (trastuzumab-dkst) requests: member must have an intolerance or contraindication Trazimera (trastuzumab-qyyp) OR Ogivri (trastuzumab-dkst) and meets below criteria: Breast Cancer: The member has a diagnosis of breast cancer and HER2 (human epidermal growth factor receptor2) positive disease (e.g., defined as IHC 3+ or IHC2+/ISH positive). Gastric Cancer: The member has a diagnosis of advanced, gastric cancer or gastroesophageal adenocarcinoma and HER2 positive disease (e.g., defined as IHC 3+ or IHC2+/ISH positive) AND trastuzumab is being used in combination with cisplatin and fluorouracil or capecitabine.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	All Medically accepted uses not listed above: The member is using a trastuzumab product for a medically accepted indication as defined by Medicare rules. For Herceptin (trastuzumab), Kanjinti (trastuzumab-anns), Herzuma (trastuzumab-pkrb), Hercessi (trastuzumab-strf), or Ontruzant (trastuzumab-dttb) requests: member must have an intolerance or contraindication to Trazimera (trastuzumab-qyyp) or Ogivri (trastuzumab-dkst).
Part B Prerequisite	0

TRELSTAR - VIAL (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

TREMFYA - SYRINGE (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Plaque Psoriasis: The member has a diagnosis of moderate to severe plaque psoriasis AND the member must weigh at least 40kg AND The member has tried at least one conventional oral systemic treatment (e.g. methotrexate, cyclosporine), or cannot use any conventional oral systemic treatments. Psoriatic Arthritis: The member has a diagnosis of active psoriatic arthritis AND the member must weigh at least 40kg. Ulcerative Colitis: The member has a diagnosis of moderate to severely active ulcerative colitis. Crohn's Disease: The member has a diagnosis of moderately to severely active Crohn's Disease.
Age Restriction	Plaque Psoriasis, Psoriatic Arthritis: The member must be 6 years of age or older. All other indications: The member must be 18 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

TREMFYA ONE-PRESS - AUTO-INJECTOR (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Plaque Psoriasis: The member has a diagnosis of moderate to severe plaque psoriasis AND the member must weigh at least 40kg AND The member has tried at least one conventional oral systemic treatment (e.g. methotrexate, cyclosporine), or cannot use any conventional oral systemic treatments. Psoriatic Arthritis: The member has a diagnosis of active psoriatic arthritis AND the member must weigh at least 40kg. Ulcerative Colitis: The member has a diagnosis of moderate to severely active ulcerative colitis. Crohn's Disease: The member has a diagnosis of moderately to severely active Crohn's Disease.
Age Restriction	Plaque Psoriasis, Psoriatic Arthritis: The member must be 6 years of age or older. All other indications: The member must be 18 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

TREMFYA PEN - PEN INJECTOR (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Plaque Psoriasis: The member has a diagnosis of moderate to severe plaque psoriasis AND the member must weigh at least 40kg AND The member has tried at least one conventional oral systemic treatment (e.g. methotrexate, cyclosporine), or cannot use any conventional oral systemic treatments. Psoriatic Arthritis: The member has a diagnosis of active psoriatic arthritis AND the member must weigh at least 40kg. Ulcerative Colitis: The member has a diagnosis of moderate to severely active ulcerative colitis. Crohn's Disease: The member has a diagnosis of moderately to severely active Crohn's Disease.
Age Restriction	Plaque Psoriasis, Psoriatic Arthritis: The member must be 6 years of age or older. All other indications: The member must be 18 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

TREMFYA PEN INDUCTION PK(2PEN) - PEN INJECTOR (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Plaque Psoriasis: The member has a diagnosis of moderate to severe plaque psoriasis AND the member must weigh at least 40kg AND The member has tried at least one conventional oral systemic treatment (e.g. methotrexate, cyclosporine), or cannot use any conventional oral systemic treatments. Psoriatic Arthritis: The member has a diagnosis of active psoriatic arthritis AND the member must weigh at least 40kg. Ulcerative Colitis: The member has a diagnosis of moderate to severely active ulcerative colitis. Crohn's Disease: The member has a diagnosis of moderately to severely active Crohn's Disease.
Age Restriction	Plaque Psoriasis, Psoriatic Arthritis: The member must be 6 years of age or older. All other indications: The member must be 18 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

tretinoin - GEL (GRAM)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Approval will be given to all members using this agent for medically necessary, FDA approved, or compendia supported, non-cosmetic indications including but not limited to the following: Acne: the member has a diagnosis of acne vulgaris, Actinic Keratosis: the member has a diagnosis of actinic keratosis.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

TRIKAFTA - TABLET, SEQUENTIAL

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Cystic Fibrosis: The member must meet ALL of the following criteria: Diagnosis of Cystic Fibrosis AND Prior lab testing with documentation of a mutation in the CFTR gene that is responsive to therapy based on clinical literature and/or in vitro assay data.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

TRULICITY - PEN INJECTOR (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member is using for weight loss.
Required Medical Information	Diabetes Mellitus. The member has a diagnosis of type 2 diabetes mellitus.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

TRUQAP - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member has progression while on Truqap (capivasertib).
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

TUKYSA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member experiences disease progression on Tukysa (tucatinib)
Required Medical Information	Breast Cancer. The member has metastatic or advanced unresectable HER 2 positive breast cancer (inclusive of brain metastases) and all of the following apply: Member has received one or more prior anti-HER2 based regimen in the metastatic setting AND Tukysa is given in combination with trastuzumab product and capecitabine as subsequent therapy. Colorectal cancer. The member has a diagnosis of RAS wild-type HER2-positive unresectable or metastatic colorectal cancer AND The member has progressed following treatment with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy AND Tuksya will be given in combination with trastuzumab product.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

TURALIO - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	NA
Exclusion Criteria	NA
Required Medical Information	Symptomatic Tenosynovial Giant Cell Tumor: The member has symptomatic tenosynovial giant cell tumor (TGCT) and the disease is associated with severe morbidity or functional limitations and the disease is not amenable to improvement with surgery and Turalio (pexidartinib) will be used as monotherapy.
Age Restriction	NA
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 months duration
Other Criteria	NA
Part B Prerequisite	0

TYENNE - SYRINGE (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	Member must be 18 years and older. For the treatment of PJIA and SJIA the member must be at least 2 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

TYENNE AUTOINJECTOR - PEN INJECTOR (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	Member must be 18 years and older. For the treatment of PJIA and SJIA the member must be at least 2 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

TYMLOS - PEN INJECTOR (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Osteoporosis in Postmenopausal Women: The member must have a diagnosis of osteoporosis AND the member must be postmenopausal AND The member must meet one of the following criteria: The member must be at a high risk of fracture defined as a history of osteoporotic fracture or The member has multiple risk factors for fracture (e.g. inflammatory joint disorders, diabetes, hyperthyroidism, liver disease) OR The member has failed or is intolerant to at least ONE other available osteoporosis therapy (e.g. bisphosphates, RANK-ligand inhibitor, parathyroid hormone analog, sclerostin inhibitor, etc.) OR The member must be at VERY HIGH RISK for fracture as defined as: recent fracture (e.g. within the past 12 months) OR fracture while on approved therapy for treatment of osteoporosis OR fracture while on drugs causing skeletal harm (e.g. systemic corticosteroids) OR experienced multiple fractures OR has a very low T-score (e.g. less than -3.0) OR is at high risk for falls or has a history of falls OR has a very high fracture probability per FRAX (e.g. greater than 30% major osteoporotic fracture, greater than 4.5% hip fracture). Osteoporosis in Men: The member must be using for osteoporosis in men AND The member is at high risk of fracture as defined as: a history of osteoporotic fracture OR multiple risk factors for fracture OR The member has new fractures or significant loss of bone mineral density despite previous treatment, contraindication or intolerance with an oral or intravenous bisphosphonate (e.g. alendronate).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

UBRELVY - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Acute Migraine: The member will be using Ubrelvy (ubrogepant) for the acute treatment of migraines AND The member has tried or cannot use ONE of the following: naratriptan tablet, rizatriptan tablet, sumatriptan tablet, eletriptan tablet.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

UDENYCA - SYRINGE (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Febrile Neutropenia Prophylaxis: Requested product will be administered more frequently than once every two weeks.
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 months duration
Other Criteria	Hematopoietic Subsyndrome of Acute Radiation Syndrome: The member has been acutely exposed to myelosuppressive doses of nontherapeutic radiation. Stem Cell Mobilization for Autologous Donors: Requested product is being used in combination with plerixafor as part of a hematopoietic stem cell mobilization regimen.
Part B Prerequisite	0

UDENYCA AUTOINJECTOR - AUTO-INJECTOR (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Febrile Neutropenia Prophylaxis: Requested product will be administered more frequently than once every two weeks.
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 months duration
Other Criteria	Hematopoietic Subsyndrome of Acute Radiation Syndrome: The member has been acutely exposed to myelosuppressive doses of nontherapeutic radiation. Stem Cell Mobilization for Autologous Donors: Requested product is being used in combination with plerixafor as part of a hematopoietic stem cell mobilization regimen.
Part B Prerequisite	0

UDENYCA ONBODY - SYRINGE, WITH WEARABLE INJECTOR

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Febrile Neutropenia Prophylaxis: Requested product will be administered more frequently than once every two weeks.
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 months duration
Other Criteria	Hematopoietic Subsyndrome of Acute Radiation Syndrome: The member has been acutely exposed to myelosuppressive doses of nontherapeutic radiation. Stem Cell Mobilization for Autologous Donors: Requested product is being used in combination with plerixafor as part of a hematopoietic stem cell mobilization regimen.
Part B Prerequisite	0

UNITUXIN - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members receiving Unituxin (dinutuximab) as monotherapy. Members that have experienced disease progression while on Unituxin (dinutuximab).
Required Medical Information	High-risk neuroblastoma:The member has a diagnosis of high-risk neuroblastoma ANDUnituxin (dinutuximab) will be used in combination with isotretinoin AND Unituxin (dinutuximab) will be used in alternating cycles of Leukine (sargramostim) and Proleukin (aldesleukin) AND The member has achieved at least a partial response to the following: Induction combination chemotherapy AND Maximum feasible surgical resection The member has had the previous procedure/therapy: Myeloablative consolidation chemotherapy followed by autologous stem cell transplantation AND Radiation therapy to residual soft tissue disease.
Age Restriction	Memebr must be 18 years of age or younger.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

UNLOXCYT - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression on or after prior PD-1/PD-L1 inhibitor (e.g., Keytruda).
Required Medical Information	Cutaneous squamous cell carcinoma: The member has a diagnosis of cutaneous squamous cell carcinoma AND The disease is locally advanced or metastatic AND The member is not a candidate for curative surgery or curative radiation Unloxcyt (cosibelimab-ipdl) will be used as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 months duration
Other Criteria	
Part B Prerequisite	0

UPTRAVI - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

VALCHLOR - GEL (GRAM)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on Valchlor (mechlorethamine).
Required Medical Information	Cutaneous T-Cell Lymphoma: The member has a diagnosis of fungoides-type cutaneous T-cell lymphoma AND The member has had prior therapy with skin-directed therapy or using as primary treatment.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

valrubicin - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has an active urinary tract infection (UTI). The member has perforated bladder or compromised bladder mucosa. The member has small bladder capacity (unable to tolerate a 75 mL instillation).
Required Medical Information	Bladder Cancer: The member has recurrent or persistent carcinoma in situ of the urinary bladder(Cis). The member has experienced disease progression, intolerance or has a contraindication to BCG therapy. The member is not a candidate for immediate cystectomy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

VANFLYTA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on or following Vanflyta (quizartinib).
Required Medical Information	Acute Myeloid Leukemia (newly diagnosed). The member has newly diagnosed acute myeloid leukemia (AML) AND The member has documented FLT3 internal tandem duplication (ITD)-positive disease AND The member will be using Vanflyta (quizartinib) in combination with standard cytarabine and anthracycline induction and cytarabine consolidation chemotherapy OR The member will be using Vanflyta (quizartinib) as maintenance monotherapy following consolidation with systemic chemotherapy (excludes maintenance monotherapy following allogeneic hemopoietic stem cell transplantation).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

VECTIBIX - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members with colorectal cancer (CRC) in which RAS mutation status is unknown. Member has disease progression on Vectibix or Erbutux.
Required Medical Information	Colorectal Cancer [Wild-Type RAS; Wild-Type KRAS/NRAS]: The member has a diagnosis of advanced or metastatic colorectal cancer (mCRC) AND The member has colorectal cancer that is documented as wild-type, non-mutated KRAS and NRAS [wild-type RAS] AND one of the criteria following applies: Vectibix (panitumumab) will be used as monotherapy in members who had disease progression on or following fluoropyrimidine (e.g., capecitabine, fluorouracil), oxaliplatin, and irinotecan-containing chemotherapy regimens . OR Vectibix (panitumumab) will be used in combination with CAPEOX, FOLFOX or FOLFIRI OR Vectibix (panitumumab) will be used concurrently with irinotecan-based therapy. Colorectal Cancer [KRAS G12C-mutated]: The member has a diagnosis of metastatic colorectal cancer (mCRC) AND The colorectal cancer is documented as KRAS G12C-mutated, as determined by an FDA-approved test AND The member has received prior systemic treatment with chemotherapy [fluoropyrimidine-based chemotherapy (e.g., 5-fluorouracil, capecitabine), oxaliplatin, irinotecan] AND Vectibix (panitumumab) will be given in combination with Lumakras (sotorasib).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

VELSIPITY - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Ulcerative Colitis: The member has a diagnosis of moderately to severely active ulcerative colitis AND the member has tried or cannot use two of the following: Preferred adalimumab product [i.e. adalimumab-adbm, adalimumab-adaz], preferred ustekinumab product (i.e., Otulfi, Yesintek), Skyrizi, Rinvoq, or Tremfya.
Age Restriction	18 years of age or older
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

VENCLEXTA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression while on Venclexta (venetoclax).
Required Medical Information	Chronic Lymphocytic Leukemia (CLL):The member has a diagnosis of Chronic Lymphocytic Leukemia (CLL) and one of the following applies: The member has received at least one prior therapy OR The request is for first-line therapy AND the member is using Venclexta (venetoclax) as part of a supported combination therapy. Mantle Cell Lymphoma: The member has a diagnosis of MCL AND one of the following applies: the member is using Venclexta (venetoclax) as second or subsequent line therapy AND the member is using Venclexta (venetoclax) as monotherapy or in combination with a rituximab product or in combination with Imbruvica (ibrutinib) OR The member has untreated, TP53 mutated disease AND the member is using Venclexta (venetoclax) in combination with Brukinsa (zanubrutinib) and Gazyva (obinutuzumab). Acute Myeloid Leukemia: The member has a diagnosis of acute myeloid leukemia (AML) AND one of the following applies: the disease is newly diagnosed AND the member is 75 years or older AND the member will be using Venclexta (venetoclax) in combination with azacitidine, or decitabine, or low-dose cytarabine OR the disease is newly diagnosed AND the member has comorbidities that preclude the use of intensive induction chemotherapy (e.g. baseline Eastern Cooperative Oncology Group (ECOG) performance status of 2-3, severe cardiac or pulmonary comorbidity, moderate hepatic impairment, or creatinine clearance less than 45 ml/min) AND the member will be using Venclexcta (ventoclax) in combination with azacitidine, or decitabine, or low-dose cytarabine OR The member has relapsed/refractory disease AND Venclexta (venetoclax) will be used as part of supported combination therapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

VENCLEXTA STARTING PACK - TABLET, DOSE PACK

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression while on Venclexta (venetoclax).
Required Medical Information	Chronic Lymphocytic Leukemia (CLL):The member has a diagnosis of Chronic Lymphocytic Leukemia (CLL) and one of the following applies: The member has received at least one prior therapy OR The request is for first-line therapy AND the member is using Venclexta (venetoclax) as part of a supported combination therapy. Mantle Cell Lymphoma: The member has a diagnosis of MCL AND one of the following applies: the member is using Venclexta (venetoclax) as second or subsequent line therapy AND the member is using Venclexta (venetoclax) as monotherapy or in combination with a rituximab product or in combination with Imbruvica (ibrutinib) OR The member has untreated, TP53 mutated disease AND the member is using Venclexta (venetoclax) in combination with Brukinsa (zanubrutinib) and Gazyva (obinutuzumab). Acute Myeloid Leukemia: The member has a diagnosis of acute myeloid leukemia (AML) AND one of the following applies: the disease is newly diagnosed AND the member is 75 years or older AND the member will be using Venclexta (venetoclax) in combination with azacitidine, or decitabine, or low-dose cytarabine OR the disease is newly diagnosed AND the member has comorbidities that preclude the use of intensive induction chemotherapy (e.g. baseline Eastern Cooperative Oncology Group (ECOG) performance status of 2-3, severe cardiac or pulmonary comorbidity, moderate hepatic impairment, or creatinine clearance less than 45 ml/min) AND the member will be using Venclexcta (ventoclax) in combination with azacitidine, or decitabine, or low-dose cytarabine OR The member has relapsed/refractory disease AND Venclexta (venetoclax) will be used as part of supported combination therapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

VEPPANU - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Estrogen Receptor (ER) positive Breast Cancer: The member has a diagnosis of ER-positive, HER2-negative advanced or metastatic breast cancer AND The breast cancer has a documented ESR1-mutation as determined by FDA approved test AND The member has experienced progressive disease following at least one prior line of endocrine therapy (e.g., aromatase inhibitor, CDK 4/6 inhibitor).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

VERQUVO - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

VERSACLOZ - SUSPENSION, ORAL (FINAL DOSE FORM)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	The member must be using clozapine ODT or Versacloz (clozapine oral solution) for treatment-resistant schizophrenia. The member has tried or cannot use generic clozapine tablets.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

VERZENIO - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Metastatic Breast cancer- initial endocrine based therapy. The member has a diagnosis of advanced or metastatic hormone receptor (HR)-positive and human epidermal growth factor receptor 2 (Her2neu)-negative breast cancer AND Verzenio (abemaciclib) is given in combination with an aromatase inhibitor (e.g., letrozole) as first line endocrine based therapy. Metastatic breast cancer combination therapy with Faslodex (fulvestrant). The member has diagnosis of advanced or metastatic hormone receptor (HR) positive human epidermal growth factor receptor 2 (HER2) negative breast cancer AND The member has experienced disease progression on endocrine therapy (e.g., anastrazole) AND Verzenio (abemaciclib) is given in combination with Faslodex (fulvestrant). Metastatic breast cancer monotherapy: The member has diagnosis of advanced or metastatic HR positive, HER2 negative breast cancer AND the member has experienced disease progression on endocrine therapy (e.g., anastrazole) and chemotherapy in the metastatic setting AND Verzenio (abemaciclib) is being used as monotherapy. Early Breast cancer - combination therapy: The member has a diagnosis of HR positive, HER2 negative, node positive, early breast cancer at high risk of recurrence AND Verzenio (abemaciclib) is given in combination with tamoxifen or aromatase inhibitor.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

vigabatrin - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Refractory Complex Partial Seizure: Member has a diagnosis of refractory complex partial seizures AND Member has been unable to achieve seizure control with at least TWO other drugs for controlling complex partial seizures (e.g. levetiracetam, topiramate, carbamazepine, gabapentin, divalproex, or lamotrigine) AND vigabatrin will be used in combination with at least one other drug for controlling complex partial seizures. Infantile Spasms: Documented diagnosis of infantile spasms.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

vigadrone - POWDER IN PACKET (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Refractory Complex Partial Seizure: Member has a diagnosis of refractory complex partial seizures AND Member has been unable to achieve seizure control with at least TWO other drugs for controlling complex partial seizures (e.g. levetiracetam, topiramate, carbamazepine, gabapentin, divalproex, or lamotrigine) AND vigabatrin will be used in combination with at least one other drug for controlling complex partial seizures. Infantile Spasms: Documented diagnosis of infantile spasms.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

VIGAFYDE - SOLUTION, ORAL

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Infantile Spasms: The member has a documented diagnosis of infantile spasms.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

vigpoder - POWDER IN PACKET (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Refractory Complex Partial Seizure: Member has a diagnosis of refractory complex partial seizures AND Member has been unable to achieve seizure control with at least TWO other drugs for controlling complex partial seizures (e.g. levetiracetam, topiramate, carbamazepine, gabapentin, divalproex, or lamotrigine) AND vigabatrin will be used in combination with at least one other drug for controlling complex partial seizures. Infantile Spasms: Documented diagnosis of infantile spasms.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

vilazodone - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Major depressive disorder: The member has a clinical diagnosis of major depressive disorder (MDD) AND The member has tried or cannot use a selective serotonin reuptake inhibitor (SSRI) (e.g., fluoxetine, citalopram, or sertraline) AND a generic bupropion product (75mg/100mg IR, 100mg/150mg/200mg SR, or 150mg/300mg XL) or mirtazapine.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

VITRAKVI - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

VIZIMPRO - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Non-small cell lung cancer (NSCLC). The member has a diagnosis of metastatic non-small cell lung cancer (NSCLC) AND the following criteria applies: The member has a documented epidermal growth factor receptor (EGFR) exon 19 deletion or exon 21 (L858R) substitution mutation AND The member is using Vizimpro (dacomitinib) as a single agent for first line therapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Six month duration
Other Criteria	
Part B Prerequisite	0

VONJO - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on Vonjo (pacritinib).
Required Medical Information	Myelofibrosis: The member has a diagnosis of primary myelofibrosis or secondary myelofibrosis (i.e. post-polycythemia vera or post-essential thrombocythemia) AND The member has one of the following risk categories, as defined by accepted risk stratification tools for myelofibrosis (e.g. International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or DIPSS-PLUS): Intermediate-risk disease OR High-risk disease AND The member will be using Vonjo (pacritinib) as monotherapy AND The member has a platelet count below 50 X 109/L.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

VORANIGO - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members have experienced disease progression while on Voranigo (vorasidenib).
Required Medical Information	Gliomas: The member has a diagnosis of grade 2 astrocytoma or oligodendroglioma AND The member has a documented isocitrate dehydrogenate-1 (IDH1) or isocitrate dehydrogenate-2 (IDH2) mutation AND The member is using Voranigo (vorasidenib) as single agent following surgery (i.e., biopsy, sub-total resection, or gross total resection).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

VOSEVI - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Retreatment of Chronic Hepatitis C. The member must have a diagnosis of chronic hepatitis C (HCV). Member must be non-cirrhotic OR have compensated cirrhosis (Child-Pugh Class A). The member must have HCV genotype documented prior to therapy. The member has relapsed after completing a full course of or cannot use generic sofosbuvir/velpatasvir. (Note: Previous treatment with sofosbuvir/velpatasvir does not apply if member has had prior treatment failure with a full course of another NS5A-inhibitor based regimen [(e.g., glecaprevir/pibrentasvir therapy)] OR a regimen containing sofosbuvir without an NS5A inhibitor).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Pending CMS Review
Other Criteria	
Part B Prerequisite	0

VOWST - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Prevention of recurrent Clostridioides difficile infection (CDI): Member has experienced at least one episode of Clostridioides difficile infection (CDI) and Vowst will be used to prevent recurrence of CDI AND Member has completed OR will complete CDI standard of care treatment (defined as 10-21 days of treatment with vancomycin 125 mg PO QID and/or fidaxomicin 200 mg PO BID) within 2-4 days before initiating Vowst AND Member will perform a bowel cleanse using magnesium citrate or polyethylene glycol electrolyte solution on the day prior to first dose of Vowst.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

VRAYLAR - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Schizophrenia: The member must be utilizing Vraylar for the treatment of schizophrenia AND The member has tried or cannot use at least one of the following: risperidone, olanzapine, quetiapine, ziprasidone, aripiprazole, or lurasidone. Bipolar I Disorder (manic or mixed episode): The member must be utilizing Vraylar for the treatment of bipolar I disorder (manic or mixed episode) AND The member has tried or cannot use at least one of the following: risperidone, olanzapine, quetiapine, ziprasidone, or aripiprazole. Bipolar 1 Disorder (Bipolar Depression): The member must have a diagnosis of bipolar 1 disorder (bipolar depression) and the member has tried or cannot use quetiapine or lurasidone. Major Depressive Disorder: The member has a clinical diagnosis of major depressive disorder (MDD) AND The member has tried or cannot use at least one antidepressant used as monotherapy for MDD AND The member has tried or cannot use at least one generic oral atypical antipsychotic that has been used with antidepressant therapy AND Vraylar must be used with antidepressant therapy and not as monotherapy.
Age Restriction	Schizophrenia: The member is 13 years of age or older. Bipolar I Disorder (manic or mixed episode): The member is 10 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

VYLOY - VIAL (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member has experienced disease progression on Vyloy (zolbetuximab-clzb).
Required Medical Information	Gastric or Gastroesophageal Junction Adenocarcinoma (GEJ): The member has a diagnosis of locally advanced unresectable or metastatic, human epidermal growth factor receptor 2 (HER2)-negative, gastric or gastroesophageal junction (GEJ) adenocarcinoma AND The disease is claudin (CLDN) 18.2 positive as determined by an FDA approved test AND Vyloy (zolbetuximab-clzb) will be used in combination with a fluoropyrimidine- (fluorouracil [5-FU], capecitabine) and platinum-containing chemotherapy as first-line therapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 months duration
Other Criteria	
Part B Prerequisite	0

WEGOVY - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

WELIREG - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	von Hippel Lindau (VHL) Disease and Renal Cell Carcinoma (RCC): Member experiences disease progression on Welireg (belzutifan).
Required Medical Information	Pending CMS Review
Age Restriction	Pheochromocytoma or Paraganglioma (PPGL): The member is 12 years and older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

WINREVAIR - KIT

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pulmonary Arterial Hypertension (PAH): The member must have a diagnosis of pulmonary arterial hypertension (WHO Group 1) confirmed by right heart catheterization AND Member has WHO functional class II to IV symptoms AND One of the following: Member is stable on combination therapy (e.g. PDE5i (e.g. sildenafil or tadalafil) or Adempas (riociguat) and an ERA (e.g. ambrisentan, bosentan, macitentan)) OR Member is stable on PAH monotherapy (e.g. PDE5i, sGC, ERA) due to intolerance of combination therapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

XALKORI - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	ALCL: The member is greater than 1 year of age up to young adult (21 years of age). IMT: The member is greater than 1 year of age and older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

XATMEP - SOLUTION, ORAL

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members with disease progression on Xatmep (methotrexate) (applies to acute lymphoblastic leukemia only).
Required Medical Information	Acute Lymphoblastic Leukemia (ALL): The member has a diagnosis of acute lymphoblastic leukemia AND The member will be using Xatmep (methotrexate) as part of a multi-phase, combination chemotherapy maintenance regimen AND The member has had previous treatment or intolerance to generic methotrexate. Polyarticular Juvenile Idiopathic Arthritis (pJIA): The member has a diagnosis of active polyarticular juvenile idiopathic arthritis (pJIA) AND The member has had previous treatment or intolerance to generic methotrexate. BvsD Coverage Determination may also be required.
Age Restriction	The member is less than 18 years of age.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

XCOPRI - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Partial-Onset (Focal) Seizures: The member has a diagnosis of partial-onset (focal) seizures AND Member has been unable to achieve seizure control with at least TWO other drugs for controlling partial-onset seizures (e.g., levetiracetam, lamotrigine, carbamazepine, zonisamide, divalproex, gabapentin or topiramate).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

XCOPRI MAINTENANCE PACK - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Partial-Onset (Focal) Seizures: The member has a diagnosis of partial-onset (focal) seizures AND Member has been unable to achieve seizure control with at least TWO other drugs for controlling partial-onset seizures (e.g., levetiracetam, lamotrigine, carbamazepine, zonisamide, divalproex, gabapentin or topiramate).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

XCOPRI TITRATION PACK - TABLET, DOSE PACK

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Partial-Onset (Focal) Seizures: The member has a diagnosis of partial-onset (focal) seizures AND Member has been unable to achieve seizure control with at least TWO other drugs for controlling partial-onset seizures (e.g., levetiracetam, lamotrigine, carbamazepine, zonisamide, divalproex, gabapentin or topiramate).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

XDEMVY - DROPS

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Demodex blepharitis: Has a diagnosis of Demodex blepharitis.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

XERMELO - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Carcinoid Syndrome. The member has a diagnosis of carcinoid syndrome diarrhea inadequately controlled with somatostatin analog (SSA) therapy AND member has received 3 months of SSA therapy (i.e., octreotide acetate, Sandostatin LAR, or Somatuline Depot) at a stable dose AND Xermelo will be used concomitantly with SSA therapy. Reauthorization: Member has not experienced severe constipation or abdominal pain with Xermelo AND Xermelo continues to demonstrate clinical benefit, as evidenced by maintained reduction in the number of daily bowel movements from baseline AND Xermelo will continue to be used concomitantly with SSA therapy.
Age Restriction	The member must be 18 years or older.
Prescriber Restriction	Licensed Practitioner.
Coverage Duration	6 months duration.
Other Criteria	
Part B Prerequisite	0

XIFAXAN - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Traveler's diarrhea: Member must have traveler's diarrhea caused by non-invasive strains of Escherichia Coli AND Member's condition is not complicated by fever or blood in the stool AND Member has tried or cannot use ciprofloxacin, levofloxacin, or azithromycin. Hepatic encephalopathy prophylaxis: Member has had at least one overt episode of hepatic encephalopathy (HE) AND Xifaxan (rifaximin) is being used to reduce the risk of HE recurrence. Irritable bowel syndrome with diarrhea (IBS-D): Member has a diagnosis of Irritable bowel syndrome with diarrhea (IBS-D).
Age Restriction	Must be age 12 or older for Travelers Diarrhea, Age 18 or older for Hepatic encephalopathy prophylaxis and IBS-D.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

XOLAIR - SYRINGE (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Chronic Idiopathic Urticaria. Member has a diagnosis of chronic idiopathic urticaria. Member has remained symptomatic despite at least 2 weeks of H1 antihistamine therapy, unless contraindicated. Member will continue to receive H1 antihistamine therapy while on Xolair, unless contraindicated. Moderate or Severe persistent asthma. The patient has a diagnosis of moderate or severe persistent asthma. The patient has evidence of specific allergic sensitivity confirmed by positive skin test (i.e. prick/puncture test) or blood test (i.e. RAST) for a specific IgE or in vitro reactivity to a perennial aeroallergen. Member must have baseline serum IgE greater than or equal to 30 IU/ml. The patient has inadequately controlled asthma despite the use of Inhaled Corticosteroids.
Age Restriction	The patient is 12 years of age or older for diagnosis of chronic idiopathic urticaria. The patient is 6 years of age or older for moderate to severe persistent asthma. The patient is 18 years of age or older for nasal polyps. The patient is 1 year or older for IgE Mediated Food Allergy.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	Nasal Polyps: The member must meet all of the following criteria: have a diagnosis of nasal polyps (e.g., Chronic Rhinosinusitis with Nasal Polyposis [CRSwNP]) AND Xolair will be used in combination with a daily intranasal corticosteroid spray AND is unable to achieve adequate control of symptoms with maximum tolerated intranasal corticosteroid therapy. IgE Mediated Food Allergy: Member has been diagnosed with IgE mediated food allergy AND Member is using Xolair to reduce allergic reactions (Type I), including anaphylaxis, that may occur with accidental exposure to one of more foods.
Part B Prerequisite	0

XOSPATA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Acute Myeloid Leukemia. The member has a diagnosis of acute myeloid leukemia AND The member has relapsed or refractory disease AND The member has documented FLT3 mutation positive disease AND The member will be using Xospata (gilteritinib) as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

XPOVIO - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression on Xpovio (selinexor).
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

XTANDI - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on Xtandi (enzalutamide).
Required Medical Information	Prostate Cancer (metastatic castration-resistant): The member has metastatic castration-resistant prostate cancer (CRPC). Prostate Cancer (non-metastatic castration-resistant): The member has a diagnosis of non-metastatic castration-resistant prostate cancer AND The member will use Xtandi (enzalutamide) in combination with androgen deprivation therapy (e.g. previous bilateral orchiectomy or luteinizing hormone-releasing hormone (LHRH) analog [i.e., LHRH agonist/ antagonist]). Prostate Cancer (metastatic castration-sensitive): the member has a diagnosis of metastatic castration-sensitive prostate cancer AND the member will use Xtandi (enzalutamide) in combination with androgen deprivation therapy (e.g. previous bilateral orchiectomy or luteinizing hormone-releasing hormone (LHRH) analog [i.e., LHRH agonist/ antagonist]). Prostate Cancer (non-metastatic castration-sensitive): The member has a diagnosis of non-metastatic castration-sensitive prostate cancer (nmCSPC) with biochemical recurrence at high risk for metastasis (biochemical recurrence at high risk for metastasis defined as prostate-specific antigen [PSA] doubling time less than or equal to 9 months, PSA greater than or equal to 1 ng/mL post radical prostatectomy, or screening PSA greater than or equal to 2 ng/mL above nadir post radiotherapy).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

YERVOY - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Melanoma: Concomitant Zelboraf (vemurafenib), Tafinlar (dabrafenib), Cotellic (cobimetnib) or Mekinist (trametinib) therapy AND The member has had progression of disease on adjuvant therapy with Yervoy (ipilimumab). MSI-H or dMMR: Disease progression while on or following previous anti-CTLA4-based combinations (e.g., Opdivo [nivolumab] with Yervoy [ipilimumab]). Hepatocellular carcinoma: Disease progression while on or following previous anti-CTLA-based combinations (e.g., Imjudo [tremelimumab-actl] with Imfinzi [durvalumab]).
Required Medical Information	Melanoma. The member has a diagnosis of unresectable or metastatic melanoma OR Adjuvant treatment of cutaneous melanoma with pathologic involvement of regional lymph nodes of more than 1 millimeter who have undergone complete resection, including total lymphadenectomy. The member is naive to Yervoy (ipilimumab).The member has an Eastern Cooperative Oncology Group (ECOG) performance status of 0-2. Melanoma - Reauthorization Criteria unresectable or metastatic. The member had stable disease, partial response or complete response for greater than 3 months following the completion of initial induction (completion of four cycles within a 16 week period. Members who were unable to tolerate or receive the complete induction regimen within 16 weeks of initiation will not receive approval). AND The member has progressive disease, necessitating reinduction therapy with Yervoy (ipilimumab). AND The member has an Eastern Cooperative Oncology Group (ECOG) performance status of 0-2. Reauth adjuvant treatment of cutaneous melanoma. The member has not had disease recurrence or unacceptable toxicity with Yervoy (ipilimumab) AND The total duration of treatment is less than 3 years AND The member has an ECOG performance status of 0-2. Renal Cell Carcinoma. The member has a diagnosis of advanced renal cell carcinoma (RCC) AND The member has predominant clear cell histology AND The member will be using Yervoy (ipilimumab) in combination with Opdivo (nivolumab) AND The member will be using for first line therapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	4 Months Duration

YERVOY - VIAL (ML)

Other Criteria	Microsatellite Instability-High (MSI-H) or Mismatch Repair Deficient (dMMR) Metastatic Colorectal Cancer. The member has a diagnosis of unresectable or metastatic colorectal cancer with documented microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR) AND The member will be using Yervoy (ipilimumab) in combination with Opdivo (nivolumab). Hepatocellular carcinoma: The member has a diagnosis of hepatocellular carcinoma AND The member will be using Yervoy (ipilimumab) in combination with Opdivo (nivolumab). Non-small cell lung cancer (NSCLC) -- First Line Therapy: The member must have a diagnosis of metastatic non-small cell lung cancer (NSCLC) AND one of the following applies: Disease with PD-L1 expression greater than or equal to 1% with no EGFR or ALK genomic tumor aberrations and given as first line therapy AND Tumor expresses PD-L1 as determined by an FDA-approved test AND Will be used in combination with Opdivo (nivolumab) OR Disease with no EGFR or ALK genomic tumor aberrations and given as first line therapy AND Will be used in combination with Opdivo (nivolumab) AND Will be used in combination with two cycles of platinum doublet chemotherapy. Malignant pleural mesothelioma: The member has a diagnosis of unresectable malignant pleural mesothelioma AND The member will be using for first-line or subsequent treatment, if not administered first-line AND Yervoy (ipilimumab) will be used in combination with Opdivo (nivolumab). Esophageal Cancer (squamous cell carcinoma): the member has unresectable advanced, recurrent, or metastatic squamous cell carcinoma of the esophagus AND Yervoy will be given as first line treatment in combination with Opdivo.
Part B Prerequisite	0

YESINTEK - SYRINGE (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	Moderate to severe chronic plaque psoriasis and psoriatic arthritis: The member must be 6 years of age or older. For Ulcerative Colitis: The member must be 18 years of age or older. For Crohn's Disease: The member must be 2 years of age or older.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

YONDELIS - VIAL (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Member experiences disease progression on Yondelis (trabectedin)
Required Medical Information	Liposarcoma/Leiomyosarcoma:The member has unresectable or metastatic liposarcoma or leiomyosarcoma AND The member has received prior anthracycline (e.g., doxorubicin) containing regimen. Soft Tissue Sarcoma. Yondelis (trabectedin) will be used as monotherapy for palliative treatment and one of the following applies: The member has a diagnosis of unresectable or progressive retroperitoneal or intraabdominal soft tissue sarcoma OR the member has a diagnosis of angiosarcoma or rhabdomyosarcoma OR the member has a diagnosis stage IV soft tissue sarcoma of the extremity/superficial trunk, head/neck, or recurrent disease with disseminated metastases.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

PA Criteria	Criteria Details
Off-Label Uses	Pending CMS Review
Exclusion Criteria	Pending CMS Review
Required Medical Information	Pending CMS Review
Age Restriction	Pending CMS Review
Prescriber Restriction	Pending CMS Review
Coverage Duration	Pending CMS Review
Other Criteria	Pending CMS Review
Part B Prerequisite	Pending CMS Review

ZARXIO - SYRINGE (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	6 Months Duration
Other Criteria	Febrile Neutropenia Prophylaxis, in members with acute myeloid leukemia receiving chemotherapy: The member must have a diagnosis Acute Myeloid Leukemia (AML). The member must be receiving either induction chemotherapy OR consolidation Chemotherapy AND The member is not administering filgrastim product earlier than 24 hours after cytotoxic chemotherapy or within 24 hours before chemotherapy. Febrile Neutropenia Prophylaxis, In non-myeloid malignancies following Hematopoietic Stem Cell Transplant (HCST): The member must have had a Hematopoietic Stem Cell Transplant (HCST) (e.g. bone marrow transplant, peripheral-blood progenitor cell (PBPC) transplant) for a non-myeloid malignancy AND The member is not administering filgrastim product earlier than 24 hours after cytotoxic chemotherapy or within 24 hours before chemotherapy Harvesting of peripheral blood stem cells: The member is using as a component of a hematopoietic stem cell mobilization protocol and one of the following applies: is scheduled for autologous peripheral-blood stem cell (PBSC) transplantation, is storing cells for a possible future autologous transplant, or is donating stem cells for an allogeneic or syngeneic PBSC transplant. Neutropenic disorder, chronic (Severe), Symptomatic: The member must have a diagnosis of congenital, cyclic, or idiopathic neutropenia. Neutropenia in AIDS patients: The member must have a diagnosis of AIDS with neutropenia. Treatment of Aplastic Anemia: The member must have a diagnosis of Aplastic Anemia. Treatment of Agranulocytosis: The member must have a diagnosis of congenital or drug induced agranulocytosis. Hematopoietic Syndrome of Acute radiation syndrome: The member has been acutely exposed to myelosuppressive doses of radiation.
Part B Prerequisite	0

ZEGFROVY - TABLET

PA Criteria	Criteria Details
Off-Label Uses	Pending CMS Review
Exclusion Criteria	Pending CMS Review
Required Medical Information	Pending CMS Review
Age Restriction	Pending CMS Review
Prescriber Restriction	Pending CMS Review
Coverage Duration	Pending CMS Review
Other Criteria	Pending CMS Review
Part B Prerequisite	Pending CMS Review

ZEJULA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members that have experienced disease progression while on or following PARP inhibitor therapy [e.g., Rubraca (rucaparib), Lynparza (olaparib), Zejula (niraparib)].
Required Medical Information	Epithelial Ovarian Cancer, Fallopian Tube, or Peritoneal Cancer subsequent line maintenance therapy: The member has a diagnosis of recurrent epithelial ovarian cancer, fallopian tube cancer, or primary peritoneal cancer AND The member is in complete or partial response to their last platinum regimen AND The member will utilize Zejula (niraparib) as a monotherapy* AND *Discontinue Bevacizumab product before initiating maintenance therapy with Zejula. Advanced Ovarian Cancer, Fallopian Tube, or Peritoneal Cancer First Line Maintenance Therapy: member has a diagnosis of advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer AND Members disease is associated with homologous recombination deficiency (HRD) positive status defined by either: a deleterious OR suspected deleterious BRCA mutation or genomic instability, as defined by an FDA approved test. Member is in complete response or partial response to first line treatment with platinum based chemotherapy AND member will utilize Zejula (niraparib) as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ZELBORAF - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Members on concomitant Yervoy (ipilimumab), Opdivo (nivolumab), Keytruda (pembrolizumab), Tafinlar (dabrafenib), Mekinist (trametinib), Braftovi (encorafenib), or Mektovi (binimetinib). Members that have experienced disease progression while on Zelboraf (vemurafenib). Members that have experienced disease progression while on prior anti-BRAF/MEK combination therapy [e.g. Braftovi (encorafenib) with Mektovi (binimetinib) or Tafinlar (dabrafenib) with Mekinst (trametinib)].
Required Medical Information	Pending CMS Review
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ZEMAIRA - VIAL (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	IgA deficient members or presence of antibodies against IgA.
Required Medical Information	The member must meet ALL of the following criteria: Diagnosis of congenital alpha1-antitrypsin deficiency with clinically evident emphysema and chronic replacement therapy is needed AND Has an alpha1-antitrypsin phenotype of PiZZ, PiZ(null), or PI (null, null) or phenotypes associated with reduced serum alpha-1 antitrypsin concentrations, AND an initial serum alpha 1-antitrypsin concentration of less than 57mg/dL when measured by nephelometry OR less than 80 mg/dL when measured by radial immunodiffusion. (These products should not be used in individuals with the PiMZ or PiMS phenotypes of alpha1-antitrypsin deficiency because these individuals appear to be at small risk of developing clinically evident emphysema.)
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ZENBEXUS - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	Pending CMS Review
Exclusion Criteria	Pending CMS Review
Required Medical Information	Pending CMS Review
Age Restriction	Pending CMS Review
Prescriber Restriction	Pending CMS Review
Coverage Duration	Pending CMS Review
Other Criteria	Pending CMS Review
Part B Prerequisite	Pending CMS Review

ZIRABEV - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Malignant Pleural Mesothelioma, Soft Tissue Sarcoma, Recurrent Primary CNS Tumor (including Glioblastoma multiforme), Renal Cell Carcinoma, Non-small cell lung cancer (non-squamous cell histology): The member has experienced disease progression while on or following a bevacizumab product. Metastatic colorectal cancer: A bevacizumab product may not be used in conjunction with Vectibix (panitumumab) or Erbitux (cetuximab).
Required Medical Information	For Avastin (bevacizumab), Alymsys (bevacizumab-maly), Jobevne (bevacizumab-nwgd), and Vegzelma (bevacizumab-adcd) oncology requests: member must have an intolerance or contraindication with Mvasi or Zirabev. Metastatic colorectal cancer: metastatic colorectal cancer AND 1 of the following applies: using a bevacizumab product in combo with fluoropyrimidine (e.g., 5-fluorouracil or capecitabine) based chemo for 1st or 2nd-line therapy OR in combo with fluoropyrimidine-irinotecan or fluoropyrimidine-oxaliplatin-based chemo for 2nd-line therapy in patients who have progressed on 1st-line bevacizumab-containing regimens. Non-small cell lung cancer (non-squamous cell histology). NSCLC with non-squamous cell histology AND the disease is unresectable, locally advanced, recurrent, or metastatic AND A bevacizumab product will be used as part of a supported combination chemotherapy. Hepatocellular carcinoma: unresectable or metastatic HCC AND a bevacizumab product will be used as 1st line therapy in combo with an atezolizumab product.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration

ZIRABEV - VIAL (ML)

Other Criteria	Renal cell carcinoma: Member has RCC and a bevacizumab product will be used in combo with interferon alpha. Recurrent Primary CNS Tumor (including Glioblastoma multiforme): Diagnosis of progressive or recurrent glioblastoma or anaplastic glioma AND a bevacizumab product is being used as a single agent or in combo with irinotecan, carmustine, lomustine or temozolomide AND the member does not have a CNS hemorrhage. Soft Tissue Sarcoma: One of the following apples: Diagnosis of angiosarcoma AND a bevacizumab product is being used as a single agent OR The member has a diagnosis of solitary fibrous tumor and a bevacizumab product is being used in combo with temozolomide. Macular Retinal Edema. Avastin is being used to treat central or branch retinal vein occlusion with macular retinal edema. Cervical Cancer: member has recurrent, or metastatic cervical cancer AND A bevacizumab product will be used as part of a supported combination therapy. Endometrial Cancer: The member has a diagnosis of endometrial cancer AND One of the following applies: The member has stage III-IV disease or recurrent disease AND a bevacizumab product will be used in combination with carboplatin and paclitaxel, followed by maintenance with a bevacizumab product maintenance OR The member has recurrent disease AND a bevacizumab product monotherapy will be used as second-line or subsequent therapy if not previously used. Malignant Pleural Mesothelioma. Diagnosis of unresectable malignant pleural mesothelioma and a bevacizumab product will be used in combo with cisplatin and pemetrexed or carboplatin and pemetrexed followed by maintenance with a bevacizumab product (for responders). Epithelial ovarian, fallopian tube, or primary peritoneal cancer: Diagnosis of epithelial ovarian, fallopian tube or primary peritoneal cancer AND One of the following applies: The disease is recurrent, platinum-resistant AND a bevacizumab product will be used in combination with paclitaxel, pegylated liposomal doxorubicin or topotecan AND the member has not received more than 2 prior chemotherapy regimens OR The disease is recurrent, platinum-sensitive AND a bevacizumab product will be used in combination with carboplatin and paclitaxel, or in combination with carboplatin and gemcitabine, followed by a bevacizumab product monotherapy AND The disease is stage III or IV AND initial surgical resection was completed AND a bevacizumab product will be used in combo with carboplatin and paclitaxel, followed by a bevacizumab product monotherapy OR . Age Related Macular Degeneration (Avastin requests only): Avastin (bevacizumab) is being used to treat age related macular degeneration. Diabetic Macular Edema (Avastin requests only): Avastin (bevacizumab) is being used to treat diabetic macular edema.
Part B Prerequisite	0

zoledronic acid-mannitol-water - INTRAVENOUS SOLUTION, PIGGYBACK (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Osteoporosis: The member has a diagnosis of osteoporosis. Has new fractures or significant loss of bone mineral density despite previous treatment, contraindication, or intolerance to any other oral bisphosphonate. Osteoporosis Prophylaxis in postmenopausal members: The member is postmenopausal. Has new fractures or significant loss of bone mineral density despite previous treatment, contraindication, or intolerance to any other oral bisphosphonate. Glucocorticoid induced Osteoporosis in men and women taking systemic glucocorticoids: Diagnosis of glucocorticoid induced osteoporosis or is either initiating or continuing systemic glucocorticoids with a daily dosage equivalent of 7.5 mg or greater of prednisone and is expected to remain on glucocorticoids for at least 12 months AND has new fractures or significant loss of bone mineral density despite previous treatment, contraindication, or intolerance to any other oral bisphosphonate . Paget's Disease: Diagnosis of Paget's disease. The member has continued elevation(s) in serum alkaline phosphatase (SAP) of two times or higher than the upper limit of normal despite previous treatment, contraindication, or intolerance to any other oral bisphosphonate. And the member: Is symptomatic OR is at risk for complications from their disease, to induce remission, or to normalize serum alkaline phosphatase. Brand Reclast request only: Members must have previous treatment, contraindication, or intolerance to generic Zoledronic acid (generic Reclast).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ZOLINZA - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Cutaneous T-Cell Lymphoma (CTCL). The member has a diagnosis of progressive, persistent, or recurrent disease or The member will be using Zolinza (vorinostat) as primary treatment or adjuvant therapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ZONISADE - SUSPENSION, ORAL (FINAL DOSE FORM)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Partial-Onset (Focal) Seizures: Member has a diagnosis of partial-onset (focal) seizures AND Zonisade (zonisamide) will be used in combination with at least one other drug for controlling partial-onset seizures AND Member has tried or cannot use zonisamide capsules AND at least one other drug for controlling partial-onset seizures (e.g., levetiracetam, lamotrigine, carbamazepine, divalproex, gabapentin or topiramate).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

ZORYVE - CREAM (GRAM)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Atopic Dermatitis: The member has a diagnosis of mild to moderate atopic dermatitis AND The member has tried or cannot use one of the following topical products: triamcinolone, mometasone, betamethasone, or clobetasol cream/ointment OR The member has tried or cannot use a topical calcineurin inhibitor (e.g. tacrolimus). Previous treatment with a calcineurin inhibitor is not required for children less than 2 years of age.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

ZTALMY - SUSPENSION, ORAL (FINAL DOSE FORM)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Seizures associated with Cyclin-dependent Kinase-Like 5 (CDKL5) Deficiency Disorder (CDD): the member has a diagnosis of cyclin-dependent kinase-like 5 CDKL5) deficiency disorder (CDD).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ZURZUVAE - CAPSULE

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Postpartum Depression. The member has a documented diagnosis of a major depressive episode as defined by DSM-5 criteria and severity as determined by an appropriate depression rating scale (e.g. HAM-D, MADRS, PHQ-9) with onset of symptoms in the third trimester of pregnancy or within 4 weeks of delivery AND The member will be receiving Zurzuvae therapy no later than one year postpartum AND The member is not pregnant at initiation of Zurzuvae therapy AND The member will be receiving no more than one 14-day course of treatment with Zurzuvae for the current episode of postpartum depression.
Age Restriction	The member is at least 18 years of age.
Prescriber Restriction	Licensed Practitioner
Coverage Duration	30 day duration
Other Criteria	
Part B Prerequisite	0

ZYDELIG - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	The member has experienced disease progression while on or following a PI3K inhibitor (e.g. idelalisib, copanlisib).
Required Medical Information	Chronic Lymphocytic Leukemia (CLL): The member must have a diagnosis of relapsed OR refractory chronic lymphocytic leukemia (CLL).
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ZYKADIA - TABLET

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Non-small Cell Lung Cancer (NSCLC): The member has a diagnosis of metastatic and documented anaplastic lymphoma kinase (ALK)-positive non-small cell lung cancer (NSCLC) AND the member has a medical reason as to why Alecensa (alectinib), Alunbrig (brigatinib) or Lorbrena (lorlatinib) cannot be started or continued AND member will be using Zykadia (ceritinib) as monotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan Year Duration
Other Criteria	
Part B Prerequisite	0

ZYMFENTRA - PEN INJECTOR KIT (EA)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	
Required Medical Information	Ulcerative Colitis: The member has a diagnosis of moderately to severely active ulcerative colitis AND The member has tried or cannot use TWO of the following: preferred adalimumab product [i.e., adalimumab-adbm, adalimumab-adaz], a preferred ustekinumab product (i.e., Otulfi, Yesintek), Skyrizi, Rinvoq, or Tremfya OR The member has completed induction doses with an intravenous Infliximab product. Crohn's Disease: The member has a diagnosis of moderately to severely active Crohn's disease AND The member has tried or cannot use TWO of the following: preferred adalimumab product [i.e., adalimumab-adbm, adalimumab-adaz], preferred ustekinumab product (i.e., Otulfi, Yesintek), Skyrizi, Rinvoq, or Tremfya OR The member has completed induction doses with an intravenous Infliximab product.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Plan year duration
Other Criteria	
Part B Prerequisite	0

ZYNYZ - VIAL (ML)

PA Criteria	Criteria Details
Off-Label Uses	
Exclusion Criteria	Anal carcinoma: Disease progression on previous anti-PD-1/PD-L1 therapy (e.g., pembrolizumab, nivolumab) AND For first line therapy: total treatment duration with Zynyz (retifanlimab-dlwr) is more than 12 months.
Required Medical Information	Merkel Cell Carcinoma: The member has a diagnosis of recurrent locally advanced or metastatic merkel cell carcinoma AND Zynyz (retifanlimab-dlwr) will be used as monotherapy. Anal carcinoma: The member has a diagnosis of inoperable locally recurrent or metastatic squamous cell carcinoma of the anal canal (SCAC) AND One of the following applies: Zynyz (retifanlimab-dlwr) will be used as first line therapy in combination with carboplatin and paclitaxel, followed by Zynyz (retifanlimab-dlwr) monotherapy OR Zynyz (retifanlimab-dlwr) will be used as a single agent after disease progression or intolerance to platinum-based chemotherapy.
Age Restriction	
Prescriber Restriction	Licensed Practitioner
Coverage Duration	Six months duration
Other Criteria	
Part B Prerequisite	0

The following drugs are subject to a Medicare Part B or D authorization depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.

PART B VERSUS PART D

Products Affected

- Akynzeo (netupitant) 300 mg-0.5 mg capsule
- aprepitant 125 mg (1)-80 mg (2) capsules in a dose pack
- aprepitant 125 MG; 125 mg (1)- 80 MG (2); 40 MG; 80 MG; capsule
- Astagraf XL 0.5 MG; 1 MG; 5 MG; capsule,extended release
- Azasan 100 MG; 75 MG; tablet
- azathioprine 100 MG; 50 MG; 75 MG; tablet
- CellCept 200 MG/ML; 250 MG; 500 MG; capsule
- CellCept 200 MG/ML; 250 MG; 500 MG; oral suspension
- CellCept 200 MG/ML; 250 MG; 500 MG; tablet
- CellCept Intravenous 500 MG; intravenous solution
- chlorpromazine 10 MG; 25 MG; tablet
- Compazine 10 MG; 5 MG; tablet
- cyclophosphamide 25 MG; 50 MG; capsule
- cyclophosphamide 25 MG; 50 MG; tablet
- cyclosporine 100 MG; 25 MG; capsule
- cyclosporine modified 100 MG; 100 MG/ML; 25 MG; 50 MG; capsule
- cyclosporine modified 100 MG; 100 MG/ML; 25 MG; 50 MG; oral solution
- dronabinol 10 MG; 2.5 MG; 5 MG; 5 MG/ML; capsule
- dronabinol 10 MG; 2.5 MG; 5 MG; 5 MG/ML; oral solution
- Emend 125 mg (1)- 80 MG (2); 125 mg (25 mg/ ML FINAL CONC.); 80 MG; capsule
- Emend 125 mg (1)-80 mg (2) capsules in a dose pack
- Emend 125 mg (25 mg/mL final conc.) oral suspension
- Envarsus XR 0.75 MG; 1 MG; 4 MG; tablet,extended release
- everolimus (immunosuppressive) 0.25 MG; 0.5 MG; 0.75 MG; 1 MG; tablet
- Gengraf 100 MG; 100 MG/ML; 25 MG; capsule
- Gengraf 100 MG; 100 MG/ML; 25 MG; oral solution
- granisetron HCl 1 MG; tablet
- Granisol 1 MG/5 ML; oral solution
- Imuran 50 MG; tablet
- Jylamvo 2 MG/ML; oral solution
- Marinol 10 MG; 2.5 MG; 5 MG; capsule
- Medrol 16 MG; 2 MG; 4 MG; 8 MG; tablet
- methotrexate sodium 2.5 MG; tablet
- methylprednisolone 16 MG; 32 MG; 4 MG; 8 MG; tablet
- Millipred 5 MG; tablet
- mycophenolate 500 MG; intravenous solution
- mycophenolate mofetil 200 MG/ML; 250 MG; 500 MG; capsule
- mycophenolate mofetil 200 MG/ML; 250 MG; 500 MG; oral powder for suspension
- mycophenolate mofetil 200 MG/ML; 250 MG; 500 MG; tablet
- mycophenolate sodium 180 MG; 360 MG; tablet,delayed release

- Myfortic 180 MG; 360 MG; tablet,delayed release
- Myhibbin 200 MG/ML; oral suspension
- Neoral 100 MG; 100 MG/ML; 25 MG; capsule
- Neoral 100 MG; 100 MG/ML; 25 MG; oral solution
- ondansetron 16 MG; 4 MG; 8 MG; disintegrating tablet
- ondansetron HCl 4 MG; 4 MG/5 ML; 8 MG; oral solution
- ondansetron HCl 4 MG; 4 MG/5 ML; 8 MG; tablet
- Orapred ODT 10 MG; 15 MG; 30 MG; disintegrating tablet
- Pediapred 5 mg base/5 mL (6.7 MG/5 ML); oral solution
- prednisolone 10 MG; 10 MG/5 ML; 15 MG; 15 mg/5 mL (3 MG/ML); 20 mg/5 mL (4 MG/ML); 25 mg/5 mL (5 MG/ML); 30 MG; 5 mg base/5 mL (6.7 MG/5 ML); disintegrating tablet
- prednisolone 15 MG/5 ML; 5 MG; oral solution
- prednisolone 15 MG/5 ML; 5 MG; tablet
- prednisolone sodium phosphate 10 MG; 10 MG/5 ML; 15 MG; 15 mg/5 mL (3 MG/ML); 20 mg/5 mL (4 MG/ML); 25 mg/5 mL (5 MG/ML); 30 MG; 5 mg base/5 mL (6.7 MG/5 ML); oral soln
- prednisolone sodium phosphate 10 MG; 10 MG/5 ML; 15 MG; 15 mg/5 mL (3 MG/ML); 20 mg/5 mL (4 MG/ML); 25 mg/5 mL (5 MG/ML); 30 MG; 5 mg base/5 mL (6.7 MG/5 ML); oral solution
- prednisone 1 MG; 10 MG; 2 MG; 2.5 MG; 20 MG; 5 MG; 5 MG/5 ML; 50 MG; oral solution
- prednisone 1 MG; 10 MG; 2 MG; 2.5 MG; 20 MG; 5 MG; 5 MG/5 ML; 50 MG; tablet
- prednisone 1 MG; 10 MG; 2 MG; 2.5 MG; 20 MG; 5 MG; 5 MG/5 ML; 50 MG; tablet,delayed release
- Prednisone Intensol 5 MG/ML; oral concentrate
- prochlorperazine maleate 10 MG; 5 MG; tablet
- Prograf 0.2 MG; 0.5 MG; 1 MG; 5 MG; capsule
- Prograf 0.2 MG; 0.5 MG; 1 MG; 5 MG; oral granules in packet
- Rayos 1 MG; 2 MG; 5 MG; tablet,delayed release
- Sandimmune 100 MG; 25 MG; capsule
- sirolimus 0.5 MG; 1 MG; 1 MG/ML; 2 MG; oral solution
- sirolimus 0.5 MG; 1 MG; 1 MG/ML; 2 MG; tablet
- Syndros 5 MG/ML; oral solution
- tacrolimus 0.5 MG; 1 MG; 5 MG; capsule, immediate-release
- tacrolimus XL 0.5 MG; 1 MG; 5 MG; capsule,extended release 24 hr
- Trexall 10 MG; 15 MG; 5 MG; 7.5 MG; tablet
- trimethobenzamide 300 MG; capsule
- Varubi 90 MG; tablet
- Veripred 20 20 mg/5 mL (4 MG/ML); oral solution
- Xatmep 2.5 MG/ML; oral solution
- Zortress 0.25 MG; 0.5 MG; 0.75 MG; 1 MG; tablet

Notice of Availability - Auxiliary Aids and Services Notice

English: Free language, auxiliary aid, and alternate format services are available. Call **1-800-794-5907 (TTY: 711)**.

العربية [Arabic]: تتوفر خدمات اللغة والمساعدة الإضافية والتنسِّيق البديل مجاناً. اتصل على الرقم **1-800-794-5907 (الهاتف النصي: 711)**.

Հայերեն [Armenian]: Հասանելի են անվճար լեզվական, աջակցման և այլընտրանքային ձևաչափի ծառայություններ: Զանգահարեք **1-800-794-5907 (TTY՝ 711)** հեռախոսահամարով:

বাংলা [Bengali]: বিনামূল্যে ভাষা, আনুষঙ্গিক সহায়তা, এবং বিকল্প বিন্যাসে পরিষেবা উপলব্ধ। ফোন করুন **1-800-794-5907 (TTY: 711)** নম্বরে।

简体中文 [Simplified Chinese]：我们可提供免费的语言、辅助设备以及其他格式版本服务。请致电 **1-800-794-5907**（听障专线：**711**）。

繁體中文 [Traditional Chinese]：我們可提供免費的語言、輔助設備以及其他格式版本服務。請致電 **1-800-794-5907**（聽障專線：**711**）。

Kreyòl Ayisyen [Haitian Creole]: Gen sèvis lengwistik gratis, èd oksilyè, ak lòt fòma sèvis ki disponib. Rele **1-800-794-5907 (TTY: 711)**.

Hrvatski [Croatian]: Dostupne su besplatne jezične usluge, usluge dodatne pomoći i alternativnog formata. Nazovite **1-800-794-5907 (TTY: 711)**.

فارسی [Farsi]: خدمات زبان رایگان، کمک های اضافی و فرمت های جایگزین در دسترس است. با **1-800-794-5907 (TTY: 711)** تماس بگیرید.

Français [French] : Des services gratuits d’assistance linguistiques, d’aide auxiliaire et de format alternatif sont disponibles. Appelez le **1-800-794-5907 (ATS : 711)**.

Deutsch [German]: Es stehen kostenlose unterstützende Hilfs- und Sprachdienste sowie alternative Dokumentformate zur Verfügung. Telefon: **1-800-794-5907 (TTY: 711)**.

Ελληνικά [Greek]: Διατίθενται δωρεάν γλωσσικές υπηρεσίες, βοηθήματα και υπηρεσίες σε εναλλακτικές προσβάσιμες μορφές. Καλέστε στο **1-800-794-5907 (TTY: 711)**.

ગુજરાતી [Gujarati]: મફત ભાષા, સહાયક સહાય અને વૈકલ્પિક ફોર્મેટ સેવાઓ ઉપલબ્ધ છે. **1-800-794-5907 (TTY: 711)** પર કૉલ કરો.

עברית [Hebrew]: שירותים אלה זמינים בחינם: שירותי תרגום, אביזרי עזר וטקסטים בפורמטים חלופיים. נא התקשר למספר **1-800-794-5907 (TTY: 711)**

Hmoob [Hmong]: Muaj kev pab txhais lus, pab kom hnov suab, thiab lwm tus qauv pab cuam. Hu **1-800-794-5907 (TTY: 711)**.

Italiano [Italian]: Sono disponibili servizi gratuiti di supporto linguistico, assistenza ausiliaria e formati alternativi. Chiamare il numero **1-800-794-5907 (TTY: 711)**.

GHHNOA2026CP_0526

日本語 [Japanese]: 言語支援サービス、補助支援サービス、代替形式サービスを無料でご利用いただけます。**1-800-794-5907 (TTY: 711)** までお電話ください。

ភាសាខ្មែរ [Khmer]៖ សេវាកម្មផ្នែកភាសា ជំនួយ និង សេវាកម្មជាទម្រង់ផ្សេងជំនួសអាចរកបាន។ ទូរសព្ទទៅលេខ **1-800-794-5907 (TTY: 711)**។

한국어 [Korean]: 무료 언어, 보조 지원 및 대체 형식 서비스를 이용하실 수 있습니다. **1-800-794-5907 (TTY: 711)** 번으로 문의하십시오.

Diné [Navajo]: Saad bee áná'álwo', bee haz'áanii, dóó naaltsoos t'áá ajiłii'ígíí bee ak'e'elyaaígíí t'áá jiik'eh hóló. **1-800-794-5907 (TTY: 711)** bich'į' hodíílnih.

Polski [Polish]: Dostępne są bezpłatne usługi językowe, pomocnicze i alternatywne formaty. Zadzwoń pod numer **1-800-794-5907 (TTY: 711)**.

Português [Brazilian Portuguese]: Estão disponíveis serviços gratuitos de ajuda linguística auxiliar e outros formatos alternativos. Ligue **1-800-794-5907 (TTY: 711)**.

ਪੰਜਾਬੀ [Punjabi]: ਮੁਫ਼ਤ ਭਾਸ਼ਾ, ਸਹਾਇਕ ਸਹਾਇਤਾ, ਅਤੇ ਵਿਕਲਪਿਕ ਫਾਰਮੈਟ ਸੇਵਾਵਾਂ ਉਪਲਬਧ ਹਨ। **1-800-794-5907 (TTY: 711)** 'ਤੇ ਕਾਲ ਕਰੋ।

Русский [Russian]: Предоставляются бесплатные услуги языковой поддержки, вспомогательные средства и материалы в альтернативных форматах. Звоните по номеру **1-800-794-5907 (TTY: 711)**.

Español [Spanish]: Los servicios gratuitos de asistencia lingüística, ayuda auxiliar y servicios en otro formato están disponibles. Llame al **1-800-794-5907 (TTY: 711)**.

Tagalog [Tagalog]: May available na mga libreng serbisyo sa wika, mga gamit na pantulong, at alternatibong format. Tumawag sa **1-800-794-5907 (TTY: 711)**.

தமிழ் [Tamil]: மொழி,, உதவிப் பொருட்கள் மற்றும் மாற்று வடிவமைப்புச் சேவைகள் இலவசமாகக் கிடைக்கின்றன. **1-800-794-5907 (TTY: 711)** எண்ணை அழைக்கவும்.

తెలుగు [Telugu]: ఉచిత భాష, సహాయక మద్దతు, మరియు ప్రత్యామ్నాయ ఫార్మాట్ సేవలు అందుబాటులో గలవు. **1-800-794-5907 (TTY: 711)** కి కాల్ చేయండి.

اردو [Urdu]: مفت زبان، معاون امداد، اور متبادل فارمیٹ کی خدمات دستیاب ہیں۔ کال **1-800-794-5907 (TTY: 711)**۔

Tiếng Việt [Vietnamese]: Có sẵn miễn phí các dịch vụ ngôn ngữ, hỗ trợ phụ trợ và định dạng thay thế. Hãy gọi **1-800-794-5907 (TTY: 711)**.

This notice is available at **CarePlusHealthPlans.com/MLI**