



2025 LI NET pharmacy point-of-sale safety edits

Effective Jan. 1, 2025

The Limited Income NET (LI NET) Program employs several point-of-sale safety edits, prompting additional safety reviews to determine if the prescribed medication is appropriate and medically necessary. Dispensing pharmacists should utilize their clinical knowledge and judgment to resolve and override with the updated drug utilization review (DUR)/professional pharmacy service (PPS) codes and International Classification of Diseases, 10th Revision (ICD-10) diagnosis code entry overrides. LI NET point-of-sale edits appear under the “Documents and forms” tab on [Humana.com/Member/Medicare-LINET-Pharmacy-Resources](https://www.humana.com/Member/Medicare-LINET-Pharmacy-Resources).

DUR/PPS code functionality allowed

Claims will display the message “Soft Reject Payer Allows DUR/PPS Code Override” or “DUR Reject” in the National Council for Prescription Drug Programs (NCPDP) field. Based on the type of safety edit, use the charts below to enter the correct “Reason for service,” “Professional service” and “Result of service” codes for successful claim adjudication. The review steps include:

1. Reviewing pharmacy records to identify the reason for rejection (therapeutic duplication, interactions, inappropriate dosage)
 - A. Rejections may result because of multiple edits occurring concurrently.
2. Consulting with the patient or their prescriber to confirm the appropriateness of the prescribed medications and determine current medications to exclude any therapy changes
3. Overriding the rejection as indicated below if pharmacy data and/or the prescriber/patient confirms appropriateness of the prescribed drug therapy and the pharmacist approves the prescription fill

For questions, please call the LI NET help desk at **800-783-1307**, Monday – Friday, 8 a.m. – 7 p.m., Eastern time.

Table A

Safety edit description	Reason for service code	Professional service code	Result of service code
<p>Morphine milligram equivalent – opioid care coordination</p> <p>The cumulative morphine milligram equivalent (MME) daily dosage is limited across all opioid prescriptions to a predetermined lower threshold between 90 MME and 200 MME (and the patient receives an opioid from more than two prescribers and two pharmacies).</p> <p>Reject codes: NCPDP 88: DUR reject error NCPDP 922: MORPHINE MILLIGRAM EQUIVALENT (MME) EXCEEDS LIMITS</p>	<p>HD: High dose</p>	<p>M0: Prescriber consulted DE: Dosing evaluation DP: Dosage evaluated</p>	<p>1B: Filled prescription as is 1D: Filled with different directions 1F: Filled with different quantity 1G: Filled with prescriber approval 4A: Prescribed with acknowledgments 4B: Dispensed, palliative care 4C: Dispensed, hospice (Florida Medicaid only) 4D: Dispensed, cancer treatment 4K: Prescriber specialty exemption-oncology or non-hospice palliative care 4L: Prescriber specialty exemption-hospice</p>
<p>Morphine milligram equivalent – opioid care coordination</p> <p>The cumulative MME daily dosage is limited across all opioid prescriptions to a predetermined upper threshold of greater than 200 MME.</p> <p>Reject codes: NCPDP 88: DUR reject error NCPDP 922: MORPHINE MILLIGRAM EQUIVALENT (MME) EXCEEDS LIMITS</p>	<p>ER: Overuse</p>	<p>M0: Prescriber consulted</p>	<p>4B: Dispensed, palliative care 4L: Prescriber specialty exemption-hospice</p>

Safety edit description	Reason for service code	Professional service code	Result of service code
<p>Polypharmacy edits</p> <ul style="list-style-type: none"> Concurrent use of two or more unique anticholinergic medications in patients 65 years old and older <p>Reject code: NCPDP 88: DUR reject error</p>	DD: Drug-drug interaction	DE: Dosing evaluation M0: Prescriber consulted MP: Patient will be monitored PE: Patient education/instruction PO: Patient consulted RO: Pharmacist consulted other source SW: Literature search/review	1A: Filled as is, false positive 1B: Filled prescription as is 1D: Filled with different directions 1F: Filled with different quantity 1G: Filled with prescriber approval 4A: Prescribed with acknowledgments 4B: Dispensed, palliative care 4D: Dispensed, cancer treatment
<p>Drug-to-disease interactions</p> <p>Potential conflict between medication claims and diagnosis in patient's history.</p> <p>Reject code: NCPDP 70: Product/service not covered – plan/benefit exclusion</p>	DC: Drug disease	DE: Dosing evaluation M0: Prescriber consulted MP: Patient will be monitored RO: Pharmacist consulted other source SW: Literature search/review	1A: Filled as is, false positive 1B: Filled prescription as is 1D: Filled with different directions 1F: Filled with different quantity 1G: Filled with prescriber approval 4A: Prescribed with acknowledgments 4B: Dispensed, palliative care 4D: Dispensed, cancer treatment
<p>Drug-to-drug interactions</p> <p>Interactions include concurrent opioid and benzodiazepine usage.</p> <p>Patient's prescription history detects potential interactions between two or more medications.</p> <p>Reject code: NCPDP 88: DUR reject error; additional messaging: This drug interacts with the patient's other drug(s).</p>	DD: Drug-drug interaction OR AT: Additive toxicity (use for opioid and benzodiazepine interaction)	DE: Dosing evaluation M0: Prescriber consulted MP: Patient will be monitored PE: Patient education/instruction PO: Patient consulted RO: Pharmacist consulted other source SW: Literature search/review	1A: Filled as is, false positive 1B: Filled prescription as is 1D: Filled with different directions 1F: Filled with different quantity 1G: Filled with prescriber approval 4A: Prescribed with acknowledgments 4B: Dispensed, palliative care 4D: Dispensed, cancer treatment

Safety edit description	Reason for service code	Professional service code	Result of service code
<p>Duplicate therapy Potential therapeutic or ingredient duplications are based on duplicate therapy classes.</p> <p>Reject code: NCPDP 88: DUR reject error; additional messaging: This drug interacts with the patient's other drug(s).</p> <p>Note: Pharmacy processing for some therapeutic duplications may vary depending on if the prescription fill attempt is the initial fill or a subsequent fill. Please see Table B below for examples.</p>	TD: Therapeutic duplication	M0: Prescriber consulted PE: Patient education/instruction PO: Patient consulted RO: Pharmacist consulted other source SW: Literature search/review TH: Therapeutic product interchange	1A: Filled as is, false positive 1B: Filled prescription as is 1D: Filled with different directions 1F: Filled with different quantity 1G: Filled with prescriber approval 4A: Prescribed with acknowledgments 4B: Dispensed, palliative care 4D: Dispensed, cancer treatment

Table B

Duplicate therapy edits	Pharmacy processing on initial prescription fill Note: For PPS eligible, see codes above under duplicate therapy	Pharmacy processing on subsequent prescription fills
Diuretics – aldosterone receptor antagonist	PPS eligible	Prior authorization required
Janus kinase inhibitors	PPS eligible	Prior authorization required
Antiplatelet and antithrombotic drugs (selected group two)	PPS eligible	Prior authorization required

Opioid naïve edit

The pharmacy system will result in a soft or hard reject, which can be overridden if a patient meets the appropriate eligible exemptions. The review steps are below:

1. Review pharmacy records to confirm the patient has not received any opioid prescriptions within the last 108 days (i.e., patient is opioid naïve).
2. If the patient is opioid naïve, identify if they have an eligible exemption using pharmacy records or consult with the patient's prescriber. Please see the table below for eligible exemptions.
3. If pharmacy data or the prescriber confirms an exemption, enter the ICD-10 diagnosis code at the point of sale to override the rejection.

Table C

Safety edit	Safety edit description and processing
Opioid naïve – seven-day supply limit	<p>Patients who have not had an opioid prescription within the past 108 days (i.e., new to opioid therapy) are limited to a supply of seven days or less.</p> <p>Reject codes: NCPDP 88: DUR reject error NCPDP 925: initial fill days supply exceeds limit DUR messaging: <i>DUR Message 1 – OPIOID NAÏVE; DUR Message 2 – <insert number> DAY MAX. FOR SICKLE CELL, CANCER, CHRONIC PAIN, USE ICD-10 CODE TO OVERRIDE.</i></p> <p>Pharmacy processing The pharmacist at the point of sale may override the rejection to allow for paid claims utilizing eligible ICD-10 codes if a patient has an appropriate exemption (sickle cell disease, cancer diagnosis, palliative care, hospice, chronic pain management diagnosis [i.e., G89, M25, M47, M50, M51 or M54]).</p> <p>Note: Patients new to LI NET plans also will trigger this edit, and appropriate override codes should be entered if they are not opioid naïve.</p> <p>Subsequent prescriptions filled within 108 days will not reject as the patient will no longer be identified as opioid naïve.</p>

Prior authorization is required for the following edits:

Table D

Safety edit	Safety edit description	Reject code
Opioid days' supply limitation	Opioid claims are limited to a 30-day supply, but a 31-day supply per fill is allowed for residents in long-term care (LTC) facilities. This includes both short-acting and long-acting medications.	NCPDP 76: Plan limitations exceeded; additional messaging: Days' supply greater than maximum allowed for this plan.
Benzodiazepine days' supply limitation	Benzodiazepine claims are limited to a 30-day supply, but a 31-day supply per fill is allowed for residents in LTC facilities.	NCPDP 76: Plan limitations exceeded; additional messaging: Days' supply greater than maximum allowed for this plan.
Antipsychotic use in patients with dementia	Patients who are 65 years old and older, have a diagnosis of dementia, and are prescribed an antipsychotic will require prior authorization.	NCPDP 88: DUR reject error; additional messaging: atypical antipsychotic alert
Concurrent use of any opioid medication with a benzodiazepine medication	Overlapping fills of opioid and benzodiazepine medication will require a coverage determination.	NCPDP 88: DUR reject error; additional messaging: This drug interacts with patient's other drug(s).

Safety edit	Safety edit description	Reject code
<p>MME threshold limits</p> <p>Note: MME thresholds may vary by line of business and/or state requirements.</p>	<p>Patients are filling opioid medication doses greater than allowed MME dosing.</p>	<p>NCPDP 88: DUR reject error; additional messaging: Cumulative morphine equivalent dose exceeds limits. NCPDP 922: Morphine milligram equivalent (MME) exceeds limits NCPDP G4: Prescriber must contact plan</p>
<p>Drug-to-disease interactions</p> <p>Potential conflict between medication claims and diagnosis in patient's history (some specific drug-to-disease interactions will require a prior authorization)</p>	<p>There is a potential conflict between medication claims and diagnosis in the patient's history.</p>	<p>NCPDP 70: Product/service not covered – plan/benefit exclusion</p>

The patient's prescriber can submit a request for a prior authorization by calling LI NET at **800-783-1307**.