

Care Plus disorder Major depression/Major depressive

Clinical overview

Definitions

The American Psychiatric Association's Diagnostic Statistical Manual of Mental Disorders, Fifth Edition (DSM-5) classifies clinical depression (major depressive disorder) as feeling sad, low or worthless most days for at least two weeks while also having other symptoms such as sleep problems, loss of interest in activities or change in appetite. This is the most severe form of depression and one of the most common forms. Providers may use criteria in the DSM-5 to diagnose this condition.1

Causes²

The exact cause is not known. As with many mental disorders, a variety of factors may be involved, such as:

- Biological differences/physical changes in the brain
- Brain chemistry, hormones or inherited traits
- Traumatic or stressful events

Signs and symptoms²

During these episodes, symptoms occur most of the day, nearly every day and may include:

- Feelings of sadness, tearfulness, emptiness or hopelessness
- Angry outbursts, irritability or frustration, even over small matters
- Sleep disturbances and lack of energy
- · Reduced appetite and weight loss or increased cravings for food and weight gain
- Anxiety, agitation or restlessness
- Trouble thinking, concentrating, making decisions and remembering things
- Frequent or recurrent thoughts of death or suicide

Complications²

- Alcohol or substance abuse
- Social isolation, family conflicts or relationship difficulties
- Excess weight or obesity, which can lead to heart disease and diabetes
- Suicidal feelings, suicide attempts or suicide

Diagnostic tools

- PHQ-9 (a nine-item patient health questionnaire which consists of the criteria upon which the diagnosis and severity of depressive disorders is based)3
- Psychological evaluation²
- Physical exam

Treatment²

- Psychotherapy/mental health counseling
- Medications (linked to diagnosis)
- Brain stimulation therapies (e.g., electroconvulsive therapy [ECT] or transcranial magnetic stimulation [TMS])



Best documentation practices for healthcare providers

Subjective

In the subjective section of the office note, document the presence or absence of any current signs or symptoms related to major depressive disorder (e.g., depressed mood, insomnia, feeling of worthlessness, etc.).

Objective

The objective section should include physical exam findings (e.g., flat affect, weight loss or gain, etc.) and related assessment results, such as PHQ-9 scores.

Assessment

Specificity: Describe each final diagnosis clearly, concisely and to the highest level of specificity. Use all applicable descriptors and include the following:

- Episode single or recurrent
- Severity mild, moderate, severe
- Presence or absence of psychosis/psychotic features
- Remission status partial or full

Treatment plan

- Document a clear and concise treatment plan for major depressive disorder linking related medications to the diagnosis.
- Indicate in the office note to whom or where any referral or consultation requests are made.
- Document when the patient will be seen again, even if only on an as-needed basis.

Coding tips

Major depression coexisting with bipolar disorder

- Major depression coexisting with bipolar disorder classifies to the applicable combination code under category F31 for bipolar disorder. Bipolar disorder includes both depression and mania and it is more important to capture the bipolar disorder. Therefore, a code for depression is not reported separately.⁴
- Category F31.- Bipolar disorder, has an *Excludes1* note advising categories F32.-; major depressive disorder, single episode and F33.- major depressive disorder, recurrent are NOT coded here! An *Excludes1* note indicates that the code excluded should never be used at the same time as the code above the *Excludes1* note. An *Excludes1* note is used when two conditions cannot occur together.⁶

Additional reminders

- While MDD is a commonly accepted medical abbreviation for major depressive disorder, this abbreviation also can be used to represent manic depressive disorder, which classifies to a different diagnosis code.
- "Chronic depression" and "depression" without further description both code to F32.A, Depression, unspecified.⁵



Coding examples

Example 1	
Medical record HPI: Depression disorder	
documentation Past medical history (PMH): Bipolar depression	
Psychiatric/Behavioral review of systems: Positive for sleep disturban	nce; patient is
nervous/anxious; depression	
Psychiatric exam: She is attentive; Affect is flat and angry.	
Assessment: Moderate episode of recurrent major depressive disorder wi	th documented
treatment.	
ICD-10-CM code F33.1 Major depressive disorder, recurrent, moderate	
Rationale Bipolar depression is documented as historical in the PMH. Only the modern	ate recurrent
depression is documented and supported in this example.	
Example 2	
Medical record HPI: Depression	
documentation Assessment: Recurrent major depression – patient is asymptomatic, symptomatic,	otoms are
stable. Patient denies suicidal ideation. Plan is to continue current manager	
EHR-inserted code: F33.0 (Major depressive disorder, recurrent, mild)	
ICD-10-CM code F33.9 Major depressive disorder, recurrent, unspecified	
Rationale The severity of mild, moderate, or severe with or without psychotic symptom	ms or features is
not documented by the provider. As there is a mismatch between the provi	
diagnosis and the EHR-inserted code descriptor, with no additional support	
record, code assignment is based on the provider's diagnostic statement.6	
Example 3	
Medical record HPI: Follow-up on chronic conditions, including depression	
documentation Problems List: Recurrent major depression	
Assessment: Endogenous depression stable on current medications	
ICD-10-CM code F33.2 Major depressive disorder, recurrent severe without psychotic fea	tures
Rationale Review of the record in its entirety supports a current diagnosis of endoger	
The coding path for endogenous depression leads to F33.2, even in the abs	
specificity of severity documentation by the provider. Depression (acute) (r	
endogenous (without psychotic symptoms) > F33.2 ⁷	,
Example 4	
Medical record Diagnosis #1: Depression	
documentation Diagnosis #5: Major depressive disorder, recurrent in partial remission	
Diagnosis #12: Major depressive disorder, recurrent in full remission	
Diagnosis #15: Major depressive disorder, single episode, severe without p	sychotic features
ICD-10-CM code Query the provider for clarification	
Rationale When multiple variations of Major depressive disorder are documented in t	he same
encounter, no ICD-10 code can be assigned without a provider query to co	
level. This example illustrates why it is important to update your problem li	
Assessment, etc., to prevent contradiction in the documentation.	



References

- Cleveland Clinic. Depression Symptoms, Causes, & Treatment Cleveland Clinic. https://my.clevelandclinic.org/health/diseases/9290-depression. Published January 13, 2023. Accessed February 3, 2025.
- 2. Mayo Clinic. Depression (Major Depressive Disorder). Mayo Clinic. https://www.mayoclinic.org/diseases-conditions/depression/symptoms-causes/syc-20356007. Published October 14, 2022. Accessed January 31, 2025.
- 3. Kroenke K, Spitzer RL, Williams, JBW. The PHQ-9: Validity of a brief depression severity measure. *Journal of General Internal Medicine*. 2001;16(9):606-613. Doi: HTTPs://doi.org/10.1046/j.1525-1497.2001.016009606.x
- 4. American Hospital Association (AHA). Bipolar disorder and recurrent major depressive disorder. *Coding Clinic, First Quarter, 2020.* Published online 2020:23.
- 5. American Hospital Association (AHA). Depression not otherwise specified. *Coding Clinic, Fourth Quarter, 2021*. Published online 2021:9-10.
- 6. CMS. *ICD-10-CM Official Guidelines for Coding and Reporting*.; 2024. https://www.cms.gov/files/document/fy-2025-icd-10-cm-coding-quidelines.pdf Accessed October 3, 2024.
- 7. AAPC. ICD-10-CM Complete Code Set 2025. AAPC; 2024.

