

Best documentation practices for healthcare

Prostate – specific antigen (PSA)

PSA is a protein produced by cells of the prostate gland that often is elevated in men with prostate cancer. The PSA laboratory test measures the level of PSA in a man's blood. PSA testing often is used with a digital rectal examination of the prostate gland to screen for prostate cancer. Other noncancerous conditions, however, can cause an elevated PSA, for example, inflammation or enlargement of the prostate gland.

Generally, a PSA level of 4.0 ng/mL or less is considered normal. Recent studies have shown, however, that some men with PSA levels below 4.0 have prostate cancer, while other men with higher levels do not have prostate cancer. There are various causes of fluctuations in PSA levels, but generally the higher the PSA level, the more likely the diagnosis of prostate cancer. A continuous rise in PSA level over time may be a sign of prostate cancer.¹

Prostate cancer treatment options²

- Surgery – radical prostatectomy or TURP (Transurethral Resection of the Prostate)
- Radiation therapy – external beam radiation, internal radiation therapy (brachytherapy), proton beam therapy
- Medical therapy – chemotherapy, hormone therapy, immunotherapy
- Other treatments – cryosurgery, HIFU (high-intensity focused ultrasound), active surveillance (watchful waiting)

Hormone therapy for prostate cancer³

Most prostate cancer cells rely on testosterone to help them grow. Hormone therapy (also known as androgen deprivation therapy [ADT] or androgen suppression therapy) stops the body from producing testosterone or stops testosterone from reaching prostate cancer cells, which often makes prostate cancers shrink or grow more slowly for a time.

Hormone therapy alone, however, does not cure prostate cancer. Hormone therapy includes:

- **Orchiectomy** (surgical castration) – Surgical removal of the testicles, which reduces testosterone levels in the body quickly and significantly. This option is permanent and irreversible.
- **Drug therapy** (sometimes called chemical or medical castration)
 - **Luteinizing hormone-releasing hormone (LHRH) agonists and antagonists** – Medications that stop the body from producing testosterone.
 - **Anti-androgens** – Medications that block testosterone from reaching prostate cancer cells. They are usually given in conjunction with LHRH agonists, as LHRH agonists can cause a temporary increase in testosterone before levels decrease.
 - **Other medications** – When prostate cancer persists or recurs despite hormone therapy, other medications can be used to block testosterone in the body. Each medication targets testosterone in the body in a different way. Examples include corticosteroids such as prednisone and the anti-fungal drug ketoconazole.

Radioactive seed implantation⁴

Radiation therapy for prostate cancer using implanted radioactive seeds is called brachytherapy. There are

two types of brachytherapy:

- Permanent (low dose rate): Implanted seeds are left in place, but they become biologically inert (no longer release radiation) after a period of weeks or months.
- Temporary (high dose rate): Seeds are implanted and left in place for about 30 minutes; the seeds and radioactive material then are removed.

Current versus historical prostate cancer

- Do not use the phrase “history of” to describe current prostate cancer. In diagnosis coding, “history of” means the condition is historical and no longer exists as a current problem.
- On the other hand, do not document a final diagnosis of simply “prostate cancer” to describe a historical prostate cancer that was previously excised or eradicated and for which there is:
 - No active treatment, *and*
 - No evidence of disease or recurrence, it is appropriate to document “history of prostate cancer” with details of past diagnosis and treatment.

Purpose of PSA blood testing¹

After successful treatment of prostate cancer that results in complete eradication of the disease, regular monitoring for recurrence typically continues and includes ongoing laboratory testing of PSA. Documentation of ongoing PSA monitoring by itself without evidence that the prostate cancer is still present does not clearly support current prostate cancer. A final impression and plan of “prostate cancer – check PSA” is vague and ambiguous; it does not clearly indicate current versus historical prostate cancer or the purpose of checking the PSA level.

Treatment with outcomes

Document the details of prostate cancer treatment along with the outcome and current status.

- Include beginning and ending dates of all treatment.
- For permanent radioactive seed implantation (also known as brachytherapy or internal radiation therapy):
 - Include the date seed implantation was performed.
 - Indicate when seeds are considered no longer active and thus, therapy is complete.
- Clearly document when prostate cancer has been eradicated.

Coding tips

Coding prostate cancer

Cancer or carcinoma of the prostate codes as follows:

- C61 Malignant neoplasm of prostate
- C79.82 Secondary malignant neoplasm, genital organs
- D07.5 Carcinoma in situ of prostate

Carefully review and follow all instructional notes.

Coding prostate cancer as current

Generally, prostate cancer is coded as current when the medical record clearly documents active treatment directed to the cancer for the purpose of cure or palliation and/or when the record clearly shows prostate cancer is still present but:

- Is unresponsive to treatment;
- The current treatment plan is observation only or “watchful waiting;” or
- The patient has refused any further treatment.

Coding prostate cancer as historical⁵

- Prostate cancer is coded as historical (Z85.46) after the prostate cancer has been excised or eradicated, there is no active treatment directed to the prostate cancer and there is currently no evidence of disease or recurrence.
- Encounter for follow-up examination after treatment for malignant neoplasm has been completed is coded as Z08. Carefully review and follow all instructional notes.

Metastatic prostate cancer⁶

For a current diagnosis of “metastatic prostate cancer” without further specification – and no more definitive information regarding the primary versus secondary site is obtained upon review of the entire medical record – the prostate is coded as the primary site (C61) with the secondary/metastatic site unknown (C79.9).

Coding examples

| Example 1 | |
|-------------------------------------|--|
| Medical record documentation | 79-year-old gentleman with a history of prostate cancer, here for yearly f/u. Patient had labs prior to today’s visit. No chest pain or SOB. No bowel or bladder complaints PMH: History of prostate cancer, prostatectomy ROS: As noted in the HPI Impression: Carcinoma of prostate. Patient did well with the treatments. PSA is low again this year. |
| ICD-10-CM codes | Query the provider for clarification regarding the current status of prostate cancer. Documentation is vague and does not provide sufficient information for accurate diagnosis code assignment. |
| Rationale | Prostate cancer is coded as historical (Z85.46) after the prostate cancer has been excised or eradicated, there is no active treatment directed to the prostate cancer and there is currently no evidence of disease or recurrence. ⁵ |

| Example 2 | |
|-------------------------------------|---|
| Medical record documentation | 66-year-old male here for refill. PMH of diabetes. Patient is currently under treatment for adenocarcinoma of the prostate with pelvic bone metastasis. A1C is currently controlled with medication. Assessment & Plan: Diabetes, current A1C 5.4%, will continue current medications. Malignant neoplasm of prostate with metastasis to the pelvic bone. Has f/u with oncologist scheduled for tomorrow-will advise his office staff to send updated report. |
| ICD-10-CM codes | E11.9 Type 2 diabetes mellitus without complications C61 Malignant neoplasm of prostate C79.51 Secondary malignant neoplasm of bone |
| Rationale | The documentation clearly identifies the prostate as the primary site and the pelvic bone as the secondary site. The plan of care describes both neoplasms as active. |

| Example 3 | |
|-------------------------------------|--|
| Medical record documentation | HPI: Mr. Smith presents today with history of prostate cancer, currently undergoing treatment for metastatic carcinoma to pelvic bone. Previous prostatectomy with no recurrence at primary site. Assessment: Prostate carcinoma with metastasis of pelvic bone |

| | |
|------------------------|---|
| ICD-10-CM codes | <p>C79.51 Secondary malignant neoplasm of bone</p> <p>Z85.46 Personal history of malignant neoplasm of prostate</p> |
| Rationale | <p>The documentation clearly identifies there is no recurrence at primary site. Prostate cancer is coded as historical (Z85.46) after the prostate cancer has been excised or eradicated, there is no active treatment directed to the prostate cancer and there is currently no evidence of disease or recurrence.⁵</p> |

References

1. National Cancer Institute. Prostate Cancer Treatment (PDQ®)—Patient Version - National Cancer Institute. www.cancer.gov. Published October 9, 2020. https://www.cancer.gov/types/prostate/patient/prostate-treatment-pdq#_142
2. National Cancer Institute. Prostate-specific antigen (PSA) test. National Cancer Institute. Published 2022. <https://www.cancer.gov/types/prostate/psa-fact-sheet>
3. Mayo Clinic. Hormone therapy for prostate cancer - Mayo Clinic. MayoClinic.org. Published 2019. <https://www.mayoclinic.org/tests-procedures/hormone-therapy-for-prostate-cancer/about/pac-20384737>
4. WebMD. Prostate Cancer: Radioactive Seed Implants. WebMD. Published March 14, 2014. Accessed April 30, 2025. <https://www.webmd.com/prostate-cancer/radioactive-seed-implants>
5. CMS. *ICD-10-CM Official Guidelines for Coding and Reporting*; 2024. Accessed October 3, 2024. <https://www.cms.gov/files/document/fy-2025-icd-10-cm-coding-guidelines.pdf>
6. Leon-Chisen N, Harper D, Love T, Young-Charles G. *ICD-10-CM and ICD-10-PCS Coding Handbook 2025 with Answers*. American Hospital Association; 2024.