

Provider clinical quality

OVERVIEW OF SCORE

Effective beneficiary care is important for both military readiness and better outcomes over time. Provider clinical quality evaluates the clinical quality of care delivered by Primary Care Managers (PCM) and specialists. The clinical quality measure helps identify those providers who offer higher clinical quality care. With a possible set of 101 measures, providers have numerous opportunities to impact care for both preventive and specialty care. Providers are compared to their peers' performance in the same geography. Those providers who perform well on their specialty-specific measure set are associated with better clinical quality of care for the TRICARE population.

Scores are calculated only where Humana Military requirements are met - when there is sufficient data volume, statistical credibility, minimum peer volume, etc.

Evidence-based measure methodology

APPROACH

Humana Military uses TRICARE East beneficiaries' claims experiences to evaluate their treatment and clinical quality of care and evaluates claims including medical, pharmacy and lab. The measures are calculated using a software solution widely used across the healthcare industry as a basis to review clinical quality delivered by providers.

GOVERNING BODIES OF THE SCORE MEASURES

The clinical quality score includes measures from the following governing bodies:

- National Committee for Quality Assurance (NCQA)
- Pharmacy Quality Alliance (PQA)
- Centers for Medicare and Medicaid Services (CMS)
- National Quality Forum (NQF)

Score process

EVALUATED SPECIALTIES

- Advanced practice midwife
- Allergy and immunology

- Cardiovascular disease
- Emergency medicine
- Endocrinology
- Family medicine
- General practice
- Internal medicine
- Mental health nurse practitioner
- Midwife
- Nurse practitioner
- Obstetrics and gynecology
- Pediatrics
- Physician assistant
- Psychiatry and neurology
- Pulmonary disease

Note: See appendix for specialties and associated measures


MEASUREMENT LEVEL

TRICARE providers are captured at the identifier level of Tax Identification Number (TIN) and National Provider Identifier (NPI).

GEOGRAPHIC AREAS

Geographic areas include states in the TRICARE East Region of the United States.

MAP OF STATES COVERED



PEER GROUPS

Peer groups include providers of the same specialty and type of care within the same state. Providers working in multiple states will be evaluated per the volume performed in each state as outlined on the following pages.

Provider clinical quality

FREQUENCY OF EVALUATION

The clinical quality scores are updated on a quarterly basis.

MINIMUM VOLUME REQUIREMENTS

The clinical quality score is measured for all providers who have a minimum of 20 measure results for their attributed beneficiaries and at least five distinct TRICARE members.

Peer groups will contain at least five peers for providers to be individually compared against.

PROVIDER ATTRIBUTION

Providers who contribute the most to a beneficiary's care are attributed. Attribution is prioritized utilizing a hierarchy per specialty that begins with the highest visit

count, most recent visit, highest claim paid amount and highest claim allowed amount.

SCORE METHODOLOGY

Humana Military scores providers for clinical quality only when there is sufficient volume and minimum number of peers available.

Humana Military uses a weighted average approach to create a composite score for individual providers and their peer group(s). Humana Military then compares the actual score of the individual provider over the expected score of the peer group(s) to create a performance index that exhibits individual performance versus peers.

The performance index is converted using a percentile evaluated at the provider specialty/state level. Each score is assigned points based on the percentile distribution.

SCORING OUTPUT EXAMPLE

Example: Provider A – Internal medicine – Jacksonville, Florida

Measure	INDIVIDUAL PROVIDER			PEER GROUP		
	Provider compliant (numerator)	Provider total (denominator)	Compliance rate	Peer compliant (numerator)	Peer total (denominator)	Compliance rate
Breast cancer screening	12	15	80.0%	233	280	83.2%
Bronchitis, acute, avoidance of antibiotic	14	15	93.3%	250	280	89.3%
Diabetes HbA1c testing	18	20	90.0%	115	120	95.8%
Diabetes, Part D med adherence	50	55	90.9%	252	300	84.0%

Actual performance ¹	Expected performance ²	Performance index ³
89.5%	86.9%	1.03

¹ Actual performance: provider's individual compliance rate

² Expected performance: provider's peer group compliance rate weighted by the number of opportunities

³ Performance index: >1 = high score, 1.0 = average score, <1 = low score

Provider clinical quality

Appendix

Covers brief descriptions of each measure; measure to provider specialty assignment; and governing body over measure.

Source type	Condition	Measure description	Specialty list
1	Adherence to Antipsychotic Medications for Individuals With Schizophrenia (SAA)*	Patient(s) with schizophrenia or schizoaffective disorder who were dispensed and remained on antipsychotic medication for at least 80% of their treatment period.	Psychiatry and Neurology
1	Adults' Access to Preventive/ Ambulatory Health Services (AAP)*	Patient(s) 20 years and older that had a preventive or ambulatory care visit during the last 12 months of the report period.	Family Medicine, Internal Medicine, Nurse Practitioner, Physician Assistant, General Practice
1	Antidepressant Medication Management (AMM)*	Patient(s) with major depression who start an antidepressant medication that remained on treatment for at least 12 weeks (effective acute phase treatment).	Mental Health Nurse Practitioner, Psychiatry and Neurology, General Practice, Internal Medicine, Family Medicine, Nurse Practitioner, Physician Assistant
1	Antidepressant Medication Management (AMM)*	Patient(s) with major depression who start an antidepressant medication that remained on treatment for at least 6 months (effective continuation phase treatment).	Physician Assistant, Mental Health Nurse Practitioner, Psychiatry and Neurology, General Practice, Internal Medicine, Family Medicine, Nurse Practitioner
1	Appropriate Testing for Pharyngitis (CWP)*	Patient(s) treated with an antibiotic for pharyngitis that had a Group A streptococcus test.	General Practice, Pediatrics, Nurse Practitioner, Family Medicine, Physician Assistant, Internal Medicine

*Indicates **HEDIS** measure; **Condition source type key:** 1 = NCQA measure, 2 = Modified NQF measure, 3 = CMS measure, 4 = PQA measure Measures generated with Optum Evidence Based Medicine (EBM) software.

Provider clinical quality

Source type	Condition	Measure description	Specialty list
1	Appropriate Treatment for Upper Respiratory Infection (URI)*	Patient(s) with a diagnosis of upper respiratory infection (URI) that did not have a prescription for an antibiotic on or within three days after the initiating visit.	Allergy and Immunology, Emergency Medicine, Family Medicine, General Practice, Internal Medicine, Nurse Practitioner, Pediatrics, Physician Assistant
2	Asthma	Adult(s) with presumed uncontrolled or partly controlled asthma using an inhaled corticosteroid or acceptable alternative.	Family Medicine, Internal Medicine, Nurse Practitioner, Physician Assistant, Allergy and Immunology, Pediatrics, General Practice
2	Asthma	Patient(s) exhibiting problematic asthma control who had pulmonary or allergy evaluation in last 12 reported months.	Nurse Practitioner, Pediatrics, Physician Assistant, Allergy and Immunology, General Practice, Family Medicine, Internal Medicine
2	Asthma	Patient(s) that did not have an asthma related emergency department encounter or observation stay in last 12 reported months.	Family Medicine, Internal Medicine, Nurse Practitioner, Pediatrics, Physician Assistant, Allergy and Immunology, General Practice
2	Asthma	Patient(s) that did not have an asthma related hospitalization in last 12 reported months	Family Medicine, Internal Medicine, Nurse Practitioner, Pediatrics, Physician Assistant
2	Asthma	Patient(s) that had an ambulatory visit for asthma care in last 12 reported months.	Allergy and Immunology, Pulmonary Disease, Emergency Medicine

*Indicates HEDIS measure; **Condition source type key:** 1 = NCQA measure, 2 = Modified NQF measure, 3 = CMS measure, 4 = PQA measure Measures generated with Optum Evidence Based Medicine (EBM) software.

Provider clinical quality

Source type	Condition	Measure description	Specialty list
2	Asthma	Pediatric patient(s) with presumed uncontrolled or partly controlled asthma using an inhaled corticosteroid or acceptable alternative.	Allergy and Immunology, Family Medicine, Nurse Practitioner, Pediatrics, Physician Assistant, Pulmonary Disease, Internal Medicine, General Practice
1	Asthma Medication Ratio (AMR)*	Patient(s) between the ages of 5 and 64 with an asthma medication ratio ≥ 0.50 during the report period.	Allergy and Immunology, Family Medicine, Internal Medicine, Nurse Practitioner, Pediatrics, Physician Assistant, Pulmonary Disease, General Practice
1	Avoidance of Antibiotic Treatment for Acute Bronchitis/ Bronchiolitis (AAB)*	Patient(s) with a diagnosis of acute bronchitis/bronchiolitis that did not have a prescription for an antibiotic on or within three days after the initiating visit.	Allergy and Immunology, Family Medicine, Internal Medicine, Nurse Practitioner, Pediatrics, Physician Assistant, General Practice
1	Breast Cancer Screening (BCS)*	Patient(s) 52 - 74 years that had a screening mammogram in last 27 reported months.	Family Medicine, Internal Medicine, Nurse Practitioner, Obstetrics and Gynecology, Physician Assistant, General Practice
2	Cerebral Vascular Accident and Transient Cerebral Ischemia - Part 1	Patient(s) with a LDL cholesterol test in last 24 reported months.	Psychiatry and Neurology, Internal Medicine, Nurse Practitioner, Physician Assistant, Family Medicine

*Indicates HEDIS measure; **Condition source type key:** 1 = NCQA measure, 2 = Modified NQF measure, 3 = CMS measure, 4 = PQA measure Measures generated with Optum Evidence Based Medicine (EBM) software.

Provider clinical quality

Source type	Condition	Measure description	Specialty list
2	Cerebral Vascular Accident and Transient Cerebral Ischemia - Part 2	Patient(s) with a recent acute cerebral ischemic event that had a head computerized axial tomography (CT) scan or magnetic resonance imaging (MRI) test soon after the acute event.	Physician Assistant, Nurse Practitioner, Psychiatry and Neurology, Family Medicine, Internal Medicine
2	Cerebral Vascular Accident and Transient Cerebral Ischemia - Part 4	Patient(s) with a recent hospitalization or observation stay for an acute cerebral ischemic event that had any provider visit within 30 days of hospital discharge.	Internal Medicine, Psychiatry and Neurology, Physician Assistant, Family Medicine, Nurse Practitioner
1	Cervical Cancer Screening (CCS)*	Women that had appropriate screening for cervical cancer (Commercial enrollment).	Family Medicine, General Practice, Internal Medicine, Midwife, Nurse Practitioner, Obstetrics and Gynecology, Physician Assistant
1	Child and Adolescent Well-Care Visits (WCV)*	Patient(s) 12 - 17 years that had at least one comprehensive well-care visit with a PCP or an OB/GYN practitioner in the last 12 reported months.	Family Medicine, Internal Medicine, Nurse Practitioner, Pediatrics, Physician Assistant, General Practice
1	Child and Adolescent Well-Care Visits (WCV)*	Patient(s) 18 - 21 years that had at least one comprehensive well-care visit with a PCP or an OB/GYN practitioner in the last 12 reported months.	Family Medicine, Internal Medicine, Nurse Practitioner, Pediatrics, Physician Assistant, General Practice
1	Child and Adolescent Well-Care Visits (WCV)*	Patient(s) 3 - 11 years that had at least one comprehensive well-care visit with a PCP or an OB/GYN practitioner in the last 12 reported months.	Family Medicine, Nurse Practitioner, Pediatrics, Physician Assistant, General Practice
1	Child and Adolescent Well-Care Visits (WCV)*	Patient(s) 3 - 21 years that had at least one comprehensive well-care visit with a PCP or an OB/GYN practitioner in the last 12 reported months.	Family Medicine, Internal Medicine, Nurse Practitioner, Pediatrics, Physician Assistant, General Practice

*Indicates HEDIS measure; **Condition source type key:** 1 = NCQA measure, 2 = Modified NQF measure, 3 = CMS measure, 4 = PQA measure Measures generated with Optum Evidence Based Medicine (EBM) software.

Provider clinical quality

Source type	Condition	Measure description	Specialty list
1	Childhood Immunization Status (CIS)*	Patient(s) 2 years old at the end of the report period that had a varicella immunization between their 1st and 2nd birthday.	Pediatrics, Family Medicine, Nurse Practitioner, Physician Assistant, General Practice
1	Childhood Immunization Status (CIS)*	Patient(s) 2 years old at the end of the report period that had an MMR immunization between their 1st and 2nd birthday.	Pediatrics, Family Medicine, Nurse Practitioner, Physician Assistant, General Practice
1	Childhood Immunization Status (CIS)*	Patient(s) 2 years old at the end of the report period that had four DTaP immunizations by their 2nd birthday.	Pediatrics, Family Medicine, Nurse Practitioner, Physician Assistant, General Practice
1	Childhood Immunization Status (CIS)*	Patient(s) 2 years old at the end of the report period that had four pneumococcal conjugate immunizations by their 2nd birthday.	Pediatrics, Family Medicine, Nurse Practitioner, Physician Assistant, General Practice
1	Childhood Immunization Status (CIS)*	Patient(s) 2 years old at the end of the report period that had one hepatitis A immunization between their 1st and 2nd birthday.	Pediatrics, Family Medicine, Nurse Practitioner, Physician Assistant, General Practice
1	Childhood Immunization Status (CIS)*	Patient(s) 2 years old at the end of the report period that had the required number of rotavirus immunizations by their 2nd birthday.	Pediatrics, Family Medicine, Nurse Practitioner, Physician Assistant, General Practice
1	Childhood Immunization Status (CIS)*	Patient(s) 2 years old at the end of the report period that had three hepatitis B immunizations by their 2nd birthday.	Pediatrics, Family Medicine, Nurse Practitioner, Physician Assistant, General Practice

*Indicates HEDIS measure; **Condition source type key:** 1 = NCQA measure, 2 = Modified NQF measure, 3 = CMS measure, 4 = PQA measure Measures generated with Optum Evidence Based Medicine (EBM) software.

Provider clinical quality

Source type	Condition	Measure description	Specialty list
1	Childhood Immunization Status (CIS)*	Patient(s) 2 years old at the end of the report period that had three HiB immunizations by their 2nd birthday.	Pediatrics, Family Medicine, Nurse Practitioner, Physician Assistant, General Practice
1	Childhood Immunization Status (CIS)*	Patient(s) 2 years old at the end of the report period that had three polio vaccinations by their 2nd birthday.	Pediatrics, Family Medicine, Nurse Practitioner, Physician Assistant, General Practice
1	Childhood Immunization Status (CIS)*	Patient(s) 2 years old at the end of the report period that had two influenza immunizations by their 2nd birthday.	Pediatrics, Family Medicine, Nurse Practitioner, Physician Assistant, General Practice
1	Chlamydia Screening in Women (CHL)*	Patient(s) 16 - 24 years that had a chlamydia screening test in last 12 reported months.	Family Medicine, Internal Medicine, Midwife, Nurse Practitioner, Obstetrics and Gynecology, Pediatrics, Physician Assistant, General Practice
1	Colorectal Cancer Screening (COL)*	Patient(s) 50 - 75 years that had appropriate screening for colorectal cancer.	Family Medicine, Internal Medicine, Nurse Practitioner, Physician Assistant, General Practice
1	Comprehensive Diabetes Care (CDC)*	Patient(s) 18 - 75 years of age that had a HbA1c test in last 12 reported months.	Endocrinology, Family Medicine, Internal Medicine, Nurse Practitioner, Pediatrics, Physician Assistant, General Practice

*Indicates **HEDIS** measure; **Condition source type key:** 1 = NCQA measure, 2 = Modified NQF measure, 3 = CMS measure, 4 = PQA measure Measures generated with Optum Evidence Based Medicine (EBM) software.

Provider clinical quality

Source type	Condition	Measure description	Specialty list
1	Comprehensive Diabetes Care (CDC)*	Patient(s) 18 - 75 years of age that had an annual screening test for diabetic retinopathy.	Family Medicine, General Practice, Internal Medicine, Nurse Practitioner, Ophthalmology, Pediatrics, Physician Assistant
1	Comprehensive Diabetes Care (CDC)*	Patient(s) 18 - 75 years of age that had annual screening for nephropathy or evidence of nephropathy.	Endocrinology, Family Medicine, Internal Medicine, Nurse Practitioner, Pediatrics, Physician Assistant, General Practice
2	Depression	Patient(s) 18 years of age or older taking a medication for depression treatment that had an annual provider visit.	Family Medicine, Psychiatry and Neurology, Nurse Practitioner, Internal Medicine, Physician Assistant, Mental Health Nurse Practitioner, Pediatrics, General Practice
2	Depression	Patient(s) hospitalized for depression that had a mental health evaluation or a visit with a primary care provider for depression within 30 days of discharge.	Psychiatry and Neurology
2	Depression	Patient(s) hospitalized for depression that had mental health evaluation or visit with a primary care provider for depression within 7 days after discharge.	Psychiatry and Neurology
2	Depression	Patient(s) who are currently taking lithium or an antipsychotic-containing medication that had a psychiatric evaluation in last 6 reported months.	Psychiatry and Neurology
3	Developmental Screening in the First Three Years of Life (National Standard)	Children 1- 3 years of age at the end of the report period that were screened for risk of developmental, behavioral, and social delays using a standardized tool.	Family Medicine, Nurse Practitioner, Pediatrics, Physician Assistant, General Practice

*Indicates HEDIS measure; **Condition source type key:** 1 = NCQA measure, 2 = Modified NQF measure, 3 = CMS measure, 4 = PQA measure Measures generated with Optum Evidence Based Medicine (EBM) software.

Provider clinical quality

Source type	Condition	Measure description	Specialty list
4	Diabetes Medications-Part D Medication Adherence (National Standard)	Patient(s) compliant with all prescribed diabetes medications (minimum compliance 80% or higher).	Family Medicine, Internal Medicine, Nurse Practitioner, Pediatrics, Physician Assistant, General Practice
2	Diabetes Mellitus	Patient(s) compliant with prescribed statin-containing medication (minimum compliance 80%).	Endocrinology, Family Medicine, Internal Medicine, Nurse Practitioner, Physician Assistant, General Practice
1	Diabetes Monitoring for People With Diabetes and Schizophrenia (SMD)*	Patient(s) with schizophrenia or schizoaffective disorder and diabetes who had a HbA1c and LDL-C test during the report period.	Psychiatry and Neurology
1	Diabetes Screening for People With Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications (SSD)*	Patient(s) with schizophrenia, schizoaffective disorder or bipolar disorder taking an antipsychotic medication who were screened for diabetes during the report period.	Psychiatry and Neurology
1	Follow-Up After Emergency Department Visit for Alcohol and Other Drug Abuse or Dependence (FUA)*	Patient(s) 13 years and older with an ED visit for alcohol and other drug abuse or dependence that had a follow-up visit within 30 days.	Psychiatry and Neurology

*Indicates **HEDIS** measure; **Condition source type key:** 1 = NCQA measure, 2 = Modified NQF measure, 3 = CMS measure, 4 = PQA measure Measures generated with Optum Evidence Based Medicine (EBM) software.

Provider clinical quality

Source type	Condition	Measure description	Specialty list
1	Follow-Up After Emergency Department Visit for Alcohol and Other Drug Abuse or Dependence (FUA)*	Patient(s) 13 years and older with an ED visit for alcohol and other drug abuse or dependence that had a follow-up visit within 7 days.	Psychiatry and Neurology
1	Follow-Up After Emergency Department Visit for Mental Illness (FUM)*	Patient(s) six years of age or older with an ED visit for mental illness or intentional self-harm that had a follow-up visit within 30 days.	Psychiatry and Neurology
1	Follow-Up After Emergency Department Visit for Mental Illness (FUM)*	Patient(s) six years of age or older with an ED visit for mental illness or intentional self-harm that had a follow-up visit within 7 days.	Psychiatry and Neurology
1	Follow-Up After Hospitalization for Mental Illness (FUH) - Part 2*	Patient(s) hospitalized for mental illness or intentional self-harm that had a follow-up encounter with a mental health provider within 30 days after discharge.	Mental Health Nurse Practitioner, Psychiatry and Neurology
1	Follow-Up After Hospitalization for Mental Illness (FUH) - Part 2*	Patient(s) hospitalized for mental illness or intentional self-harm that had a follow-up encounter with a mental health provider within 7 days after discharge.	Mental Health Nurse Practitioner, Psychiatry and Neurology
1	Follow-Up Care for Children Prescribed ADHD Medication (ADD)*	Patient(s) with an outpatient, intensive outpatient or partial hospitalization follow-up visit with a prescribing provider during the 30 days after the initial ADHD prescription, AND two follow-up visits during the 31 days through 300 days after the initial ADHD prescription.	Psychiatry and Neurology, Family Medicine, Nurse Practitioner, Pediatrics, Physician Assistant
1	Follow-Up Care for Children Prescribed ADHD Medication (ADD)*	Patient(s) with an outpatient, intensive outpatient or partial hospitalization follow-up visit with a prescribing provider during the 30 days after the initial ADHD prescription.	Physician Assistant, Family Medicine, Nurse Practitioner, Pediatrics

*Indicates **HEDIS** measure; **Condition source type key:** 1 = NCQA measure, 2 = Modified NQF measure, 3 = CMS measure, 4 = PQA measure Measures generated with Optum Evidence Based Medicine (EBM) software.

Provider clinical quality

Source type	Condition	Measure description	Specialty list
2	Heart Failure - Part 1	Patient(s) currently taking an ACE-inhibitor or acceptable alternative.	Cardiovascular Disease, Internal Medicine, Family Medicine, Nurse Practitioner, Physician Assistant
2	Hypertension	Patient(s) compliant with prescribed ACE-inhibitor-containing medication (minimum compliance 80%).	Family Medicine, Internal Medicine, Nurse Practitioner, Physician Assistant, Cardiovascular Disease, Pediatrics, General Practice
2	Hypertension	Patient(s) compliant with prescribed angiotensin receptor blocker (ARB)-containing medication (minimum compliance 80%).	Family Medicine, Internal Medicine, Nurse Practitioner, Physician Assistant, Cardiovascular Disease, Pediatrics, General Practice
2	Hypertension	Patient(s) compliant with prescribed beta-blocker-containing medication (minimum compliance 80%).	Internal Medicine, Family Medicine, Nurse Practitioner, Physician Assistant, Cardiovascular Disease, Pediatrics, General Practice
2	Hypertension	Patient(s) compliant with prescribed diuretic-containing medication (minimum compliance 80%).	Family Medicine, Internal Medicine, Nurse Practitioner, Physician Assistant, Cardiovascular Disease, Pediatrics, General Practice
2	Hypertension	Patient(s) taking an ACE-inhibitor, angiotensin receptor blocker (ARB), diuretic, or aldosterone receptor antagonist-containing medication that had a serum potassium in last 12 reported months.	Family Medicine, General Practice, Internal Medicine, Nurse Practitioner, Pediatrics, Physician Assistant

*Indicates HEDIS measure; **Condition source type key:** 1 = NCQA measure, 2 = Modified NQF measure, 3 = CMS measure, 4 = PQA measure Measures generated with Optum Evidence Based Medicine (EBM) software.

Provider clinical quality

Source type	Condition	Measure description	Specialty list
2	Hypertension	Patient(s) that had a serum creatinine in last 12 reported months.	Physician Assistant, Family Medicine, General Practice, Internal Medicine, Nurse Practitioner, Pediatrics
2	Hypertension	Patient(s) that had an annual physician visit.	Family Medicine, General Practice, Internal Medicine, Nurse Practitioner, Pediatrics, Physician Assistant
1	Immunizations for Adolescents (IMA)*	Patient(s) 13 years old at the end of the report period that had the meningococcal vaccine by their 13th birthday.	Pediatrics, Family Medicine, Nurse Practitioner, Physician Assistant, General Practice
1	Immunizations for Adolescents (IMA)*	Patient(s) 13 years old at the end of the report period that had the Tdap vaccine by their 13th birthday.	Pediatrics, Family Medicine, Nurse Practitioner, Physician Assistant, General Practice
1	Initiation and Engagement of Alcohol and Other Drug Abuse or Dependence Treatment (IET)*	Patient(s) with a new episode of alcohol or drug (AOD) abuse or dependence who initiated treatment and engaged in ongoing treatment within 34 days of the initiation visit.	Psychiatry and Neurology
1	Initiation and Engagement of Alcohol and Other Drug Abuse or Dependence Treatment (IET)*	Patient(s) with a new episode of alcohol or drug (AOD) abuse or dependence who initiated treatment within 14 days of the diagnosis.	Psychiatry and Neurology

*Indicates **HEDIS** measure; **Condition source type key:** 1 = NCQA measure, 2 = Modified NQF measure, 3 = CMS measure, 4 = PQA measure Measures generated with Optum Evidence Based Medicine (EBM) software.

Provider clinical quality

Source type	Condition	Measure description	Specialty list
1	Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM)*	Patient(s) 1 - 17 years who had two or more antipsychotic medications and had blood glucose and cholesterol testing during the report period.	Psychiatry and Neurology
2	Migraine Headache	Adult(s) with an imaging study of the head that was not medically indicated.	Family Medicine, Internal Medicine, Psychiatry and Neurology, Nurse Practitioner, Pediatrics, Physician Assistant, General Practice
2	Migraine Headache	Patient(s) without frequent use of acute medications.	Family Medicine, Psychiatry and Neurology, Nurse Practitioner, Internal Medicine, Physician Assistant, Pediatrics, General Practice
2	Multiple Sclerosis	Patient(s) that had neurology evaluation in last 12 reported months.	Psychiatry and Neurology
1	Non-Recommended Cervical Cancer Screening in Adolescent Females (NCS)*	Patient(s) 16-20 years of age that had a cervical cancer screening (cervical cytology or HPV test) in the last 12 reported months.	Family Medicine, Pediatrics, Nurse Practitioner, Obstetrics and Gynecology, Physician Assistant, Internal Medicine, Midwife, General Practice
2	Obesity and Overweight	Adult(s) with a HDL cholesterol test in last 36 reported months.	Family Medicine, Internal Medicine, Nurse Practitioner, Physician Assistant, General Practice

*Indicates **HEDIS** measure; **Condition source type key:** 1 = NCQA measure, 2 = Modified NQF measure, 3 = CMS measure, 4 = PQA measure Measures generated with Optum Evidence Based Medicine (EBM) software.

Provider clinical quality

Source type	Condition	Measure description	Specialty list
2	Obesity and Overweight	Adult(s) with a LDL cholesterol test in last 36 reported months.	Family Medicine, Internal Medicine, Nurse Practitioner, Physician Assistant, General Practice
2	Obesity and Overweight	Adult(s) with a triglyceride test in last 36 reported months.	Family Medicine, Internal Medicine, Nurse Practitioner, Physician Assistant, General Practice
2	Obesity and Overweight	Patient(s) that had a blood glucose or hemoglobin A1C test in last 36 reported months.	Family Medicine, Internal Medicine, Nurse Practitioner, Physician Assistant, Pediatrics, General Practice
2	Otitis Externa, Acute	Patient(s) 2 years of age and older with acute otitis externa who were NOT prescribed systemic antimicrobial therapy.	Family Medicine, Pediatrics, Nurse Practitioner, Physician Assistant, General Practice
1	Persistence of Beta-Blocker Treatment After a Heart Attack (PBH)*	Patient(s) hospitalized with an acute myocardial infarction (AMI) persistently taking a beta-blocker for six months after discharge.	Cardiovascular Disease, Family Medicine, Internal Medicine, General Practice
1	Pharmacotherapy Management of COPD Exacerbation (PCE)*	Patient(s) 40 years of age and older with COPD exacerbation that received a bronchodilator within 30 days of the hospital or ED discharge.	Family Medicine, Internal Medicine, Nurse Practitioner, Physician Assistant, Pulmonary Disease, General Practice
1	Pharmacotherapy Management of COPD Exacerbation (PCE)*	Patient(s) 40 years of age and older with COPD exacerbation that received a systemic corticosteroid within 14 days of the hospital or ED discharge.	Family Medicine, Internal Medicine, Nurse Practitioner, Physician Assistant, Pulmonary Disease, General Practice

*Indicates HEDIS measure; **Condition source type key:** 1 = NCQA measure, 2 = Modified NQF measure, 3 = CMS measure, 4 = PQA measure Measures generated with Optum Evidence Based Medicine (EBM) software.

Provider clinical quality

Source type	Condition	Measure description	Specialty list
2	Pregnancy Management	Pregnant HIV-positive women that received HIV antiretroviral therapy.	Advanced Practice Midwife, Obstetrics and Gynecology
2	Pregnancy Management	Pregnant women less than 25 years of age that had chlamydia screening.	Advanced Practice Midwife, Family Medicine, Midwife, Obstetrics and Gynecology
2	Pregnancy Management	Pregnant women less than 25 years of age that had gonorrhea screening.	Advanced Practice Midwife, Family Medicine, Midwife, Obstetrics and Gynecology
2	Pregnancy Management	Pregnant women that had HBsAg testing.	Advanced Practice Midwife, Family Medicine, Midwife, Obstetrics and Gynecology
2	Pregnancy Management	Pregnant women that had HIV testing.	Advanced Practice Midwife, Family Medicine, Midwife, Obstetrics and Gynecology
2	Pregnancy Management	Pregnant women that had syphilis screening.	Advanced Practice Midwife, Family Medicine, Midwife, Obstetrics and Gynecology
2	Pregnancy Management	Pregnant women that received Group B Streptococcus testing.	Advanced Practice Midwife, Family Medicine, Midwife, Obstetrics and Gynecology
1	Prenatal and Postpartum Care (PPC)*	Women that received a prenatal visit in the appropriate time period.	Advanced Practice Midwife, Obstetrics and Gynecology, Midwife
1	Prenatal and Postpartum Care (PPC)*	Women that received postpartum care (excluding bundled postpartum services).	Advanced Practice Midwife, Obstetrics and Gynecology, Midwife

*Indicates **HEDIS** measure; **Condition source type key:** 1 = NCQA measure, 2 = Modified NQF measure, 3 = CMS measure, 4 = PQA measure Measures generated with Optum Evidence Based Medicine (EBM) software.

Provider clinical quality

Source type	Condition	Measure description	Specialty list
2	Sinusitis, Acute	Patient(s) that did not have a sinus computerized axial tomography (CT) or magnetic resonance imaging (MRI) test.	Family Medicine, Nurse Practitioner, Physician Assistant, Pediatrics, Internal Medicine, General Practice
2	Sinusitis, Acute	Patient(s) that did not have a sinus radiographic test.	Family Medicine, Nurse Practitioner, Physician Assistant, Pediatrics, Internal Medicine, General Practice
1	Statin Therapy for Patients with Cardiovascular Disease (SPC)*	Patient(s) with cardiovascular disease that received a high-intensity or moderate-intensity statin medication.	Cardiovascular Disease, Family Medicine, Internal Medicine, Nurse Practitioner, Physician Assistant, General Practice
1	Statin Therapy for Patients with Cardiovascular Disease (SPC)*	Patient(s) with statin adherence (proportion of days covered) at least 80% during the treatment period.	Cardiovascular Disease, Family Medicine, Internal Medicine, Nurse Practitioner, Physician Assistant, General Practice
1	Statin Therapy for Patients with Diabetes (SPD)*	Patient(s) 40-75 years with diabetes that received a statin medication.	Cardiovascular Disease, Endocrinology, Family Medicine, Internal Medicine, Nurse Practitioner, Physician Assistant, General Practice
1	Statin Therapy for Patients with Diabetes (SPD)*	Patient(s) with statin adherence (proportion of days covered) at least 80% during the treatment period.	Endocrinology, Family Medicine, Internal Medicine, Nurse Practitioner, Physician Assistant, General Practice

*Indicates HEDIS measure; **Condition source type key:** 1 = NCQA measure, 2 = Modified NQF measure, 3 = CMS measure, 4 = PQA measure Measures generated with Optum Evidence Based Medicine (EBM) software.

Provider clinical quality

Source type	Condition	Measure description	Specialty list
2	Tobacco Use: Screening and Cessation Intervention	Current tobacco users who received medical assistance for tobacco use cessation.	Advanced Practice Midwife, Family Medicine, General Practice, Internal Medicine, Midwife, Nurse Practitioner, Physician Assistant, Pediatrics, Obstetrics and Gynecology
2	Tobacco Use: Screening and Cessation Intervention	Patient(s) who use tobacco.	Family Medicine, Obstetrics and Gynecology, Nurse Practitioner, Internal Medicine, Physician Assistant, Pediatrics, Midwife, General Practice
1	Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics (APP)*	Patient(s) 1 - 17 years of age with a new antipsychotic medication prescription that had psychosocial care as first-line treatment.	Psychiatry and Neurology
1	Use of Imaging Studies for Low Back Pain (LBP)*	Patient(s) with uncomplicated low back pain that did not have imaging studies.	Family Medicine, Internal Medicine, Nurse Practitioner, Physician Assistant, General Practice
1	Use of Spirometry Testing in the Assessment and Diagnosis of COPD (SPR)*	Patient(s) that had appropriate spirometry testing to confirm COPD diagnosis.	Family Medicine, Internal Medicine, Pulmonary Disease, General Practice
1	Well-Child Visits in the First 30 Months of Life (W30) – Part 1*	Patient(s) that had six or more well-child visits with a PCP during the first 15 months of life.	Family Medicine, Nurse Practitioner, Pediatrics, Physician Assistant, General Practice

*Indicates **HEDIS** measure; **Condition source type key:** 1 = NCQA measure, 2 = Modified NQF measure, 3 = CMS measure, 4 = PQA measure Measures generated with Optum Evidence Based Medicine (EBM) software.